



**ANNE CLEMENTS & ASSOCIATES PTY. LIMITED**  
**(ABN 41 077 242 365, ACN 077-160-939)**  
**Environmental and Botanical Consultants**  
**Office 2, 3 Harbourview Crescent, Milsons Point 2060**  
**PO Box 1623, North Sydney 2059**  
**Phone: (02) 9955 9733, Facsimile: (02) 9957 4343**  
**Email: [mail@acabotanic.com](mailto:mail@acabotanic.com)**

13 March 2017

## **Flora and Fauna Report 2016:**

**Six parks in Ryde Local Government Area**  
**Bell Park, Brush Farm Park, Darvall Park, Field of Mars Reserve,**  
**Lambert Park and Outlook Park.**

### **Prepared by:**

Dr AnneMarie Clements  
Tony Rodd  
Jessica Gardner  
Ruth Palsson

### **with fauna section by:**

Damian Lettoof (co-ordinator, reptiles and frog)  
Henry Cook (birds specialist)  
David Martin (nocturnal assistance with surveys)  
Dr James Schlunke (invertebrates specialist)  
Glenn Hoye (bat call analysis)  
Barbara Triggs (mammal hair analysis)

### **Prepared for:**



Lifestyle and opportunity @ your doorstep

City of Ryde – North Ryde Office  
Binary Centre  
Building 0, Level 1  
3 Richardson Place,  
North Ryde NSW 2113  
Phone: 02 9952 8222

## Contents

Preamble.....	6
Section A: Background information, including environmental setting.....	7
A1.0 Introduction.....	8
A1.1 Zoning of the parks.....	8
A2.0 Environmental Setting.....	10
A2.1 Climate.....	10
A2.2 Geology and soil landscape.....	12
A2.2.1 Geology.....	12
A2.2.2 Soil landscape.....	13
A2.2.3 Onsite soil observations.....	14
A2.3 Land use.....	15
A2.3.1 Historic land use.....	15
A2.3.2 Heritage item zoning.....	18
A2.3.3 Land use changes since earliest available aerial photograph.....	18
A2.4 General observations within the parks.....	20
Section B: Flora assessment.....	23
B1.0 Introduction.....	24
B2.0 Existing mappings.....	24
B3.0 The 2016 survey.....	27
B3.1 Methods.....	29
B3.2 Observations/ Findings.....	31
B3.2.2 Exotic and non-local native species.....	35
B3.3 Ground truthing confirmation of the previous mapping.....	38
B3.4 Targeted threatened species searches.....	39
B3.4.1 Database searches.....	39
B3.4.2 Target searches in the parks.....	42
References for Sections A, B.....	43
Section C: Fauna assessment.....	45
C1.0 Introduction.....	46
C1.1 Sampling methods.....	46
C1.1.1 Reptiles.....	46
C1.1.2 Frogs.....	46
C1.1.3 Mammals.....	46
C1.1.4 Birds.....	47
C1.1.5 Fish.....	47
C1.1.6 Invertebrate fauna.....	47
C1.2 Comparisons between the 2006 and 2016 surveys.....	48
C2.0 Fauna findings.....	48
C2.1 Reptiles.....	48
C2.2 Frogs.....	50
C2.3 Mammals.....	52
C2.4 Birds.....	55
C2.5 Fish.....	62
C2.6 Invertebrate fauna.....	63
C2.6.1 Comparability with previous (Biosphere 2006) results.....	63
C2.6.2 Habitat availability in Ryde reserves.....	63

C2.6.2.1 Ground-layer vegetation.....	63
C2.6.2.2 Shrubs and mid-storey vegetation.....	64
C2.6.2.3 Leaf litter and debris.....	64
C2.6.2.4 Aquatic habitats.....	64
C2.6.2.5 <i>Eucalyptus</i> and <i>Acacia</i> spp. foliage.....	64
C2.6.2.6 Tree trunks.....	65
C2.6.3 Invertebrates recorded in the parks.....	65
Reference for Section C.....	87
Appendix C1. Method details.....	89
Acknowledgements.....	91

## Figures

### Site Location

- 1a. Boundaries of the six parks overlaid on the SIX Maps 1:25 000 topographic map
- 1a-1. Boundaries of Bell Park, Brush Farm Park, and Lambert Park overlaid on the SIX Maps 1:25 000 topographic map
- 1a-2. Boundaries of Darvall Park and Outlook Park overlaid on the SIX Maps 1:25 000 topographic map
- 1a-3. Boundary of the Field of Mars Reserve overlaid on the SIX Maps 1:25 000 topographic map
- 1b. Boundaries of the six parks overlaid on the NearMap aerial photograph (dated 19 January 2016)
- 1b-1. Boundaries of Bell Park, Brush Farm Park, and Lambert Park overlaid on the NearMap aerial photograph (dated 19 January 2016)
- 1b-2. Boundaries of Darvall Park and Outlook Park overlaid on the NearMap aerial photograph (dated 19 January 2016)
- 1b-3. Boundary of the Field of Mars Reserve overlaid on the NearMap aerial photograph (dated 19 January 2016)

### Zoning

- 2a. Boundaries of Brush Farm Park, Bell Park and Lambert Park overlaid on the Land Zoning map sheet LZN\_002 (Ryde LEP 2014)
- 2b. Boundaries of Darvall Park and Outlook Park overlaid on the Land Zoning map sheet LZN\_002 (Ryde LEP 2014)
- 2c. Boundary of the Field of Mars Reserve overlaid on the Land Zoning map sheets LZN\_009 and LZN\_010 (Ryde LEP 2014)

### Geology and soils

- 3a-1. Boundaries for Bell Park, Brush Farm Park, Darvall Park, Lambert Park, and Outlook Park overlaid on the Sydney 1:100,000 Geological Map (Herbert and West 1983)
- 3a-2. Boundary for the Field of Mars Reserve overlaid on the Sydney 1:100,000 Geological Map (Herbert and West 1983)
- 3a-3. Boundary for the Field of Mars Reserve overlaid on the Sydney 1:100,000 Geological Map (Herbert and West 1983) with contour lines
- 3a-4. Soil transects overlaid on the Sydney 1:100,000 Geological Map (Herbert and West 1983)
- 3b-1. Boundaries for Bell Park, Brush Farm Park, Darvall Park, Lambert Park, and Outlook Park overlaid on the Sydney 1:100,000 Soil Landscape Map (Chapman et al. 1989)
- 3b-2. Boundary of the Field of Mars Reserve overlaid on the Sydney 1:100,000 Soil Landscape Map (Chapman et al. 1989)
- 3b-3. Soil transects overlaid on the Sydney 1:100,000 Soil Landscape Map (Chapman et al.

1989)

#### **Heritage mapping**

- 4a. Boundaries of Bell Park, Brush Farm Park, Darvall Park, Lambert Park and Outlook Park overlaid on the Heritage Map sheet HER\_002 (Ryde LEP 2014)
- 4b. Boundary of the Field of Mars Reserve overlaid on the Heritage Map sheet HER\_006, HER\_010 (Ryde LEP 2014)

#### **Historic aerial photographs**

- 5a. Boundaries of Bell Park, Brush Farm Park, and Lambert Park overlaid on SIX Maps historical aerial photograph (dated 1943)
- 5b. Boundaries of Darvall Park and Outlook Park overlaid on SIX Maps historical aerial photograph (dated 1943)
- 5c. Boundary of the Field of Mars Reserve overlaid on the SIX Maps historical aerial photograph (dated 1943)

#### **Vegetation mapping**

- 6a-1. Boundaries of Bell Park, Brush Farm Park and Lambert Park, overlaid on the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)
- 6a-2. Boundaries of Darvall Park and Outlook Park, overlaid on the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)
- 6a-3. Boundary of the Field of Mars Reserve overlaid on the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)
- 6b-1. Native vegetation mapping of Brush Farm Park and Lambert Park (Map 5A, Appendix 5, Biosphere 2006)
- 6b-2. Native vegetation mapping of Darvall Park (Map 5B, Appendix 5, Biosphere 2006)
- 6b-3. Native vegetation mapping of the Field of Mars Reserve (Map 5C, Appendix 5, Biosphere 2006)

#### **Surveys**

- 7a. Quadrats surveyed in 2006 and 2016 in Bell Park, Brush Farm Park, and Lambert Park overlaid on the NearMap aerial photograph (dated 19 January 2016)
- 7b. Quadrats surveyed in 2006 and 2016 in Darvall Park and Outlook Park overlaid on the NearMap aerial photograph (dated 19 January 2016)
- 7c. Quadrats surveyed in 2006 and 2016 in the Field of Mars Reserve overlaid on the NearMap aerial photograph (dated 19 January 2016)

#### **Ground truthing**

- 8a. Ground truthing locations surveyed in 2016 in Bell Park, Brush Farm Park, and Lambert Park overlaid on the NearMap aerial photograph (dated 19 January 2016) and the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)
- 8b-1. Ground truthing locations surveyed in 2016 in Darvall Park overlaid on the NearMap aerial photograph (dated 19 January 2016) and the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)
- 8b-2. Ground truthing locations surveyed in 2016 in Outlook Park overlaid on the NearMap aerial photograph (dated 19 January 2016) and the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)
- 8c. Ground truthing locations surveyed in 2016 in the Field of Mars Reserve overlaid on the NearMap aerial photograph (dated 19 January 2016) and the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)

### **Targeted threatened species search**

- 9a. GPS tracking for the targeted threatened species search in the Field of Mars Reserve on the 4 November 2016, overlaid on the NearMap aerial photograph (dated 19 January 2016)
- 9b. GPS tracking for the targeted threatened species search in the Field of Mars Reserve on the 4 November 2016, with ground truthing locations surveyed in 2016, overlaid on the NearMap aerial photograph (dated 19 January 2016)

### **Tables**

- 1. Species recorded at the 11 re-surveyed quadrats, and at the 2 new quadrats, in City of Ryde parks (Bell Park, Brush Farm Park, Darvall Park, Field of Mars Reserve, Lambert Park, Outlook Park)
- 2. Species recorded greater than 2 m in height in the 11 re-surveyed quadrats, and at the 2 new quadrats, in City of Ryde parks (Bell Park, Brush Farm Park, Darvall Park, Field of Mars Reserve, Lambert Park, Outlook Park)
- 3. Braun-Blanquet cover class scores for species recorded at the 11 resurveyed quadrats and at the 2 new quadrats, in City of Ryde parks (Bell Park, Brush Farm Park, Darvall Park, Field of Mars Reserve, Lambert Park, Outlook Park)
- 4a. Species recorded in sub-quadrats and at supplementary sampling locations in Bell Park
- 4b. Species recorded in sub-quadrats and at supplementary sampling locations in Brush Farm Park
- 4c. Species recorded in sub-quadrats and at supplementary sampling locations in Darvall Park
- 4d-1. Species recorded in sub-quadrats and at supplementary sampling locations in Field of Mars Reserve
- 4d-2. Species recorded at supplementary ground truthing quadrats in Field of Mars Reserve
- 4e. Species recorded in sub-quadrats and at supplementary sampling locations in Lambert Park
- 4f. Species recorded in sub-quadrats and at supplementary sampling locations in Outlook Park
- 5. Species recorded greater than 2 m in height in the supplementary sampling locations, including spot locations, ground truthing locations, and ground truthing quadrats, in City of Ryde parks (Bell Park, Brush Farm Park, Darvall Park, Field of Mars Reserve, Lambert Park, Outlook Park)

### **Appendices**

- 1. City of Ryde Land Register (dated 8 July 2014)
- 2. Climatic record at time of the 2006 and 2016 fauna surveys
- 3. Soil report by Dr Pamela Hazelton for soil transects in the Field of Mars Reserve
- 4. Photo file for autumn flora studies 2016
- 5. Statistical report
- 6. Comparison of species composition in 2006 and 2016 recorded in the fixed quadrats

## **Preamble**

This report consists of three sections:

Section A	Background information including the environmental setting.....	7 - 22
Section B	Flora Assessment.....	23 - 44
Section C	Fauna Assessment.....	45 - 90

## **Section A: Background information, including environmental setting**

## A1.0 Introduction

The aim of the current report is to replicate and reconcile the flora and fauna studies of 2006 to 2008 using the same methodology and include a number of new sites. Stage 1 of the City of Ryde Flora and Fauna Studies focuses on the first six sites. The report was prepared at the request of the City of Ryde.

This report assesses the vegetation of these six sites (park locations shown in Figure 1a on the 1:25 000 topographic map and Figure 1b on the aerial photograph). The total area of these six parks is approximately 62 ha (Figures 1a-1, 1a-2, 1a-3 show the park locations on the 1:25 000 topographic map and Figure 1b-1, 1b-2, 1b-3 on the aerial photographs). The parks vary in size from 0.456 ha to 46.7 ha (details in Appendix 1), namely:

Park name	Suburb	Area (m <sup>2</sup> )
Bell Park	West Ryde	4,560
Brush Farm Park	Eastwood	53,614
Darvall Park	West Ryde	60,886
Field of Mars Reserve	North Ryde	467,246
Lambert Park	Eastwood	25,936
Outlook Park	Eastwood	6,978

In addition to required field surveys, the climate differences at time of sampling, zoning, landform, geology and soil landscape and historic land uses of the six parks have been reviewed as part of the environmental setting.

### A1.1 Zoning of the parks

Under the Ryde Local Environment Plan 2014 (LEP 2014) (Figures 2a, 2b, 2c), the zoning of the six parks is as follows:

- All of the areas with tree canopy in the six parks have been zoned E2 Environmental Conservation, and in Brush Farm Park includes an area of SP2 Infrastructure (Classified Road); and
- Four of the six parks include an area of RE1 Public Recreation (not in Lambert and Outlook Parks). The area/s zoned RE1 Public Recreation in three of these four parks is on level fill with batter slopes upslope of the E2 zoning, with:

Park	Fill location
Brush Farm Park	Large lobe protruding from Lawson Street boundary to about 150 m south, with extensive mown areas including playing field, the surrounding batter slopes beneath tree canopy; other batter slopes around park perimeter, e.g. below bowling club.
Darvall Park	Playing fields and sports clubs in south end of park are on fill placed on creek flat, adjoining belt of planted trees also partly on fill. Some fill created by playground construction on north side of Kinson Crescent. Some narrow batter slopes below backyards of Chatham Road properties where they fall into the headwater gully.
Bell Park	Gully with tree canopy in the north partially filled with batter slopes below adjacent roads and residential backyards.
Field of Mars Reserve	Upslope in Cemetery site, with fill in elevated area near south-west edge but also in north-east and south-east edges; extensive batter slopes beneath tree canopy below south-west edge of Cemetery;



Park	Fill location
	fill evident in denuded area south-west of Wellington Road extension and deeper fill with warning signs due to contamination in gully to south-east of denuded area ground. Mown picnic areas on fill placed on flats on both banks of Buffalo Creek in eastern edge of the Reserve.

- All of the six parks have edge (perimeter) impacts, indicated by zoning of land along the perimeters, with length of perimeter adjoining each zone and percentage of total perimeter length as follows:

Park	Zoning on perimeter						
	E2	RE1	RE2	R2	B4	SP1	SP2
Bell Park	10% (105 m)	15% (165 m)	-	75% (830 m)	-	-	-
Brush Farm Park	-	-	15% (180 m)	25% (290 m)	-	-	Road 60% (720 m)
Darvall Park	8% (270 m)	19% (600 m)	3% (100 m)	49% (1580 m)	2% (65 m)	-	Rail 17% (560 m); Telephone 1% (40 m)
Lambert Park	25% (330 m)	-	-	65% (800 m)	-	-	Road 10% (140 m)
Outlook Park	5% (20 m)	-	-	95% (490 m)	-	-	-
Field of Mars Reserve	5% (240 m)	15% (840 m)	-	60% (3040 m)	-	Cemetery 20% (1190 m)	-

**Zoning along park edges:**

E2 Environmental Conservation

RE1 Public Recreation

RE2 Private Recreation

R2 Low Density Residential

B4 Mixed Use

SP1 Special Activities (Cemetery)

SP2 Infrastructure (Classified Road, Railways. Telephone Exchange)

- In the case of Field of Mars Reserve, the Field of Mars Cemetery occupies 20% of its perimeter to the northwest. The access road mapped as Wellington Road and the northwestern ends of Cressy Road and Westminster Road, zoned as RE1 Public Recreation, occupy 15% of the perimeter to the southwest, northwest and southeast respectively. Wellington Road adjoins the cemetery and provides a buffer to nutrient downwash from the cemetery to the northeast. There are some edge effects from the road, with the most obvious being the weedy road batters.

Edge-to-area ratios are high for all parks, except Brush Farm Park and Field of Mars Reserve, namely:

Park name	Edge (m) – E2 Environmental Conservation (m)	Area (m <sup>2</sup> )	Area (ha)	Edge-to-area ratio (m/ha)
Bell Park	1000	4,560	0.46	2,173
Brush Farm Park	1,180	81,600	8.16	145
Darvall Park	2,940	90,930	9.09	323
Field of Mars Reserve	5,070	467,246	46.73	109

Park name	Edge (m) – E2 Environmental Conservation (m)	Area (m <sup>2</sup> )	Area (ha)	Edge-to-area ratio (m/ha)
Lambert Park	950	25,936	2.59	367
Outlook Park	490	6,978	0.70	700

## A2.0 Environmental Setting

Ryde Local Government Area (LGA) is mapped on the Parramatta River 1:25,000 topographic map (Figure 1a). Ryde LGA is within the Sydney Basin Bioregion.

The landform of most of the parks includes gullies and adjoining slopes that historically have been too steep for low-cost residential development (Figures 1a-1 to 1a-3, 2a, 2b, 2c).

There are two mapped creeks, Buffalo Creek and Strangers Creek, in the Field of Mars Reserve and unmapped ephemeral watercourses in the gullies of the parks (Figures 1a-1 to 1a-3). The unmapped ephemeral watercourses in Brush Farm Park, Bell Park and Lambert Park are headwaters of Archer Creek, while the one in Darvall Park is a headwater of Smalls Creek.

In many of the parks there has been dumping of fill in earlier years, often followed by creation of playing fields and other recreation spaces on the resulting levelled surfaces. Batter slopes on the fill edges have spilled down into adjoining gullies beneath existing tree canopy, or the canopy of subsequent tree plantings.

### A2.1 Climate

The rainfall records from two nearby meteorological stations, Macquarie Park (Willandra Village, Station No. 66156) and Parramatta North (Masons Drive, Station No. 66124) (Bureau of Meteorology website, [www.bom.gov.au](http://www.bom.gov.au), accessed November 2016), are consistent with the general trend of decreasing rainfall with increasing distance from the coast. The mean annual rainfall at Parramatta North is lower than that at Macquarie Park, namely:

Station	Distance inland	Mean annual rainfall	Opened	Operational
Macquarie Park	17 km	1156 mm	1970	45 years
Parramatta North	25 km	971 mm	1965	50 years

From the climate statistics, the lowest monthly rainfall recorded at both stations was less than 10 mm and highest monthly rainfall greater than 200 mm, hence expected drought stress during low rain and potential erosion risk during high rain events.

#### Rainfall data for Macquarie Park (Willandra Village)

Statistic	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
Mean	126	140	129	114	83	121	56	61	60	84	93	88	1156
Lowest	6	17	14	7	2	3	1	0	0.2	0.4	7	11	639
Highest	370	655	339	562	345	430	215	399	219	306	356	274	2011

#### Rainfall data for Parramatta North (Masons Drive)

Statistic	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
Mean	106	121	107	94	70	91	46	57	53	68	87	74	971
Lowest	7	9	8	3	2	5	0	0	0.4	0.2	8	6	513.2
Highest	319	673	311	476	250	374	166	440	324	237	236	271	1713

Prior to and at the times of flora and fauna surveys (Autumn and Spring 2006 and Autumn and Spring 2016), there was:

Time of survey	Rainfall condition
Autumn 2006	More than 20 mm below monthly mean
Spring 2006	Variable
Autumn 2016	More than 20 mm below monthly mean
Spring 2006	Variable

#### Monthly rainfall data recorded at Macquarie Park (Willandra Village)

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec	Annual
2005	88	104		29	41	87		3		53	141		
2006	95	67	36	7	15	127	75		219	10	63	74	

2015	189	52	78	362	118	69	33	42	55	54	132	69	1253
2016	370	40	84	33	7	346	102	149	60	28			
Mean	126	140	129	114	83	121	56	61	60	84	93	88	1156

#### Monthly rainfall data recorded at Parramatta North (Masons Drive)

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec	Annual
2005	89	141	69	25	36	74	34	3	51	54	146	24	746
2006	110	68	33	4	12	108	64	48	192	8	46	67	761

2015	139	62	41	407	81	60	38	40	34	46	121	65	1133
2016	319	9	47	31	10	324	84	112	56	19			
Mean	106	121	107	94	70	91	46	57	53	68	87	74	971

	> 20 mm below the monthly average
	Within 20 mm of the monthly average
	> 20 mm above the monthly average

The 2006 autumn fauna surveys were undertaken by Biosphere Environmental Consultants on 26 to 30 April and 1, 4-9, 11 to 17 May 2006. The 2016 autumn fauna surveys were undertaken on 4 to 8, 11 to 15, 25 to 29 April and 2 May 2016. The daily temperatures and rainfall during the 2016 fauna surveys were higher than during the 2006 fauna surveys (Appendix 2), with:

#### Climatic conditions recorded during Autumn surveys

Records from Parramatta North (Masons Drive) meteorological station	26 April to 17 May 2006	4 April and 2 May 2016
<b>Temperature</b>		
Daily minimum range	4.6 to 12.4°C	9 to 19°C
Daily maximum range	17.8 to 26.0°C	20 to 35.2
<b>Rainfall</b>		
Mean	0.3 mm	0.8 mm
Range	0 to 2.4 mm	0 to 9.4 mm

The 2006 spring fauna surveys were undertaken on 9 to 14, 19 to 28 October and 1 to 6 November, and the 2016 autumn fauna surveys were undertaken on 19 to 23, 25 to 29 September. The daily temperatures during the 2016 fauna surveys were lower than during the 2006 fauna surveys, and the average daily rainfall similar in 2006 and 2016 (Appendix 2), with:

### Climatic conditions recorded during Spring surveys

Records from Parramatta North (Masons Drive) meteorological station	9 October and 6 November 2006	19 and 29 September 2016
<b>Temperature</b>		
Daily minimum range	6.2 to 23.0°C	5.4 to 14.2°C
Daily maximum range	16.0 to 37.2°C	20.0 to 23.0
<b>Rainfall</b>		
Mean	1.8 mm	1.9 mm
Range	0 to 25.5 mm	0 to 15 mm

## A2.2 Geology and soil landscape

### A2.2.1 Geology

The geology of the Sydney 1:100 000 map sheet was mapped by Herbert and West (1983) (Figure 3a-1), with:

#### Geological mapping of the parks

Park	Quaternary Alluvial Deposits (map unit Qha)	Hawkesbury Sandstone (map unit Rh)	Wianamatta Group Shales	
			Ashfield Shale (map unit Rwa)	Bringelly Shale (map unit Rwb)
Bell Park			X	
Brush Farm Park			X	X
Darvall Park			X	
Field of Mars Reserve (see Figure 3a-2, 3a-3)	Below 10-20 m AHD	Slopes below approx. 40 m AHD and above 10-20 m AHD	Above approx. 40 m AHD	
Lambert Park			X	
Outlook Park			X	

The geological map units are described by Herbert and West (1983) as follows:

#### Wianamatta Shales Group

- Ashfield Shale (Rwa): *Black to dark grey shale and laminate;*
- Bringelly Shale (Rwb): *Shale, carbonaceous claystone, laminate, fine to medium-grained lithic sandstone, rare coal;*

#### Hawkesbury

- Hawkesbury Sandstone (Rh): *Medium to coarse grained quartz sandstone, very minor shale and laminate lenses; and*

#### Quaternary Deposits

- Quaternary Holocene Alluvium (Qha): *Silty to peaty quartz sand, silt, and clay. Ferruginous and humic cementation in places. Common shell layers.*

## A2.2.2 Soil landscape

The soil landscapes of the Sydney 1:100 000 map sheet were mapped by Chapman et al. (1989). The soil landscapes vary in topographic position and parent material (Chapman and Murphy 1989), with:

Soil landscape	Parent material	Landform
Glenorie (gn)	Wianamatta group shales	Undulating to rolling low hills
West Pennant Hills (wp)	Wianamatta group shales and shale colluvium (shale downwash)	Rolling to steep side slopes
Lucas Heights (lh)	Mittagong Formation (alternating bands of shale and fine-grained sandstones)	Gently undulating crests and ridges on plateau surfaces
GyMEA (gy)	Hawkesbury Sandstone (underlying the Wianamatta group shales)	Undulating to rolling rises and low hills
Hawkesbury (ha)	Hawkesbury Sandstone	Rugged, rolling to very steep hills
Lane Cove (lc)	Alluvial floodplain draining both the Wianamatta Group shales and Hawkesbury Sandstone	Level to gently undulating

**Disturbed terrain (xx)** occurs on *level plain to hummocky terrain, extensively disturbed by human activity, including complete disturbance, removal or burial of soil.*

The parks were mapped (Figures 3b-1, 3b-2) as follows:

Park/ Reserve	Soil landscape map units
Bell Park	Glenorie (gn) on gentle lower slopes, West Pennant Hills (wp) on steeper upper slopes includes the gully
Brush Farm Park	Glenorie (gn) on gentle upper slopes, West Pennant Hills (wp) on steeper slopes includes the gully
Darvall Park	Glenorie (gn) on the gentle slope of most of the park West Pennant Hills (wp) in the upper northern tip Central gully through the park
Field of Mars Reserve	Disturbed terrain (xx) associated with the Field of Mars Cemetery and extending into the reserve Lucas Heights (lh) in northern tip of the reserve GyMEA (gy) in the north-west corner of the reserve Hawkesbury (ha) on steep slopes Lane Cove (lc) downslope of slopes
Lambert Park	Glenorie (gn) on gentle lower slopes, West Pennant Hills (wp) on steeper upper slopes includes the gully
Outlook Park	West Pennant Hills (wp) on slopes includes the gully

Most of the soil landscapes in the parks have erosion hazards (Chapman and Murphy 1989). The limitations for Disturbed terrain (xx) are dependant upon the nature of the fill material. Erosion hazards are exacerbated by high rainfall events. The limitations of the soil landscapes are as follows:

<b>Soil landscape</b>	<b>Hazard</b>
Glenorie (gn)	<i>High soil erosion hazard, localised impermeable highly plastic subsoil, moderately reactive. High soil erosion.</i>
GyMEA (gy)	<i>Localised steep slopes, high soil erosion hazard, rock outcrop, shallow highly permeable soil, very low soil fertility.</i>
Hawkesbury (ha)	<i>Extreme soil erosion hazard, mass movement (rock fall) hazard, steep slopes, rock outcrop, shallow, stony, highly permeable soil, low soil fertility.</i>
Lane Cove (lc)	<i>Flooding, high soil erosion hazard, seasonal waterlogging.</i>
Lucas Heights (lh)	<i>Stony soil, low soil fertility, low available water capacity.</i>
West Pennant Hills (wp)	<i>Mass movement hazard, steep slopes, high soil erosion hazard, localised seasonal waterlogging, impermeable plastic shrink-swell subsoil.</i>
Disturbed terrain (xx)	<i>Dependant on nature of fill material. Mass movement hazard, unconsolidated low wet-strength materials, impermeable soil, poor drainage, localised very low fertility and toxic materials.</i>

### **A2.2.3 Onsite soil observations**

There were clay soils observed in all parks, except for the Field of Mars Reserve. The observed soils are consistent with the geology and soil landscape mappings (Figures 3a-1, 3a-2, 3b-1, 3b-2).

In the Field of Mars Reserve, the soil landscape boundaries were investigated by sampling soils using a 30 mm stainless steel auger to a depth of approximately 300 mm at 10 m intervals along four transects at right angles to the contours and at two additional spot locations GTb, GTc (Figures 3a-4, 3b-3). The soil samples from the auger were separated by colour and depth recorded. The soil samples were analysed by the soil scientist, Dr Pamela Hazelton (recording in Appendix 3).

#### **Soil findings of Dr Pamela Hazelton at the Field of Mars Reserve**

<b>Soil sample</b>	<b>Geology mapping</b>	<b>Soil landscape</b>	<b>Observed</b>
<b>Transect 1</b>			
0 m	Ashfield Shale	Disturbed	Ashfield Shale
10 m	Hawkesbury Sandstone	Disturbed	Ashfield Shale
20 m	Hawkesbury Sandstone	Disturbed	Ashfield Shale with Hawkesbury Sandstone
30 m	Hawkesbury Sandstone	Disturbed	Ashfield Shale with Hawkesbury Sandstone
40 m	Hawkesbury Sandstone	Disturbed	Hawkesbury Sandstone
<b>Transect 2</b>			
0 m	Ashfield Shale	Disturbed	Disturbed
10 m	Hawkesbury Sandstone	Disturbed	Disturbed
20 m	Hawkesbury Sandstone	Disturbed	Disturbed
30 m	Hawkesbury Sandstone	Disturbed	Disturbed
40 m	Hawkesbury Sandstone	Disturbed	Disturbed
<b>Transect 3</b>			
0 m	Ashfield Shale	Disturbed	Hawkesbury Sandstone derived
10 m	Ashfield Shale	Disturbed	Hawkesbury Sandstone derived
20 m	Ashfield Shale	Disturbed	Hawkesbury Sandstone derived

Soil sample	Geology mapping	Soil landscape	Observed
30 m	Ashfield Shale	Disturbed	Hawkesbury Sandstone derived
40 m	Ashfield Shale	Disturbed	Hawkesbury Sandstone derived
<b>Transect 4</b>			
0 m	Ashfield Shale	Lane Cove	Minimal topsoil, shale fragments
10 m	Ashfield Shale	Lane Cove	Hawkesbury Sandstone derived
20 m	Ashfield Shale	Lane Cove	Hawkesbury Sandstone derived
30 m	Ashfield Shale	Lane Cove	Hawkesbury Sandstone rock
40 m	Hawkesbury Sandstone	Lane Cove	Hawkesbury Sandstone derived
50 m	Hawkesbury Sandstone	Lane Cove	Hawkesbury Sandstone rock
<b>GTb</b>			
	Hawkesbury Sandstone	Lane Cove	Disturbed
<b>GTc</b>			
	Ashfield Shale	Disturbed	Disturbed. Subsoil with no topsoil

The mapping of Lane Cove soil landscape in Transect 4 and Spot location GTb is unlikely to be correct as the landform for the Lane Cove soil landscape for this soil is *level to gentle undulating Alluvial floodplain*. Transect 4 and Spot location GTb were on a hillslope. The likely soil landscape is either Hawkesbury (map unit ha) which occurs on *rugged to very steep on Hawkesbury Sandstone*, or possibly Gynea soil landscape.

## A2.3 Land use

### A2.3.1 Historic land use

The Ryde LGA is the traditional land of the Wallumedegal people (Smith 2005).

The land uses of Brush Farm Park, Darvall Park, Lambert Park and the Field of Mars Reserve are as follows.

**Brush Farm Park** was formally established as a park in 1914. The original land grants that covered the park area date from 1794. The park is part of the former "Brush Farm Estate". From 1806, the land was being used for cattle and viticulture.

In 1834 Baron Charles von Hugel (Clark 1994) concerning the vegetation of Brush Farm, noted: *Near the house, there is a deep valley with a type of vegetation all its own, containing a number of plants which I found only in the Illawarra.*

Benson and Howell (1990, p124, 125) described the vegetation of Brush Farm Park as: *Remnants of the higher rainfall Blue Gum High Forest can be seen in Darvall Park in Denistone and in Brush Farm Park at Eastwood. These are situated in parts of the municipality where the shale soils are deep... At Brush Farm Park, similar trees grow on the upper slope [as at Darvall Park], but in the steep-sided sheltered gully, fertile shale-derived and high rainfall support a rainforest vegetation with species not found together in other sheltered sandstone gullies or on Wianamatta Shale soils of northern Sydney. Named Brush Farm by early settlers because of this rainforest brush its species included trees of Cryptocarya glaucescens, Euodia micrococca, Guioa semiglauc, Schizomeria ovata, shrubs of Alectryon subcinereus and Eupomatia laurina, and the*

*climber Aphanopetalum resinosum. There is a very large Trochocarpa laurina 12 m high. The moist fertile gully has been particularly susceptible to weed invasion.*

### **Darvall Park**

A comprehensive account of this park's history is provided in the State Heritage Register website at <http://www.environment.nsw.gov.au/heritageapp/ViewHeritageItemDetails.aspx?ID=2340177>

The park represents a residue of the 19th-century Darvall Estate after nearly all the estate was subdivided and sold in the early 20th century. The park land was acquired by Ryde Council between 1926 and 1929. Its listing on the State Heritage Register was mainly on account of its historical association with Major Edward Darvall, also because there is preserved in it the endangered ecological community Turpentine-Ironbark Forest [in the Sydney Basin Bioregion].

In the Friends of Darvall Park website (<http://friendsofdarvallpark.weebly.com/park-history.html>, accessed 9 November 2016), it is stated that:

*In the late 1850s Edward Darvall built Ryedale house and surrounded it with 20 ha of orange orchards. The rest of the land was used for growing vegetables, some cereals, corn, potatoes and steeper sections were used as pastures.*

*Major Darvall died in 1869. ... The Ryedale estate was left to Jane [his wife] who remained at the house till her death in 1899. ... she sold parcels of land to the Commissioner of railways in 1885, Eastwood public school in 1889 and the West Ryde Masonic Temple in 1898. Just before her death she made a gift of Ryedale house and the remaining land to her son Anthony William Darvall.*

*Although Darvall Park was known as a park since 1923, not until 1936 the council could be persuaded to buy the 18 acres, which at first it thought of calling "Kings Park." ... It was partly bush partly open space. The lower flatter section was and still is used for sport and recreation. At the time of inception it was considered to be sports ground for girls.*

Benson and Howell (1990, p124) described the vegetation of Darvall Park as:

*Remnants of the higher rainfall Blue Gum High Forest can be seen in Darvall Park in Denistone and in Brush Farm Park at Eastwood. These are situated in parts of the municipality where the shale soils are deep. At Darvall Park near Denistone station, there are tall trees of smooth-barked Eucalyptus saligna, along with the rough-barked Eucalyptus pilularis, Eucalyptus paniculata, Eucalyptus resinifera, Eucalyptus acmenoides (near its southern limit here) and Syncarpia glomulifera. Much of the understorey has been replaced with grass patches, but amongst the woody shrubs and trees (Ligustrum, Lantana, Salix, Erythrina and Cinnamomum), many native understorey species are present, and their growth is being encouraged in areas by log barriers from mowing and trampling. Small trees of Backhousia myrtifolia and Glochidion ferdinandi, shrubs of Bursaria spinosa, Helichrysum diosmifolium [now Ozothamnus diosmifolius], at least four species of Acacia, and moisture-loving vines including Morinda jasminoides and Celastrus subspicatus, Cissus antarctica and Cissus hypoglauca are amongst the native plants present in these regenerating areas.*

### **Lambert Park**

From the Brush Farm Park and Lambert Park Plan of Management (City of Ryde, 2009):



*Brush Farm Park has been established since 1914 and Lambert Park since 1984. ... Significant works have been undertaken within both Parks including bush regeneration, sports ground maintenance and creek line rehabilitation works. ...*

*Lambert Park forms a continuation of the natural area from Brush Farm Park. ...*

*Lambert Park is characterised in the western upper area by fill material with a crushed sandstone capping and revegetation. There are small pockets of local vegetation on the upper slopes and in the southern area where the original soil layer still persists. ...*

*Both Parks are important historically as they formed part of the Brush Farm House Estate with Brush Farm House immediately to the north of Brush Farm Park. The earliest European occupancy of the land dates back to 1794.*

### **Field of Mars Reserve:**

From the Plan of Management (City of Ryde 2009), it is stated that:

*The Field of Mars is the largest reserve under the care, control and management of the City of Ryde. The reserve is dedicated for public recreation and promotion of the study and the preservation of native flora and fauna and accordingly it is used for environmental education and passive recreation, involving walking and appreciation of the natural environment. This has been the major focus of management for recreation within the Reserve since the 1960's.*

The history of the reserve is summarised as follows (City of Ryde 2009):

Prior to European settlement, the area was occupied by the Wallumedegal people, with:

- creeks and estuary providing a major food source;
- moist gullies providing fruits such as figs, lillypilly and berries;
- drier areas providing nectar from flowers such as banksia and waratah; and small animals such as bandicoots, bush rats and possums;
- tall forests of the shale ridges with their grassy and bracken understorey were areas which attracted larger mammals such as kangaroos and wallabies.

The first land grant in the Ryde area was along the northern bank of the river between Sydney and Parramatta in January 1792. The area was named by Governor Phillip the 'Field of Mars', because part of it was reserved for use as an army encampment.

In 1804 a large area of public land for use by local inhabitants was set aside. Six commons were gazetted, including The Field of Mars Common, an area of approximately 5,050 acres located north of the Field of Mars and the Eastern Farms, and which covered most of the Ryde Municipality.

In 1874, the Common, by then a reported place of undesirables, was resumed as Crown land and subsequently cleared for the laying out of allotments and streets.

In 1884 25 acres were allocated for the Field of Mars Cemetery.

In the Government Gazette of 3 December 1887, 85 acres were proclaimed for Public recreation. This is the origin of today's Field of Mars Reserve. Ryde Council became trustee of the land in 1889.

The reserve remained a source of building and domestic materials for local residents, provided grazing land for cattle, a refuge during the depression years and a popular place for swimming, fishing and prawning until 1954.

In response to a waste disposal problem as residential development grew, the Field of Mars was identified as a location for a major putrescible waste tip in 1965. This was to prove the catalyst for significant resident mobilisation in relation to wildlife conservation.

The Ryde-Hunter’s Hill Flora and Fauna Preservation Society was formed in February 1966 to advocate wildlife conservation and to specifically preserve, manage and develop the Field of Mars Reserve as a flora and fauna sanctuary.

On 9 May 1975, the reserve was proclaimed a “Wildlife Refuge” under the National Parks and Wildlife Act, 1974. (*Field of Mars Wildlife Refuge No.339*).

### **A2.3.2 Heritage item zoning**

Under Part 1 of Schedule 5 of the Ryde Local Environment Plan 2014 (LEP 2014), three of the six parks are listed as heritage items of local significance, and are mapped as follows:

- Brush Farm Park: ‘C1’ Conservation Area – General, and Item – General (Figure 4a);
- Darvall Park: Item – General (Figure 4a); and
- The Field of Mars Reserve: Item – General (Figure 4b).

Heritage conservation area is defined in the Ryde LEP (2014) as follows:

*... an area of land of heritage significance:*

*(a) shown on the Heritage Map as a heritage conservation area, and*

*(b) the location and nature of which is described in Schedule 5, and includes any heritage items situated on or within that area.*

A heritage item is defined in the Ryde LEP (2014) as follows:

*... a building, work, place, relic, tree, object or archaeological site the location and nature of which is described in Schedule 5.*

Heritage significance is defined in the Ryde LEP (2014) as follows:

*... historical, scientific, cultural, social, archaeological, architectural, natural or aesthetic value.*

### **A2.3.3 Land use changes since earliest available aerial photograph**

From a comparison with the 1943 aerial photographs (Figures 5a, 5b, 5c) with the 2016 aerial photographs (Figures 1b-1, 1b-2, 1b-3), it was observed that:

<b>Park</b>	<b>Land use in 1943</b>	<b>Land use in 2016</b>
Bell Park	Cleared with trees along Winbourne Street to the east, in rural environment.	Area of canopy trees in north and scattered trees in southern 2/3, surrounded by residential.
Brush Farm Park	Rural cleared land upslope with scattered canopy trees in gullies and on the perimeter.  Urban development to the east and	Canopy trees throughout, except raised recreational grounds to the northwest and in the central north, as well as a carpark and Scout Hall.

Park	Land use in 1943	Land use in 2016
	<p>rural to north and west.</p> <p>Canopy in the gully to the south separating the park from Lambert Park.</p>	<p>Surrounded by urban development, with canopy to the south separating the park from Lambert Park.</p>
Darvall Park	<p>Cleared along boundary with railway, and in southern half, with canopy coverage in northern half and scattered trees in south.</p>	<p>Canopy trees, except in mown recreation field in the south.</p>
Field of Mars Reserve	<p>Fully vegetated, except for:</p> <ul style="list-style-type: none"> <li>- the cleared agricultural land on the alluvial flats adjoining Buffalo Creek;</li> <li>- the cleared upper slopes in the south and south-west.</li> </ul> <p>There are tracks through the canopy vegetation, providing access to the alluvial flats.</p> <p>There is a narrow band of vegetation along Buffalo Creek.</p> <p>Surrounded to northwest and northeast by vegetation, to the southwest and north by rural land use, and to the south by urban development.</p> <p>Field of Mars Cemetery was established in central area to north west of the Reserve.</p>	<p>Remains fully vegetated, with:</p> <ul style="list-style-type: none"> <li>- scattered trees and mown grass on the formerly cleared agricultural land on the alluvial flats adjoining Buffalo Creek;</li> <li>- grass growth and rehabilitation being undertaken on the formerly cleared upper slopes in the south and southwest.</li> </ul> <p>Track less visible than in 1943.</p> <p>Growth along the creekline.</p> <p>Dense canopy coverage with scattered clearing to southeast and at the end of Wellington Road. Surrounded by urban development with the Lane Cove National Park to the east and Pidding Park and canopy trees to the northwest.</p> <p>Field of Mars Cemetery expanded in central area to north west, with graves and tombs extending on level fill to boundary with the Reserve.</p>
Lambert Park	<p>Cleared in the southern third with canopy cover in the northern two thirds. Dense canopy along the gully.</p> <p>Surrounded by rural to the south, west with urban development to the east and by canopy cover to the north separating the park from Brush Farm Park.</p>	<p>Similar to 1943, except the cleared southern third now supporting canopy cover.</p> <p>The former rural to the south, west now urban development.</p> <p>The gully appears to have been filled.</p>
Outlook Park	<p>Canopy cover throughout and surrounded by urban development with scattered canopy trees.</p>	<p>Similar to 1943.</p>

## A2.4 General observations within the parks

**Bell Park** slopes to the south, with a children's playground and an old fig tree (*Ficus rubiginosa*) in the southernmost section. The vegetation of Bell Park consists mostly of mown grass recreational area, with dense belts of planted and weedy tree and shrub vegetation following the line of a gully.

**Brush Farm Park** slopes to the southeast, and includes a mown oval on the central ridge extending into areas of more roughly mown grass to the south. The north-western edge of the park is occupied by a car park and netball courts. The vegetation of this park consists of native bushland except for some large mown or paved areas fronting onto Lawson Street, including the Scout Hall and its grounds.

A broad belt of bushland occupies the long gully head between the oval and Brush Road. The south-flowing gully joins another gully from the northwest, with their confluence being outside the park's southern boundary.

The broad gully to the northwest is densely forested, with large, mature eucalypts dominating the south-facing slopes behind the Scout Hall and netball courts, most of them *E. paniculata* (Grey Ironbark). Deeper in the gully the forest becomes more mesic, approaching a true rainforest except for the persistence of emergent mature eucalypts. Towards the southwest on the slope running up to the bowling club there is disturbance possibly from deposition of fill and mature canopy trees are absent; rather, there are stands of even-aged eucalypts that appear to have been planted in recent decades. Examination of fallen fruits revealed that all or most of the planted eucalypts are *E. grandis* (Flooded Gum), a North Coast species of very similar appearance to *E. saligna* and often confused with it.

**Darvall Park** occupies the head of a gully running south from near Eastwood station on the west side of the Main Northern Railway. It is irregular in shape, with the northern half tapering into a point where Chatham Road converges with the railway. It is broadest in the middle, south from which it diverges from the railway with housing around Kinson Crescent and Anthony Road intervening. Further south it narrows to a strip along the creek, which in this part is low-lying and swampy. The final 500 m to the southern extremity is largely cleared with mown grass for sporting activities, though with an interrupted fringe of trees in parts. The highest-quality bushland is concentrated in the central, broadest part and features quite a number of impressively large and mature eucalypts, many with hollow branch stubs currently sheltering a variety of birds including king parrots, corellas and white cockatoos. Most frequent of the large eucalypts is *E. saligna* (Sydney Blue Gum) followed by *E. acmenoides* (White Mahogany) and *E. paniculata* (Grey Ironbark). Further north and south much of the forest is degraded and very weedy, with large trees much sparser. The low-lying parts of the creek support large swathes of wetland weeds, with massive weed growth extending up the banks. At both northern and southern ends of the bushland area there is evidence of tree planting in recent decades, dominated by eucalypt species inappropriate to the Sydney region, namely *Eucalyptus grandis* (Flooded Gum) and *E. microcorys* (Tallowwood).

### Field of Mars Reserve

The bushland, now Field of Mars Reserve, wraps around Field of Mars Cemetery on three sides. The cemetery occupies the broad flat end of the ridge system, further flattened by deposition of fill in parts, with the reserve falling away from its boundaries into gullies. There are two creeks in the reserve, Strangers Creek in the north-east and Buffalo Creek in the south.

By far the largest part of the reserve is on Hawkesbury sandstone with many low rock ledges and boulders outcropping, exceptions being some narrow fringes adjacent to the cemetery which are on sediments transitional between shale and sandstone, and alluvial deposits along the lower reaches of Buffalo Creek and Strangers Creek.

The vegetation of the reserve is largely remnant and in many parts in almost pristine condition. However the creeklines are generally weedy due to eutrophication from neighbouring residential areas upslope and to a lesser degree from the cemetery.

Southwest of the cemetery on the slope between Wellington Road and Buffalo Creek, there are some very weedy areas of bushland, in particular in a gully below the end of Wellington Road. The parcels of land at 2-14 Wellington Road have been used for a variety of residential, commercial and industrial purposes since the 1920s. A former electrical factory built in the 1960s was occupied by Council from 1975 as their works depot. Remediation works to deal with soil contamination at the site included removal of primary contamination sources, capping with clean soil and rock, and revegetation in 2009. Past land use and disturbance has led to the weed infestations downslope from this area. (See details at <http://www.ryde.nsw.gov.au/Environment-and-Waste/Remediation-Projects>). The revegetation plantings in this area are an assortment of native trees and shrubs, many of them not local native species. Past land use and disturbance has led to weed infestations downslope from this area. Areas in the south-east of the reserve have been slashed and mown

Apart from the widespread sandstone slopes forest/woodland communities characteristic of Hawkesbury Sandstone, there are narrow zones of several other vegetation communities present, namely:

- mangroves and saltmarsh along the lowest part of Buffalo Creek in the eastern corner; representing the upstream limit of these communities which are much more extensive closer to Lane Cove River on the opposite side of Pittwater Road;
- warm-temperate rainforest dominated by *Ceratopetalum apetalum* (Coachwood) and *Tristaniopsis laurina* (Water Gum) in a very narrow zone along rocky bank of Buffalo Creek, a short distance upstream from the limit of its alluvial flat; also on Strangers Creek though there Coachwood and Red Cedar appear possibly to have been planted;
- a narrow fringe of forest on transitional soils along Wellington Road, dominated by *Eucalyptus resinifera*, *E. paniculata* and *Syncarpia glomulifera*; and
- riparian vegetation along Buffalo Creek beyond upstream limit of mangroves and related salinity tolerant vegetation, but below the first rock bars; this vegetation is exceptionally weedy but appears to have included paperbark *Melaleuca* spp. and *Casuarina glauca*.

As with most such bushland reserves surrounded by residential development, the creeklines have had sewer mains constructed in or beside their beds in the mid-20th century. The sewers have contributed greatly to weed invasion.

### **Lambert Park**

The park is effectively a southern extension of Brush Farm Park running downstream along the same gully. Technically the two parks are not contiguous, being separated by a road reserve (a never-built western extension of Rutledge Street) but the road reserve is managed by Council as part of its bushland. The gully is not very steep-sided except toward the northern end (the road reserve) where the creekline is more deeply incised into the broader valley profile. Like Brush Farm Park, Lambert Park is located entirely on Wianamatta Group shales, which have given rise to gentler slopes than the Hawkesbury Sandstone.

An unmapped creek, apparently semi-permanent with shallow pools, runs through the lower part of the park. It is one of the headwaters of Archer Creek which joins Parramatta River at Meadowbank. There is a Girl Guides building in the edge of the park about 50 metres north-west of the junction of Brush Road and Warrawong Street. South of this point, the park flattens out and the creek winds between tall trees, nearly all of which appear to be of planted origin, dating from within about the last 40 years judging by their trunk diameters. Around the west and north sides of the Girl Guides building there is a patch of remnant rainforest, dominated by mature trees of *Acmena smithii* (Lilly-pilly) and some large (but not necessarily old) trees of *Ficus rubiginosa* (Port Jackson Fig). Some smaller local rainforest trees, shrubs and climbers survive despite the moderately abundant weeds here. North from the Girl Guides building and up to the road reserve most of the vegetation is highly disturbed and trees more widely spaced, nearly all apparently of planted origin; the groundlayer is mostly weed species and there are various exotic shrubs, small trees and climbers closer to the creekline.

### **Outlook Park**

This small park, descends steeply from an entrance on Trelawney Street, then broadens in the centre into a shallow gully flanked by a flatter bench, narrowing again to an entrance on Chatham Road. Apart from the road entrances it is bordered entirely by private house blocks, though many of these have large gardens containing remnant trees. A staircase descends through the steep upper section.

Remnant native vegetation survives mainly in the form of some very large, mature trees of *Eucalyptus acmenoides* (White Mahogany), *E. saligna* (Sydney Blue Gum) and *Syncarpia glomulifera* (Turpentine). The occurrence of *Eucalyptus acmenoides* here (and in nearby Darvall Park), represents the species virtually at its southern limit of natural occurrence. The groundlayer is mown in parts though with the native grass *Microlaena stipoides* abundant, but above the central bench there is a large patch of tall, dense exotic weeds, notably *Hedychium gardnerianum* (Yellow Ginger).

## **Section B: Flora assessment**

## B1.0 Introduction

The main aim of this 2016 survey is to replicate and reconcile previous studies using the same methodologies, and to undertake ground truthing of vegetation communities through on-site analysis.

The specific methods include:

- Survey in Autumn and Spring 2016 using the same methodology and quadrats set up for the previous studies [Biosphere 2006].
- Representative of mapped vegetation communities and using the same methodologies in the previous studies.
- Flora and fauna studies must be carried out in accordance with OEH guidelines
- Survey data of species recorded (species lists) is to be presented in an Access database
- Ground truthing of the vegetation communities, reviewing previous assessments by Biosphere and current Office of Environment and Heritage (OEH) mapping.
- Methodology should be in accordance with the Native Vegetation Interim Type Standard (DECCW 2009).
- GIS data is to be provided for individual threatened species, and rare or significant species.

## B2.0 Existing mappings

**Office of Environment and Heritage (OEH)** mapped the vegetation of the Sydney Metropolitan Area (OEH 2013, VIS\_ID 3817). The six parks were mapped with differing accuracy, with some mapped polygons visited by the mappers, and others assessed using aerial photographs.

The six parks are mapped by OEH (2013) (Figures 6a-1, 6a-2, 6a-3), namely:

### OEH (2013) mapping of the parks

Park / reserve	Vegetation community	Community Code	Accuracy assessment	VEGCODE
Bell Park	Blue Gum High Forest	S_WSF01	2: High: Not visited, photo pattern consistent	190210244, 190220314
Brush Farm Park	Blue Gum High Forest	S_WSF01	1: Very high: Site visited by mappers	190110203
			2: High: Not visited, photo pattern consistent	190220204
	Coastal Warm Temperate Rainforest	S_RF03	1: Very high: Site visited by mappers	253113203
			4: Low: Not visited, photo pattern unexplained	253413204
	Sydney Turpentine-Ironbark Forest	S_WSF09	1: Very high: Site visited by mappers	250110132
			2: High: Not visited, photo pattern consistent	250210133
Darvall Park	Blue Gum High Forest	S_WSF01	1: Very high: Site visited by mappers	190111132
			2: High: Not visited, photo pattern consistent	190224244



Park / reserve	Vegetation community	Community Code	Accuracy assessment	VEGCODE
Lambert Park	Blue Gum High Forest	S_WSF01	2: High: Not visited, photo pattern consistent	190210244, 190220204, 253113203
	Coastal Warm Temperate Rainforest	S_RF03	4: Low: Not visited, photo pattern unexplained	253413204
Field of Mars Reserve	Coastal Enriched Sandstone Dry Forest	S_DSF04	2: High: Not visited, photo pattern consistent	183212132, 189212131, 189212203, 199212203
			3: Medium: Not visited, photo pattern reasonable	183311243, 199312131
	Coastal Enriched Sandstone Moist Forest	S_WSF02	2: High: Not visited, photo pattern consistent	164210132, 182210132, 189210131, 189210152
			3: Medium: Not visited, photo pattern reasonable	182310134
	Coastal Sandstone Gallery Rainforest	S_RF02	1: Very high: Site visited by mappers	157110132
	Coastal Shale-Sandstone Forest	S_WSF06	1: Very high: Site visited by mappers	199111131
			2: High: Not visited, photo pattern consistent	199211203, 199212132, 199213203
			3: Medium: Not visited, photo pattern reasonable	199212132
	Estuarine Mangrove Forest	S_SW01	1: Very high: Site visited by mappers	180100000
	Estuarine Reedland	S_FrW06	4: Low: Not visited, photo pattern unexplained	120433134
	Estuarine Swamp Oak Forest	S_FoW08	0: Not assessed	181000000
			5: Very high: Site visited by others	119500000
	Sydney Turpentine-Ironbark Forest	S_WSF09	2: High: Not visited, photo pattern consistent	189211143
			4: Low: Not visited, photo pattern unexplained	189411132
	Weeds and Exotics	Weed_Ex	0: Not assessed	902000000
1: Very high: Site visited by mappers			902100000	
Outlook Park	Blue Gum High Forest	S_WSF01	2: High: Not visited, photo pattern consistent	190210241, 190210243

**Biosphere Environmental Consultants (2006)** used the following methods:

*Vegetation communities were determined by assessing colour aerial photographs supplied by Council and then ground-truthed. Geology and soil types were also determined. It was stipulated by Council that the methods used for this biodiversity survey were to be the same as used by the National Parks and Wildlife Service (NSW). The model for this survey was taken from Tozer (2003). Quadrats were to be 400 m<sup>2</sup>*

(0.04ha) in area. In most cases this was achievable with 20m x 20m quadrats except in two sites where 40m x 10m quadrats were necessary, either due to the narrowness of the vegetation community or the reserve. One extra 10 X 10m quadrat was set up in a small native remnant in Lambert Park. Quadrats were placed in areas of highest diversity of local native plants with consideration of the required size of the quadrat and the narrowness of the reserves. In order to assess abundance a Braun-Blanquet scale was used. While this method involves a subjective or qualitative description, it also provides for a quantitative, or measurable documentation for comparison of plant community characteristics, especially species richness. Therefore, an inventory of plant species and approximate species numbers was completed for each quadrat then each species was assigned a Braun-Blanquet Cover Class. In order to adequately assess the foliage projective cover of tree species, which may have a dominant effect on other plants within the quadrat, the diameter at breast height (dbh) of the dominant tree species as well as an assessment of the Specht Vegetation Structure (Table 6.1 in Recher, Lunney & Dunn, 1986) is also provided.

Finally, species contained in the quadrats were compared to species listed in the map units described by Tozer (2003) for classification purposes.

The Biosphere (2006) quadrat data has been supplied to OEH and is present on the VIS database.

Biosphere (2006) mapped the vegetation of four of the six parks, as follows (Figures 6b-1, 6b-2, 6b-3), which is not consistent with OEH (2013) mapping. The common map units between the two mapping are highlighted in **blue**.

#### Comparison of mapping by Biosphere (2006) and by OEH (2013) of the parks

Park/ Reserve	Native vegetation (Appendix 5, Biosphere 2006)	OEH 2013 map units
Bell Park	Not part of the 2006 study area	
Brush Farm Park	<b>Sydney Turpentine Ironbark Forest</b> (entire park)	Coastal Warm Temperate Rainforest Blue Gum High Forest <b>Sydney Turpentine-Ironbark Forest</b> Urban Exotic/Native
Darvall Park	<b>Sydney Turpentine Ironbark Forest</b> (entire park)	Blue Gum High Forest Urban Exotic/Native
Field of Mars Reserve	<b>Estuarine Complex</b> HSS Gully Forest HSS Ridgetop Woodland Shale-Sandstone Transition Forest (high SS influence) Turpentine Ironbark Margin Forest disturbed soils/weeds	Coastal Enriched Sandstone Dry Forest Coastal Enriched Sandstone Moist Forest Coastal Sandstone Gallery Rainforest Coastal Shale-Sandstone Forest <b>Estuarine Reedland</b> <b>Estuarine Mangrove Forest</b> <b>Estuarine Swamp Oak Forest</b> Sydney Turpentine-Ironbark Forest Urban Exotic/Native Weeds and Exotics
Lambert Park	Sydney Turpentine Ironbark Forest	Coastal Warm Temperate Rainforest Blue Gum High Forest Urban Exotic/Native
Outlook Park	Not part of the 2006 study area	Blue Gum High Forest

### B3.0 The 2016 survey

The 2016 flora survey replicated and reconciled the flora survey of Biosphere (2006), using the same methodology. In addition to re-surveying of the Biosphere (2006) quadrats, two additional parks, Bell Park and Outlook Park, were surveyed (Figures 7a, 7b, 7c). The data recorded by Biosphere (2006) are on the OEH VIS dataset, with the re-surveyed 2016 quadrats as follows:

Park / reserve	Biosphere 2006		OEH VIS Data (Biosphere 2006)	ACA 2016
	Quadrat	Name	Code	Quadrat
Bell Park	-	-	-	BP#1
Brush Farm	1	Sydney Blue Gum	RYDESBG1	BF#1
Brush Farm	2	Schizomeria	RYDE_S1	BF#2
Brush Farm	3	Turpentine	RYDE_TI1	BF#3
Brush Farm	4	Turpentine-Cassine	RYDE_TC1	BF#4
Darvall	6	Darvall	RYDE_DP1	DP#1
Field of Mars	7	Estuarine	RYDE_E1	FoM#1
Field of Mars	8	Scribbly Gum	RYDE_SG1	FoM#4
Field of Mars	9	Wellington Road	RYDE_WR1	FoM#2
Field of Mars	10	Coachwood	RYDE_C1	FoM#3
Field of Mars	11	Burnt Sclerophyll	RYDE_BS1	FoM#5
Lambert	5	Lilli Pilly	RYDE_LP1	LP#1
Outlook Park	-	-	-	OP#1

The 2016 survey data were used to verify the previous vegetation mapping with additional data recorded from four extra ground truthing (GT) quadrats in Field of Mars Reserve (Quadrat FoM GTQA, FoM GTQB, FoM GTQC, FoM GTQD) and extra spot locations in the parks.

A total of 354 species (233 local native, 28 non-local native and 93 exotic) were recorded in the six parks (Tables 1, 4a to 4f) by Tony Rodd, Jessica Gardner, Dr AnneMarie Clements, Ruth Palsson and Lucy Bonanno. From the percentage of local native to total species recorded in 0.04 ha quadrats, the species composition of the parks varied from:

- predominantly local native composition (>75% local native) in Brush Farm Park and Field of Mars Reserve; to
- predominately exotic and/or non-local native species recorded (<25% local native) in Bell Park.

The remainder of the 0.04 ha quadrats had between 52 to 73% of the species recorded being local native, namely:

2016 sampling locations	2016 survey date	Total number of species recorded	Number of local native species recorded	Number of exotic species recorded	Number of non-local native species recorded	% Local native to total recorded	% Exotic to total recorded	% Non-local native to total recorded
<b>Bell Park</b>								
Quadrat BP#1	15/04	20	3	11	6	15	55	30
Spot location A	15/04	11	3	7	1	27	64	9
Spot location B	15/04	16	3	9	4	19	56	25
Spot BP GTa	15/04	20	11	5	4	55	25	20
Spot BP GTb	15/04	16	3	10	3	19	63	19
<b>Total</b>		<b>59</b>	<b>18</b>	<b>28</b>	<b>13</b>	<b>31</b>	<b>48</b>	<b>22</b>
<b>Brush Farm Park</b>								
Quadrat BF#1	8/04	54	32	18	4	59	33	7
Quadrat BF#2	13/04, 20/05	38	29	7	2	76	18	5
Quadrat BF#3	13/04	29	20	6	3	69	21	10
Quadrat BF#4	19/04	48	32	15	1	67	31	2
Spot BF GTa	15/04	34	26	6	2	77	18	6
Spot BF GTb	15/04	23	15	5	3	65	22	13
Spot BF GTc	20/05	30	26	0	4	87	0	13
<b>Total</b>		<b>116</b>	<b>75</b>	<b>32</b>	<b>9</b>	<b>65</b>	<b>28</b>	<b>8</b>
<b>Darvall Park</b>								
Quadrat DP#1	19/04	51	31	18	2	61	35	4
Spot DP GTa	29/07	19	11	3	5	58	16	26
Spot DP GTb	29/07	11	9	2	0	82	18	0
Spot DP GTc	29/07	14	10	4	0	71	29	0
Spot DP GTd	29/07	16	11	4	1	69	25	6
Spot DP GTe	29/07	33	29	3	1	88	9	3
Spot DP GTf	29/07	16	11	3	2	69	19	13
Spot DP GTg	29/07	13	4	9	0	31	69	0
<b>Total</b>		<b>91</b>	<b>56</b>	<b>28</b>	<b>7</b>	<b>62</b>	<b>31</b>	<b>8</b>
<b>Field of Mars Reserve</b>								
Quadrat FoM#1	30/03	31	16	15	0	52	48	0
Quadrat FoM#2	30/03	46	33	12	1	72	26	2
Quadrat FoM#3	30/03	37	27	10	0	73	27	0
Quadrat FoM#4	7/04	53	47	6	0	89	11	0
Quadrat FoM#5	7/04	65	60	5	0	92	8	0
Quadrat FoM GTQA	6/10	54	42	10	2	78	19	4
Quadrat FoM GTQB	5/10	56	53	3	0	95	5	0
Quadrat FoM GTQC	5/10	51	49	2	0	96	4	0
Quadrat FoM GTQD	6/10	61	53	8	0	87	13	0
Spot location FoM A	5/10	13	4	9	0	31	69	0
Spot location FoM B	5/10	4	2	1	1	50	25	25
Spot FoM GTa	5/10	2	2	0	0	100	0	0
Spot FoM GTb	6/10	23	13	9	1	57	39	4
Spot FoM GTc	6/10	16	10	6	0	63	38	0
<b>Total</b>		<b>217</b>	<b>171</b>	<b>42</b>	<b>4</b>	<b>79</b>	<b>19</b>	<b>2</b>


2016 sampling locations	2016 survey date	Total number of species recorded	Number of local native species recorded	Number of exotic species recorded	Number of non-local native species recorded	% Local native to total recorded	% Exotic to total recorded	% Non-local native to total recorded
<b>Lambert Park</b>								
Quadrat LP#1	7/04	48	25	17	6	52	35	13
Spot LP GTa	29/07	13	10	2	1	77	15	8
Spot LP GTb	29/07	14	7	6	1	50	43	7
Spot LP GTc	29/07	9	6	3	0	67	33	0
Spot LP GTd	29/07	11	9	2	0	82	18	0
Spot LP GTe	29/07	21	13	4	4	62	19	19
Spot LP GTf	29/07	21	15	3	3	71	14	14
<b>Total</b>		<b>85</b>	<b>47</b>	<b>28</b>	<b>10</b>	<b>55</b>	<b>33</b>	<b>12</b>
<b>Outlook Park</b>								
Quadrat OP#1	22/04	50	33	11	6	66	22	12
Spot OP GTa	22/04	37	24	9	4	65	24	11
Spot OP GTb	22/04	5	3	2	0	60	40	0
Spot OP GTc	22/04	5	4	1	0	80	20	0
<b>Total</b>		<b>70</b>	<b>44</b>	<b>18</b>	<b>8</b>	<b>63</b>	<b>26</b>	<b>11</b>

Note: GT = Ground truthing


### B3.1 Methods

The flora methods used in the 2016 survey are consistent with Biosphere (2006) with the locations of the 2006 quadrats relocated from the GIS registered figures in Biosphere (2006).

The 0.04 ha quadrats (Figures 7a, 7b, 7c) consisted of four contiguous 10 m x 10 m sub-quadrats. The quadrats and sub-quadrats were laid out as follows:



In Darvall Park Quadrat DP#1, and Field of Mars FoM#3, the 0.04 ha quadrats were laid out as follows:



Supplementary species data consisted of species present in an approximately 10 m radius of Spot locations being recorded (Tables 4a to 4f). During the ground truthing of the vegetation communities, the dominant species in the strata were recorded in Ground Truthing Spots (see Figures 8a, 8b-1, 8b-2, 8c, Tables 4a to 4f).

Biosphere (2006) used a 7-stage Braun-Blanquet scale technique for species abundance in each quadrat, namely:

*While this method involves a subjective or qualitative description, it also provides for a quantitative, or measurable documentation for comparison of plant community characteristics, especially species richness.*

Cover abundance scores (1 to 7) for the Braun-Blanquet Cover Classes are given in Biosphere (2006, page 16) and interpreted in the 2016 survey as follows:

Cover Class	Biosphere (2006)		Interpreted in 2016 as
1	Rare	few individuals (three or less) and cover <5%	≤ 3 individuals AND <5% cover
2	Uncommon	more than three but not consistently throughout the plot) and cover <5%	>3 individuals AND cover <5%
3	Common	consistent throughout the plot and cover <5%	cover <5% throughout the quadrat
4		Very abundant cover <5% OR cover >5% but <20%	6–19% cover in quadrat
5		Cover >20% but <50%	21% – 49% cover in quadrat
6		Cover >50% but <75%	51% – 74% cover in quadrat
7		Cover >75% but <100%	76% – 99% cover in quadrat

The Braun-Blanquet Cover Classes score for the 11 re-surveyed quadrats in 2016 are given in Table 3.

Given the subjective estimates of the Braun-Blanquet abundance scores, the statistician Dr Margaret Donald advised that for the 2016 dataset the percent projected foliage cover for each species be recorded in the four 10 m x 10 m sub-quadrats in the quadrats. The Braun-Blanquet abundance scores were estimated in 2016 survey (Table 3) but not used in the statistical analyses.

In addition, for each quadrat the maximum height and number of individuals for each species greater than 2 m high being recorded in each of the 10 m x 10 m sub-quadrats (Tables 2, 5).

Ground truthing methods were consistent with the Native Vegetation Interim Standard (DECCW 2009). The field work consisted of an inspection, additional Ground Truth (GT) quadrats and Rapid survey at Ground Truth (GT) spot locations (as required) of each of the OEH (2013) mapped vegetation communities (Figures 8a, 8b-1, 8b-2, 8c).

In the Field of Mars Reserve, there were additional quadrats along with soil transects (Figures 3a-4, 3b-3) to assist in confirming consistency of the soil with the soil descriptions in the Final Determination of the vegetation communities.

Sampling locations were photographed at the time of survey (Appendix 4) and GPS co-ordinates were recorded using a hand held *Garmin GPSMAP® 78* unit.

A targeted search for rare plants known to flower in Spring was undertaken in the Spring survey.


Nomenclature is consistent with Harden (1990, 1992, 1993, 2002), Harden and Murray (2000) and subsequent taxonomic changes as published in *Telopea*, the Sydney Royal Botanic Gardens' journal of systematic botany, and in other Australian taxonomic literature. The Royal Botanic Gardens' PlantNet website ([plantnet.rbgsyd.nsw.gov.au](http://plantnet.rbgsyd.nsw.gov.au)) incorporating Flora Online is the major source for updated taxonomy.

## B3.2 Observations/ Findings


### B3.2.1 Comparison of 2006 and 2016 surveys

In terms of species composition recorded in 2006 and 2016 surveyed quadrats in four of the six parks (statistician report in Appendix 5, data summary table Appendix 6), it was found that:

- The Venn diagrams for “all parks” show the numbers of species in common between the 2006 and 2016 surveys with 171 of the 374 species recorded common to both 2006 and 2016, and more species recorded in 2016 than in 2006. The difference between years in the four parks may be a response by the exotic species to better rainfall prior to the 2016 Autumn survey.





All flora species across the four parks



All **native** flora species across the four parks

All Parks: Exotic Flora spp.

All Parks: NLN Flora spp.




All **exotic** flora species across the four parks

All **non-local native** flora species across the four parks

The Venn diagrams below overlapping native species between the four parks in 2006 and 2016 show that Field of Mars Reserve is clearly distinct from the other parks on Wianamatta Shale derived soil in its native species composition. There were 98 species unique to Field of Mars Reserve in 2016, related to the presence of species associated with low nutrient Hawkesbury Sandstone derived soils on rugged sandstone slopes and of estuarine species.


2016



Venn diagram 2016, with BFP = Brush Farm Park, FoM = Field of Mars, LP = Lambert Park, DP = Darvall Park.



2006



Venn diagram 2006, with BFP = Brush Farm Park, FoM = Field of Mars, LP = Lambert Park, DP = Darvall Park.

For the four parks in the 2006 and 2016 surveyed quadrats the species composition was as follows:

**Comparison of number of species recorded in 2006 and 2016 in the parks**

	Only 2006	2006 and 2016	Only 2016
<b>All parks</b>			
Total number of species	88	171	115
Number of local native species	65	128	57
Number of non-local	10	3	18
Number of exotic	23	30	50
<b>Brush Farm Park</b>			
Total number of species	28	79	38
Number of local native species	21	53	23

	Only 2006	2006 and 2016	Only 2016
Number of non-local	7	3	6
Number of exotic	8	15	17
<b>Darvall Park</b>			
Total number of species	49	30	24
Number of local native species	29	17	17
Number of non-local	6	0	2
Number of exotic	19	8	10
<b>Field of Mars Reserve</b>			
Total number of species	55	97	58
Number of local native species	44	78	44
Number of non-local	5	1	0
Number of exotic	11	13	19
<b>Lambert Park</b>			
Total number of species	22	24	24
Number of local native species	13	12	13
Number of non-local	2	0	6
Number of exotic	9	10	7

In order to statistically compare the flora composition in the 2006 quadrats and 2016 re-surveyed quadrats, the subjective estimates of the Braun-Blanquet abundance scores (class covers) were converted to percent projected foliage covers.

Of the four parks surveyed in both 2006 and 2016, only Brush Farm Park (four quadrats) and the Field of Mars Reserve (five quadrats) had sufficient quadrats to enable comparison between the two years in the parks, resulting in following comparisons being undertaken:

2006	vs.	2016
Brush Farm Park (four quadrats)	vs.	Brush Farm Park (four quadrats)
Field of Mars Reserve (five quadrats)	vs.	Field of Mars Reserve (five quadrats)
<b>All parks (11 quadrats):</b> Brush Farm Park (4 quadrats) Darvall Park (1 quadrat) Field of Mars Reserve (5 quadrats) Lambert Park (1 quadrat)	vs.	<b>All parks (11 quadrats):</b> Brush Farm Park (4 quadrats) Darvall Park (1 quadrat) Field of Mars Reserve (5 quadrats) Lambert Park (1 quadrat)

At the 5% significance level, it was found that:

Analysis	Parks compared		
	Brush Farm Park	Field of Mars Reserve	“All parks”
% projected foliage of native cover between 2006 and 2016	No statistically significant difference	No statistically significant difference	No statistically significant differences
% projected foliage of exotic cover between 2006 and 2016	No statistically significant difference	No statistically significant difference	A statistically significant difference indicating a decrease in exotic cover

Overall, when comparing the data from 2006 to 2016, the cover of native species has remained constant, and the cover by exotic species has declined.

### B3.2.2 Exotic and non-local native species

From the 2016 recorded data for the six parks, the extent of planting and subsequent colonisation by non-local native species and the infestation by exotic species varied between the parks. The Blue Gum High Forest mapping by OEH (2013) was affected by the extent of planting of non-local native species *Eucalyptus grandis*, leading to over mapping of the extent of the Blue Gum High Forest mapping.

In the tables below, (\*) denotes exotic and (#) denotes non-local native species. Values for quadrats (BP#1, BF#1 etc.) are the average percent (%) projected foliage cover for each quadrat (calculated from the percent (%) projected foliage cover recorded for each sub-quadrat). Only those species with an average percent projected foliage cover greater than or equal to 1% ( $\geq 1\%$ ) are listed. Species are listed in order from highest to lowest average percent (%) projected foliage cover  $\geq 1\%$ .

**Bell Park** is extensively colonised by non-local trees including *Corymbia citriodora*, *Eucalyptus botryoides*, *Eucalyptus grandis* and *Eucalyptus microcorys*. This park was not found to support Blue Gum High Forest as mapped by OEH (2013), but planted non-local native *Eucalyptus grandis* rather than naturally occurring *Eucalyptus saligna* and *Eucalyptus pilularis*.

The percent projected foliage cover for *Lantana camara* in patches of this park is above the 75% weed cover threshold (Gooden *et al.* 2009a, 2009b). Gooden *et al.* (2009b) found that the number of native species declined rapidly at locations where *Lantana* percent projected foliage cover exceeded 75%.

#### Bell Park average percent projected foliage cover ( $\geq 1\%$ only) per species in the surveyed quadrat

Exotic/ NLN	Species	Common names	Percentage cover BP#1
*	<i>Lantana camara</i>	Lantana	77.5
#	<i>Eucalyptus grandis</i>	Flooded Gum	17.5
*	<i>Ipomoea indica</i>	Morning Glory	9.5
*	<i>Jasminum polyanthum</i>	Jasmine	7.0
#	<i>Eucalyptus botryoides</i>	Bangalay	6.3
#	<i>Corymbia citriodora</i>	Lemon-scented Gum	5.0
*	<i>Ligustrum lucidum</i>	Broad-leaved Privet	3.0
#	<i>Eucalyptus scoparia</i>	Wallangarra White Gum	2.5
#	<i>Grevillea robusta</i>	Silky Oak	1.3
#	<i>Melaleuca armillaris</i>	Bracelet Honey-myrtle	1.3
*	<i>Delairea odorata</i>	Cape Ivy	1.0

**Brush Farm Park** has had some planting and subsequent colonisation by non-local native species *Eucalyptus grandis* and *Eucalyptus microcorys*, but not as extensive as in Bell Park.

The recorded vegetation is consistent with the OEH (2013) mapping. The mapped occurrence of Temperate Rainforest is predominantly the weedy native species *Pittosporum undulatum*

(Clements 1983, Gleadow and Ashton 1981, Gleadow 1982, Gleadow and Rowan 1982, Gleadow *et al.* 1983, Gleadow and Narayan 2007).

**Brush Farm Park average percent projected foliage cover ( $\geq 1\%$  only) per species in the surveyed quadrats**

Exotic / NLN	Species	Common name	Percentage cover			
			BF#1	BF#2	BF#3	BF#4
*	<i>Ehrharta erecta</i>	Panic Veldgrass	15	1.3	2	47.5
*	<i>Tradescantia fluminensis</i>	Wandering Jew	45	7.5	-	0.5
*	<i>Chlorophytum comosum</i>	Spider Plant	-	-	-	11.8
#	<i>Eucalyptus microcorys</i>	Tallowwood	5	-	-	-
*	<i>Lantana camara</i>	Lantana	-	-	-	4.8
#	<i>Brachychiton acerifolius</i>	Illawarra Flame Tree	0.5	2	2.3	1.3
*	<i>Passiflora suberosa</i>	Cork Passionflower	-	-	-	1.8
*	<i>Ochna serrulata</i>	Micky Mouse Plant	-	1.5	-	0.025
#	<i>Eucalyptus grandis</i>	Flooded Gum	1.3	-	-	-
*	<i>Hedera helix</i>	English Ivy	0.025	-	1.3	-
*	<i>Jacaranda mimosifolia</i>	Jacaranda	1.3	-	-	-

**Darvall Park** has also been planted with *Eucalyptus grandis* and *Eucalyptus microcorys* in the northern part of the park nearest the railway. The most frequently occurring weed recorded in this park is the understorey cover of *Tradescantia fluminensis* (Wandering Jew), followed by *Ehrharta erecta* (Panic Veld-grass). Outside the quadrats, there was extensive infestations of *Lantana camara* in the central and southern areas.

Despite the planting of *Eucalyptus grandis*, there are large remnant trees of *Eucalyptus saligna* and the park is found to be relatively accurately mapped by OEH (2013) as Blue Gum High Forest. It could equally be mapped as Sydney Turpentine Ironbark Forest or an intergrade of the two map units.

**Darvall Park average percent projected foliage cover ( $\geq 1\%$  only) per species in the surveyed quadrat**

Exotic / NLN	Species	Common name	Percentage cover DP#1
*	<i>Tradescantia fluminensis</i>	Wandering Jew	42.5
*	<i>Ehrharta erecta</i>	Panic Veldgrass	9.3
#	<i>Eucalyptus grandis</i>	Flooded Gum	2.5
*	<i>Cardiospermum grandiflorum</i>	Balloon Vine	1.3
#	<i>Eucalyptus microcorys</i>	Tallowwood	1.3
*	<i>Megathyrsus maximus</i>	Guinea Grass	1.3
*	<i>Rubus laudatus</i>	Plains Blackberry	1.0

**Field of Mars Reserve** has little or no planting of non-local native species, except in the former depot site, the edge of the Field of Mars Cemetery and in the mown areas in the south eastern part of the reserve.

The extent of exotic species is concentrated in the south-west, especially downslope of the former depot and the cemetery, as well as at the rear of adjoining houses. The occurrence of *Lantana camara* is generally infrequent within the sandstone vegetation. The bird drop weeds (especially *Lantana camara*, *Ligustrum* spp.) pose a major risk in areas with nutrient enrichment.

The vegetation data recorded are consistent with the OEH (2013) mapping.

**Field of Mars Reserve average percent projected foliage cover ( $\geq 1\%$  only) per species in the surveyed quadrats**

Exotic/ NLN	Species	Common name	Percentage cover								
			FoM #1	FoM# 2	FoM# 3	FoM# 4	FoM# 5	FoM GTQ A	FoM GTQ B	FoM GTQ C	FoM GTQ D
*	<i>Ligustrum lucidum</i>	Broad-leaved Privet	-	0.08	2	-	-	0.26	-	-	0.25
#	<i>Cyathea cooperi</i>	Straw Tree-fern	-	-	-	-	-	1.75	-	-	-
*	<i>Ligustrum sinense</i>	Chinese Privet	-	0.05	1.5	0.3	0.05	1.5	-	-	0.75
*	<i>Ochna serrulata</i>	Micky Mouse Plant	-	0.1	0.28	0.05	0.05	1.25	0.05	0.03	0.3
*	<i>Lantana camara</i>	Lantana	-	0.05	-	0.03	0.05	1	-	-	0.75

**Lambert Park** has been planted and colonised by *Eucalyptus grandis* and *Casuarina cunninghamiana*.

The vegetation data recorded was not consistent with the OEH (2013) mapping of Blue Gum High Forest, as there are no naturally occurring *Eucalyptus saligna* but extensive plantings.

**Lambert Park average percent projected foliage cover ( $\geq 1\%$  only) per species in the surveyed quadrat**

Exotic/ NLN	Species	Common name	Percentage cover LP#1
*	<i>Tradescantia fluminensis</i>	Wandering Jew	8
#	<i>Eucalyptus grandis</i>	Flooded Gum	3.8
#	<i>Casuarina cunninghamiana</i>	River Oak	2.5
*	<i>Ehrharta erecta</i>	Panic Veldgrass	2.0

**Outlook Park** has been extensively planted with *Eucalyptus microcorys* in the west of the park.

The vegetation data recorded was not consistent with the OEH (2013) mapping of Blue Gum High Forest as there were only two remnant *Eucalyptus saligna* in the park (recorded in Quadrat OP#1) and extensive planting.

**Outlook Park average percent projected foliage cover ( $\geq 1\%$  only) per species in the surveyed quadrat**

Exotic/ NLN	Species	Common name	Percentage cover OP#1
#	<i>Grevillea robusta</i>		2.5
#	<i>Brachychiton acerifolius</i>		1.5

### B3.3 Ground truthing confirmation of the previous mapping

The Biosphere 2006 and OEH (2013) mapping has been assessed (see section B3.1 for ground truthing methodology) using the 2016 survey data .

Park/ Reserve	Native vegetation (Appendix 5, Biosphere 2006)	OEH 2013 map units	Finding of the 2016 survey data
Bell Park	Not part of the 2006 study area	Blue Gum High Forest	NOT confirmed as Blue Gum High Forest.
		Urban Exotic/Native	Confirmed as Urban Exotic/Native.
Brush Farm Park	Sydney Turpentine Ironbark Forest (entire park)	Coastal Warm Temperate Rainforest	Predominantly the weedy native <i>Pittosporum undulatum</i> .
		Blue Gum High Forest	Confirmed as Blue Gum High Forest with remnant <i>Eucalyptus saligna</i> .
		Sydney Turpentine- Ironbark Forest	Confirmed Sydney Turpentine-Ironbark Forest.
		Urban Exotic/Native	Confirmed as Urban Exotic/Native associated with Sportsground and Scout Hall.
Darvall Park	Sydney Turpentine Ironbark Forest (entire park)	Blue Gum High Forest	Confirmed as Blue Gum High Forest but with plantings and weed dominated close to property boundaries and swampy depression.
		Urban Exotic/Native	Confirmed as Urban Exotic/Native.
Field of Mars Reserve	Estuarine Complex HSS Gully Forest HSS Ridgetop Woodland Shale-Sandstone Transition Forest (high SS influence) Turpentine Ironbark Margin Forest disturbed soils/weeds	Coastal Enriched Sandstone Dry Forest Coastal Enriched Sandstone Moist Forest Coastal Sandstone Gallery Rainforest Coastal Shale- Sandstone Forest Estuarine Reedland Estuarine Mangrove Forest Estuarine Swamp Oak Forest Sydney Turpentine- Ironbark Forest Urban Exotic/Native	Confirmed.

Park/ Reserve	Native vegetation (Appendix 5, Biosphere 2006)	OEH 2013 map units	Finding of the 2016 survey data
		Weeds and Exotics	
Lambert Park	Sydney Turpentine Ironbark Forest	Coastal Warm Temperate Rainforest	Confirmed as Coastal Warm Temperate Rainforest, but with high occurrence of <i>Pittosporum undulatum</i> .
		Blue Gum High Forest	Not confirmed. Only <i>Eucalyptus saligna</i> planted in south.
		Urban Exotic/Native	Confirmed as Urban Exotic/Native.
Outlook Park	Not part of the 2006 study area	Blue Gum High Forest	Not confirmed as Blue Gum High Forest, but planted canopy with only two remnant <i>Eucalyptus saligna</i> .

### B3.4 Targeted threatened species searches

The conservation significance of species that may be present in the parks were assessed at a:

1. National level against the schedules of the Commonwealth *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act).
2. State level against the schedules of the *Threatened Species Conservation Act 1995* (TSC Act).

#### B3.4.1 Database searches

Database searches were undertaken to determine the possible threatened species in the Ryde LGA, with:

- 19 plant species listed in the Protected Matters Search (EPBC Act 1999) (<https://www.environment.gov.au/epbc/protected-matters-search-tool>, accessed 30 November 2016), and
- 46 plant species listed under the NSW *Threatened Species Conservation Act 1995* are recorded within the Ryde LGA (NSW Office of Environment and Heritage Bionet - Atlas of NSW Wildlife, <http://www.bionet.nsw.gov.au>, accessed 30 November 2016).

Species	Status under EPBC Act	Status under TSC Act
<i>Acacia bynoeana</i>	Vulnerable	Endangered
<i>Acacia pubescens</i>	Vulnerable	Vulnerable
<i>Acacia terminalis</i> subsp. <i>terminalis</i>	Endangered	Endangered
<i>Allocasuarina glareicola</i>	Endangered	-
<i>Allocasuarina portuensis</i>	-	Endangered
<i>Asterolasia elegans</i>	Endangered	-
<i>Caladenia tessellata</i>	Vulnerable	Endangered
<i>Callistemon linearifolius</i>	-	Vulnerable
<i>Chamaesyce psammogeton</i>	-	Endangered

<b>Species</b>	<b>Status under EPBC Act</b>	<b>Status under TSC Act</b>
<i>Cryptostylis hunteriana</i>	Vulnerable	Vulnerable
<i>Cynanchum elegans</i>	-	Endangered
<i>Darwinia biflora</i>	Vulnerable	Vulnerable
<i>Deyeuxia appressa</i>	Endangered	Endangered
<i>Dillwynia tenuifolia</i>	-	Vulnerable
<i>Epacris purpurascens</i> var. <i>purpurascens</i>	-	Vulnerable
<i>Eucalyptus camfieldii</i>	-	Vulnerable
<i>Eucalyptus nicholii</i>	-	Vulnerable
<i>Genoplesium baueri</i>	Endangered	Endangered
<i>Grammitis stenophylla</i>	-	Endangered
<i>Grevillea caleyi</i>	-	Critically Endangered
<i>Grevillea parviflora</i> subsp. <i>parviflora</i>	-	Vulnerable
<i>Gyrostemon thesioides</i>	-	Endangered
<i>Haloragodendron lucasii</i>	-	Endangered
<i>Hibbertia puberula</i>	-	Endangered
<i>Hibbertia</i> sp. Bankstown	-	Critically Endangered
<i>Hibbertia spanantha</i>	-	Critically Endangered
<i>Hibbertia superans</i>	-	Endangered
<i>Leptospermum deanei</i>	Vulnerable	Vulnerable
<i>Leucopogon exolasius</i>	-	Vulnerable
<i>Maundia triglochinooides</i>	-	Vulnerable
<i>Melaleuca biconvexa</i>	Vulnerable	-
<i>Melaleuca deanei</i>	Vulnerable	Vulnerable
<i>Microtis angusii</i>	-	Endangered
<i>Pelargonium</i> sp. Striatellum (G.W.Carr 10345)	Endangered	-
<i>Persoonia hirsuta</i>	-	Endangered
<i>Persoonia nutans</i>	-	Endangered
<i>Pimelea curviflora</i> var. <i>curviflora</i>	Vulnerable	Vulnerable
<i>Pimelea spicata</i>	Endangered	Endangered
<i>Pomaderris prunifolia</i>	-	Endangered Population
<i>Prasophyllum fuscum</i>	-	Critically Endangered
<i>Prostanthera marifolia</i>	-	Critically Endangered
<i>Pterostylis saxicola</i>	Endangered	Endangered
<i>Pterostylis</i> sp. Botany Bay	-	Endangered
<i>Pultenaea pedunculata</i>	-	Endangered
<i>Senecio spathulatus</i>	-	Endangered
<i>Syzygium paniculatum</i>	Vulnerable	Endangered
<i>Tetratheca glandulosa</i>	-	Vulnerable
<i>Thesium australe</i>	Vulnerable	-
<i>Wahlenbergia multicaulis</i>	-	Endangered Population
<i>Wilsonia backhousei</i>	-	Vulnerable
<i>Zannichellia palustris</i>	-	Endangered



From the searches of Australia's Virtual Herbarium (specimens held in herbaria in Australia, website <http://avh.ala.org.au/>, accessed 30 November 2016), the records for Ryde LGA of listed threatened species are as follows:

Species	Record location	Collection date	Collector	Comm. status	NSW status
<i>Callistemon linearifolius</i>	Lane Cove in Ryde LGA	Not given	Fletcher, J.J.		Vulnerable
<i>Darwinia biflora</i>	Hunters Hill	Supplied date "1892-08"	Andrew	Vulnerable	Vulnerable
	SE corner of Riverside Corporate Park, North Ryde, beside Epping Road	Supplied date "1998-09"	Rodd, A.N., Moore, R.; Clements, A.		
<i>Epacris purpurascens</i> var. <i>purpurascens</i>	Field of Mars Reserve, Ryde Municipality	1989-01-05	Gibson, C.		Vulnerable
<i>Genoplesium baueri</i>	Not specified, but in Ryde LGA	Supplied date "1884-04"	Deane, H.	Endangered	Endangered
	Not specified, but in Ryde LGA	Supplied date "1884-05"	Deane, H.		
	Not specified, but in Ryde LGA	Supplied date "1885-03"	Deane, H.		
	Not specified, but in Ryde LGA	Supplied date "1887-01"	Deane, H.		
	Not specified, but in Ryde LGA	Supplied date "1887-02"	Deane, H.		
	Not specified, but in Ryde LGA	Supplied date "1986-03-30"	Clements, M.A.		
<i>Grammitis stenophylla</i>	Field of Mars [NB - this name applied to most of Ryde LGA plus Epping in 1884!]	Supplied date "1884-04"	Deane, H.		Endangered
<i>Melaleuca deanei</i>	Not specified, but in Ryde LGA	1930-10-07	Supplied as "[unknown]"	Vulnerable	Vulnerable
<i>Pimelea curviflora</i> var. <i>curviflora</i>	Not specified, but in Ryde LGA	Supplied date "1884-05"	Deane, H.	Vulnerable	Vulnerable
	Hunters Hill	Supplied date "1914-12"	Musson, C.T.		
	Gladesville	Supplied date "1883-08"	Deane, H.		
<i>Prostanthera marifolia</i>	Not specified, but in Ryde LGA	Supplied date "1903-11"	Dixon, W.A.	Critically Endangered	Critically Endangered
	Not specified, but in Ryde LGA	Supplied date "1906-09"	Hamilton, A.A.		

Species	Record location	Collection date	Collector	Comm. status	NSW status
<i>Wilsonia backhousei</i>	Ermington, Spurway St. Edge of Parramatta river	1986-10-19	Bishop, W.		Vulnerable
	RAN Armaments Depot, Parramatta River, W of Homebush Bay, c. 3 miles E of Parramatta	1964-11-17	Constable, E.F.		

### B3.4.2 Target searches in the parks

Despite extensive survey in the five parks, only one of the listed threatened species from the database search was recorded:

Species	Status under EPBC Act	Status under TSC Act	Location found
# <i>Syzygium paniculatum</i>	Vulnerable	Endangered	OP#1 BP GTa

\* These values are approximate only as BP GTa is a ground truthing location, and does not have a fixed area.

As *Syzygium paniculatum* is a non-local native species, it is highly likely that the individuals recorded in the parks were planted.

The more intact Field of Mars Reserve was searched (see Figures 9a, 9b). None of the listed threatened species from the database searches were recorded in the quadrats, spot location or search in the 2016 surveys, though it is considered likely that there may habitat for the following species, namely:

Species	Habitat from PlantNET
<i>Callistemon linearifolius</i>	Grows in dry sclerophyll forest on the coast and adjacent ranges, chiefly from Georges R. to the Hawkesbury R.
<i>Darwinia biflora</i>	Grows in heath on sandstone or in the understorey of woodland on shale-capped ridges; Cheltenham to Hawkesbury R., rare.
<i>Epacris purpurascens</i> var. <i>purpurascens</i>	Grows in sclerophyll forest, scrubs and swamps on sandstone from Gosford and Sydney districts.
<i>Genoplesium baueri</i>	Grows in sparse sclerophyll forest and moss gardens over sandstone; from the Hunter Valley to Nowra district.
<i>Grammitis stenophylla</i>	Grows on rocks in rainforest and in wet sclerophyll forest.
<i>Melaleuca deanei</i>	Grows in wet heath on sandstone; uncommon, in coastal districts from Berowra to Nowra.
<i>Pimelea curviflora</i> var. <i>curviflora</i>	Confined to coastal areas around Sydney on sandstone.
<i>Prostanthera marifolia</i>	Woodland dominated by <i>Eucalyptus sieberi</i> and <i>Corymbia gummifera</i> . In deeply weathered clay soil with ironstone nodules.
<i>Wilsonia backhousei</i>	Grows in coastal saltmarshes; chiefly in the Sydney district, also common at Jervis Bay.

## References for Sections A, B

Benson D. and Howell J. (1990) *Taken for Granted: The Bushland of Sydney and its Suburbs*. Kangaroo Press, Sydney.

Biosphere Environmental Consultants (2006) *Ryde Flora and Fauna Study – Brush Farm Park, Lambert Park, Darvall Park and Field of Mars Reserve*. Prepared for City of Ryde.

Chapman G.A. and Murphy C.L. (1989) *Soil Landscapes of the Sydney 1: 100 000 sheet*. Soil Conservation Service of NSW, Sydney.

Chapman G.A., Murphy C.L., Tille P.J., Atkinson G., and Morse R.J. (1989) Accompanying map to *Soil Landscapes of the Sydney 1:100 000 sheet*. Soil Conservation Service of NSW, Sydney.

Gleadow R.M., Rowan K.S., Ashton D.H. (1983) Invasion by *Pittosporum undulatum* of the forests of Central Victoria. IV shade tolerance, *Australian Journal of Botany* [P], vol 31, issue 2, CSIRO Publications, Collingwood Australia, pp. 151-160.

Gleadow R.M. (1982) Invasion by *Pittosporum undulatum* of the forests of Central Victoria. II Dispersal, germination and establishment, *Australian Journal of Botany* [P], vol 30, issue 2, CSIRO Publications, Collingwood Australia, pp. 185-198.

Gleadow R.M., Rowan K.S. (1982) Invasion by *Pittosporum undulatum* of the forests of Central Victoria. III Effects of temperature and light on growth and drought resistance, *Australian Journal of Botany* [P], vol 30, issue 3, CSIRO Publications, Collingwood Australia, pp. 347-357.

Gleadow R.M., Ashton D.H. (1981) Invasion by *Pittosporum undulatum* of the forests of Central Victoria. I Invasion patterns and plant morphology, *Australian Journal of Botany* [P], vol 29, issue 6, CSIRO Publications, Collingwood Australia, pp. 705-720.

Gleadow R.M., Narayan I. (2007) Temperature thresholds for germination and survival of *Pittosporum undulatum*: implications for management by fire, *Acta Oecologica*, vol 31, issue 2, Gauthier-Villars/Editions Elsevier, Paris France, pp. 151-157.

Gooden B., French K., Turner P.J. and Downey P.O. (2009a) Impact threshold for an alien plant invader, *Lantana camara* L., on native plant communities. *Biological Conservation*. 142(11): 2631–2641.

Gooden B., French K. and Turner P.J. (2009b) Invasion and management of a woody plant, *Lantana camara* L., alters vegetation diversity within wet sclerophyll forest in southeastern Australia. *Forest Ecology and Management*. 257(3): 960–967.

Harden G.J. (1990-93, 2002) *Flora of New South Wales Volumes 1 to 4*. University of New South Wales Press, Kensington.

Harden G.J. and Murray L.J. (2000) *Supplement to Flora of New South Wales – Volume 1*. University of New South Wales Press, Kensington.

Herbert C. and West J.L. (1983) *Sydney 1:100 000 Geological Sheet 9130*, 1st edition. Geological Survey of New South Wales, Sydney.

Office of Environment and Heritage (2013) *The native vegetation of the Sydney metropolitan area*. Mapping file VIS\_ID 3817.

Office of Environment and Heritage (2013a) *The native vegetation of the Sydney metropolitan area. Volume 1: Technical Report*. Version 2.0. Office of Environment and Heritage, Department of Premier and Cabinet. Dated October 2013.

Smith K.V. (2005) *Wallumedegal an Aboriginal history of Ryde*. Prepared for City of Ryde. Dated December 2005.

## **Section C: Fauna assessment**

## C1.0 Introduction

The fauna survey replicated biodiversity surveys conducted by Biosphere in 2006 consisting of replicated survey Autumn and Spring in four parks and 2016 Autumn and Spring in two additional parks.

The 2016 surveys were conducted by the following ecologists:

Ecologist	Expertise
Damian Lettoof	Project fauna manager, herpetofauna surveys, nocturnal surveys, bird surveys, hair tubing
David Martin	Nocturnal surveys
Henry Cook	Bird surveys, fish surveys
Dr James Schlunke	Invertebrate surveys
Glenn Hoye	Bat surveys (ANABAT analysis)
Barbara Triggs	Mammal hair analysis

## C1.1 Sampling methods

### C1.1.1 Reptiles

Reptiles were actively and visually searched for on warm, dry days. The 2016 replicated flora quadrats of each park were surveyed for 20 minutes each. In addition, in the parks all walking tracks and potential shelter sites were also surveyed, and opportunistic sightings from the bushland regeneration contractors included. Each park was surveyed for two hours one morning and one afternoon per season.

### C1.1.2 Frogs

Frog surveys were conducted for 20 minutes per quadrat during the nocturnal surveys if there was recent rain, otherwise on specific raining nights. Calling frogs were identified aurally; non-calling frogs were identified by visual inspection. Searches of the area were carried out using head torches. During daylight hours, hand-netting was carried out to search for tadpoles.

### C1.1.3 Mammals

**For small ground mammals**, single entrance, baited hair tubes were used in the remnant bushland areas of each park. The tubes were installed and remained on site for a five days before being collected and hair samples analysed by Barbara Triggs. The number of hair tubes used at each site depended on the size of the reserve, with 5 hair tubes used in eastern bushland of the 0.7 ha Outlook Park to 125 hair tubes in the 46.7 ha Field of Mars Reserve, namely:

Hair tube numbers and location

Park	Size of park	No. of hair tubes	Location of tubes
Bell Park	0.46 ha	10	Scattered in northern bushland and alongside riparian zone of the park
Brush Farm Park	5.4 ha	50	Alongside all walking tracks in park
Darvall Park	6.1 ha	25	Set in a line running from north to south through centre of the park

Park	Size of park	No. of hair tubes	Location of tubes
Field of Mars Reserve	46.7 ha	125	25 traps set around each survey quadrat site
Lambert Park	2.6 ha	15	Around the outer boundaries of the park
Outlook Park	0.7 ha	5	Scattered in the eastern bushland of park

Animal tracks, diggings, burrows were searched for, and shed fur/feathers and scats collected opportunistically, sometimes with the assistance of bush regeneration contractors.

**Arboreal mammals** were detected by nocturnal spotlighting surveys, conducted during the first three hours after dark, over two nights per park. Generally, all of the walking tracks and open areas of the parks were slowly walked by two ecologists, panning spotlights into the trees canopies but also all surrounding areas. If an animal was detected, it was approached until it could be identified visually. Calling species (e.g. flying foxes) were identified by call. In addition, diurnal examinations of trees for scratch marks and drays were conducted.

### **Bats**

Flying foxes were detected by spotlighting at night whereas insectivorous bats were detected using ultra-sonic (ANABAT SD1) bat recorders. The recorders are hand-held and carried through the reserves at night while spotlighting was in progress. Recorded bat calls were later analysed using ANABAT 5.0 software.

#### **C1.1.4 Birds**

**Diurnal birds** (birds active during the day) were surveyed in the early mornings using the survey quadrat areas. Two mornings of twenty minutes survey time (in each survey period) was devoted to recording the birds that were seen or heard. In addition, other bird species detected opportunistically outside the quadrat were also recorded.

#### **Owls and nocturnal birds**

Owl surveys were conducted at night during nocturnal surveys by spotlighting and using a small portable amplifier. Owl calls were broadcast at night for the two local owl species: boobook owls and powerful owls, calls were also played for potential other owls including: sooty owls, masked owls and barn owls. Calls were played at suitable sites each night and the amplifier was aimed away from nearby residences before the sounds were played. A listening period of 2 minutes followed the playing of each tape. If it was possible to visually identify the responding owl, all attempts were made to do so. Other night birds, such as tawny frogmouths were also spotlighted during night surveys.

#### **C1.1.5 Fish**

Small hand nets were used to sample for fish in the drainage lines in Lambert Park, the creeklines in Darvall Park, Bell Park and Field of Mars Reserve. Due to the ephemeral nature of Archer Creek in Brush Farm Park and piped creekline in Outlook Park, there were no water bodies to survey for fish.

#### **C1.1.6 Invertebrate fauna**

Invertebrates were surveyed within the quadrats (two mornings of twenty minutes search effort per season) as well as habitat specific searches. This involved sweep-netting creeks, searching leaf-litter and beneath rocks/logs for spiders, insects and other soft-bodied creatures and using

small battery operated night lights for two evenings per park to collect nocturnal insects. Invertebrates were identified to genus if possible, otherwise class or order. The number of different species within each group collected was also recorded.

## **C1.2 Comparisons between the 2006 and 2016 surveys**

Fauna records have been separated and listed per taxa instead of per park. This is to give an overall insight as to species preference of each park and due to the available habitat. Most species detected should be present across all parks.

The park species lists presented in the Biosphere 2006 report appeared to contain species detected during primary surveys, yet also species historically recorded in a several km radius of the parks. To eliminate confusion of a dramatic drop in diversity we have listed comparative tables of each fauna within each park using only the species detected in the primary surveys (2006/2016), as well as suggesting species likely to be present within the parks yet undetected during the surveys.

The purpose of the 2016 study was to recreate another snap-shot of the fauna present in these parks. We have also listed a maximum number of individuals per species detected on any one day, which portrays a comparison to ten years ago. Microbat counts are not listed as they were detected by call recordings so abundance cannot be confirmed.

Detailed 2016 fauna data was submitted to the City of Ryde in a Microsoft Access database, and subsequently all records submitted to the NSW Office of Environment and Heritage Atlas of NSW Wildlife.

## **C2.0 Fauna findings**

The following fauna data has been divided into different taxa (such as reptiles, frogs, mammals, etc.) with tables showing the presence of individual species found in 2006 and/or 2016 surveys, with a summary of each taxa detected across the Ryde reserves. The final column of each table lists maximum number of individuals detected in any 2016 single survey period, representing abundance.

### **C2.1 Reptiles**

Reptile biodiversity across the bushland parks of Ryde surveyed in 2016 has barely changed in the last ten years. All skink species detected are common within urban Sydney and are mostly still present within the parks. Elegant snake-eyed skinks were not detected in several parks during these surveys but are common around brick houses in the area, as these houses have replaced the rocky outcrop habitat once used by these skinks. Weasel skinks and water skinks are common in every park surveyed, and were not previously detected in some parks in 2006. Interestingly, Eastern blue-tongued lizards (*Tiliqua scincoides*) were not detected during any of the surveys, despite persisting in the yards of surrounding suburbia (Koenig *et al.* 2001).

Photographs of a land mullet in Brush Farm Park were submitted by a bush care volunteer in Autumn, but the skink was never detected during surveys. Despite Brush Farm Park providing a suitable rainforest habitat for the skink, their natural populations do not exist south of the Hawkesbury River and it is most certainly an escaped pet or has been released there by a member of the public. It is highly unlikely the individual will survive predation by a dog or cat, or survive the local climate.



Only one dragon species, the Eastern water dragon, persists in the area. Previously, they were present in Brush Farm Park and Field of Mars Reserve, but as this species relies heavily on a permanent water source to avoid predation, the ephemeral nature of Archer Creek has led to the loss of water dragons from Brush Farm Park. Their population appears still healthy in the Field of Mars Reserve, and perhaps they will colonise Darvall Park in the future. The single species of gecko, the broad-tailed gecko, also has a healthy population within the Field of Mars Reserve. This species is another rocky outcrop specialist that has adapted to urbanisation and is found commonly within local houses (Mo 2014), and the abundant outcrops of the Field of Mars Reserve.

Eastern long-necked turtles were detected in the Field of Mars Reserve (as they were previously), but also in Darvall Park. These turtles have been resilient to urbanisation (Stokeld et al. 2013), and populations may continue to grow in the Ryde region if suitable permanent water bodies persist within parks and nest predation from foxes is minimised (Kennett et al. 2009).

The red-bellied black snake was the only snake detected during these surveys, another healthy population within the Field of Mars Reserve. However, due to the cryptic nature of snakes and the survey effort (two days/nights per park per season), it is likely that both other species detected in 2006 (the swamp snake and golden-crowned snake) are still present in the larger parks. These small elapids specialise in feeding on lizards which are abundant within all parks, but will probably be in low densities due to frequent predation by cats and dogs (Shine & Koenig 2001). Based on recent records (Rob Ambrose: Sydney Snake Catchers, pers. comm. 2016) several other species are present in the area and likely to be present in the reserves, especially within the Field of Mars Reserve. These include green tree snakes (*Dendrelaphis punctulatus*), diamond pythons (*Morelia spilota*), and yellow-faced whip snakes (*Demansia psammophis*).

#### Reptiles of Brush Farm Park

Species	Common name	2006	2016	Maximum (2016)
<i>Lampropholis delicata</i>	Garden skink	X	X	13
<i>Lampropholis guichenoti</i>	Grass skink	X	X	3
<i>Eulamprus quoyii</i>	Eastern water skink	X	X	4
<i>Saproscincus mustelinus</i>	Weasel skink	X	X	10
<i>Cryptoblepharus pulcher</i>	Elegant snake-eyed skink	X		-
<i>Intellagama lesueurii</i>	Eastern water dragon	X		-
<i>Hemiaspis signata</i>	Swamp snake	X		-
<i>Bellatorias major</i>	Land mullet		X	1

#### Reptiles of Lambert Park

Species	Common name	2006	2016	Maximum (2016)
<i>Lampropholis delicata</i>	Garden skink	X	X	40
<i>Lampropholis guichenoti</i>	Grass skink	X	X	6
<i>Eulamprus quoyii</i>	Eastern water skink	X	X	4
<i>Saproscincus mustelinus</i>	Weasel skink		X	3
<i>Cryptoblepharus pulcher</i>	Elegant snake-eyed skink	X	X	2

#### Reptiles of Darvall Park

Species	Common name	2006	2016	Maximum (2016)
<i>Lampropholis delicata</i>	Garden skink	X	X	30

Species	Common name	2006	2016	Maximum (2016)
<i>Lampropholis guichenoti</i>	Grass skink	X	X	4
<i>Eulamprus quoyii</i>	Eastern water skink	X	X	7
<i>Saproscincus mustelinus</i>	Weasel skink		X	10
<i>Cryptoblepharus pulcher</i>	Elegant snake-eyed skink	X	X	1
<i>Chelodina longicollis</i>	Eastern long-necked turtle		X	1

#### Reptiles of the Field of Mars Reserve

Species	Common name	2006	2016	Maximum (2016)
<i>Lampropholis delicata</i>	Garden skink	X	X	50
<i>Lampropholis guichenoti</i>	Grass skink	X	X	50
<i>Eulamprus quoyii</i>	Eastern water skink	X	X	6
<i>Saproscincus mustelinus</i>	Weasel skink	X	X	1
<i>Cryptoblepharus pulcher</i>	Elegant snake-eyed skink	X	X	3
<i>Saiphos equalis</i>	Three-toed skink		X	2
<i>Intellagama lesueurii</i>	Eastern water dragon	X	X	5
<i>Phyllurus platurus</i>	Broad-tailed gecko	X	X	11
<i>Hemiaspis signata</i>	Swamp snake	X		-
<i>Cacophis squamulosus</i>	Golden-crowned snake	X		-
<i>Pseudechis porphyriacus</i>	Red-bellied black snake	X	X	2
<i>Chelodina longicollis</i>	Eastern long-necked turtle		X	1

#### Reptiles of Bell Park

Species	Common name	2006	2016	Maximum (2016)
<i>Lampropholis delicata</i>	Garden skink	N/A	X	5
<i>Lampropholis guichenoti</i>	Grass skink	N/A	X	13
<i>Eulamprus quoyii</i>	Eastern water skink	N/A	X	5
<i>Saproscincus mustelinus</i>	Weasel skink	N/A	X	5

#### Reptiles of Outlook Park

Species	Common name	2006	2016	Maximum (2016)
<i>Lampropholis delicata</i>	Garden skink	N/A	X	1
<i>Lampropholis guichenoti</i>	Grass skink	N/A	X	3
<i>Eulamprus quoyii</i>	Eastern water skink	N/A	X	2
<i>Saproscincus mustelinus</i>	Weasel skink	N/A	X	3
<i>Cryptoblepharus pulcher</i>	Elegant snake-eyed skink	N/A	X	2

## C2.2 Frogs

The frog fauna of the Ryde parks has altered over ten years, but overall has improved. The most noticeable change is the presence of Leaf-green tree frogs in each of the parks, except Brush Farm Park. These small frogs are restricted to creeks lined with vegetation (Hoskin et al. 2015). Although not all parks containing these frogs had active flowing creeks, they still provide adequate habitat for populations of leaf-green tree frogs. In some instances (such as in Bell Park and Lambert Park) the frogs were restricted to the small drains and channels of the parks. The ephemeral nature of these channels may not provide breeding habitat for the frogs as no

tadpoles were found, and they have potentially been washed down to these sites from previous rain events. Archer Creek in Brush Farm Park was completely dry during the survey periods which explain the absence of these frogs from the park, and their presence in the more permanent, vegetated creek downstream in Lambert Park.

Peron's tree frogs are now present in every park. These larger frogs shelter in tree hollows and pipes, and are not as reliant on permanent water bodies except for breeding. They are a common garden species of the area. Striped marsh frogs are also present in all parks except Outlook Park. These frogs are extremely common and live in still water bodies, as small as a bird bath or puddle. As Outlook Park currently has no water bodies, it will not support these frogs.

Interestingly, the presence of common Eastern froglets has declined across all parks except the Field of Mars Reserve. This tiny frog is very common across South-Eastern Australian bushland and farmland, but in urban Sydney now appears to be restricted to parks with adjacent bushland. They also live in still water bodies and will breed in puddles. One theory is they may desiccate easily and not disperse so easily due to their small size (Grant Webster, pers. comm. 2016).

Previous surveys (Biosphere 2006) detected less frog diversity and density than current surveys, and suggested frogs will struggle to survive within certain parks until water quality and habitat can be improved. Besides the loss of common Eastern froglets, which may be due to location and not direct habitat factors of the park, the frog fauna has improved in the last ten years.

#### Frogs of Brush Farm Park

Species	Common name	2006	2016	Maximum (2016)
<i>Limnodynastes peronii</i>	Striped marsh frog	X	X	2
<i>Crinia signifera</i>	Common Eastern froglet	X		-
<i>Litoria peronii</i>	Peron's tree frog		X	1
<i>Litoria phyllochroa</i>	Leaf-green tree frog	X		-

#### Frogs of Lambert Park

Species	Common name	2006	2016	Maximum (2016)
<i>Limnodynastes peronii</i>	Striped marsh frog	X	X	1
<i>Crinia signifera</i>	Common Eastern froglet	X		-
<i>Litoria peronii</i>	Peron's tree frog		X	1
<i>Litoria phyllochroa</i>	Leaf-green tree frog		X	3

#### Frogs of Darvall Park

Species	Common name	2006	2016	Maximum (2016)
<i>Limnodynastes peronii</i>	Striped marsh frog	X	X	2
<i>Crinia signifera</i>	Common Eastern froglet	X		-
<i>Litoria peronii</i>	Peron's tree frog		X	1
<i>Litoria phyllochroa</i>	Leaf-green tree frog		X	-

#### Frogs of Field of Mars Reserve

Species	Common name	2006	2016	Maximum (2016)
<i>Limnodynastes peronii</i>	Striped marsh frog	X	X	1

Species	Common name	2006	2016	Maximum (2016)
<i>Crinia signifera</i>	Common Eastern froglet	X	X	5
<i>Litoria peronii</i>	Peron's tree frog	X	X	2
<i>Litoria phyllochroa</i>	Leaf-green tree frog	X	X	20

#### Frogs of Bell Park

Species	Common name	2006	2016	Maximum (2016)
<i>Limnodynastes peronii</i>	Striped marsh frog	N/A	X	5
<i>Litoria peronii</i>	Peron's tree frog	N/A	X	1
<i>Litoria phyllochroa</i>	Leaf-green tree frog	N/A	X	1

#### Frogs of Outlook Park

Species	Common name	2006	2016	Maximum (2016)
<i>Litoria peronii</i>	Peron's tree frog	N/A	X	2

### C2.3 Mammals

The mammal biodiversity of the parks of Ryde has seen some interesting changes in the last ten years. One of the noticeable differences is the substantial increase in ring-tailed possum abundance. Previous surveys detected on average two to three possums of each species per park, yet current surveys indicate at least four times the amount of ring-tailed compared to brush-tailed, and most parks containing upwards of 15 individual ringtails. Ring-tailed possums may have an advantage over brush-tailed possums by living communally, being smaller and therefore requiring less food, and will make their own dreys (nests), not being completely reliant on hollows (Inions 1989). Although ring-tail density has increased in these reserves both possum species densities generally have decreased across their natural range (Friedlander 2007), it would be interesting to see if this is a common trend across Sydney urban remnant bushland parks. Generally brush-tailed possums have been the common urban invader (Matthews et al. 2003), but perhaps it has just taken some more time for ringtails to exploit human resources.

The second noticeable difference is the decline of feral mammal detections. Black rats and dogs were the only species still detected within every park, however all dogs were pets walked by owners and are not residents of the parks and thus would likely have very little impact on resident fauna. Interestingly no house mice were detected at all in 2016. This may be due to their preference for urban habitat and not dense bushland (Lunney 1987). They are probably still present in all of the parks but at low densities. Cats were also detected in only a handful of parks, but are likely present in all parks and not detected due to their cryptic nature. The main decline of feral species has been rabbits and foxes. While both species can be cryptic their evidence is not, diggings and scats are very easy to detect. Rabbits are now only residents in a couple of parks and foxes appear to now be restricted to the Field of Mars Reserve.

Long-nosed bandicoots have now returned to the Field of Mars Reserve and appear to be wide spread across the entire bushland area of the reserve. Despite foxes being a main threat to bandicoot populations (Dexter & Murray 2009; Scott et al. 1999) and being present in the park, the coordinated approach to fox control across Sydney has caused the bandicoot populations to rise, and start to spread back into their former historic range (Chen 2013).

The Field of Mars Reserve contains the highest mammal (and all taxa) diversity not only due to its size and complex of habitats, but also its connectivity to Lane Cove National Park. Although

no other novel native mammals were detected in the Field of Mars Reserve besides a swamp wallaby, it has the most potential to support populations of native bush rats (*Rattus fuscipes*), brown antechinus (*Antechinus stuartii*), Eastern pygmy possums (*Cercertetus nanus*) and sugar gliders (*Petards crevices*).

Generally the bat diversity has increased in the last ten years, including five new species utilising the parks. Common bat species detected in almost every park were grey-headed flying foxes, Eastern bent-wing bats (both threatened species) and Gould's wattled bat. Flying foxes key threatening processes are loss of roosting and foraging sites and conflict with humans. This species was found to be foraging on every site surveyed hence the parks are providing stable resources for the population. Eastern bent-wing bats are cave roosting specialists which have begun to utilise stormwater tunnels/bridges and other human structures, and hunt in forested areas above the canopy. Primary threats to their populations are loss of quality roosting sites and productive foraging habitat.

Unfortunately two species of microbat appear to no longer exist in this area of Ryde, the lesser long-eared bat and the mouse-eared fishing bat. Lesser long-eared bats may not have been detected due to surveys being conducted at a less active time of year but they may also be regionally extinct, their presence was previously very low (Glenn Hoyer, pers. comm. 2016). Fishing bats are a threatened species due to loss or disturbance of roosting sites, and a reduction of stream water quality and fish. Only a single fishing bat was detected ten years ago appearing to be resident of the Lane Cove River valley, foraging occasionally in the Field of Mars Reserve.

*Mormopterus ridei* and *Vespadelus darlingtoni* calls were possibly/probably detected at each park in 2016, however due to their calls being too weak to confidently identify they were not included in the tables. Both species are present in the North Sydney region.

#### Mammals of Brush Farm Park

Species	Common name	2006	2016	Maximum (2016)
<i>Trichosurus vulpecula</i>	Brush-tailed possum	X	X	4
<i>Pseudocheirus peregrinus</i>	Ring-tailed possum	X	X	18
<i>Rattus rattus</i>	Black rat*	X	X	4
<i>Mus musculus</i>	House mouse*	X		-
<i>Canis lupis familiaris</i>	Domestic dog*	X	X	2
<i>Vulpes vulpes</i>	European red fox*	X		-
<i>Felis catus</i>	Domestic cat*	X		-
<i>Oryctolagus cuniculus</i>	European rabbit*	X		-
<i>Pteropus poliocephalus</i>	Grey-headed flying fox	X	X	3
<i>Chalinolobus gouldii</i>	Gould's wattled bat	X	X	-
<i>Nyctophilus geoffroyi</i>	Lesser long-eared bat	X		-
<i>Nyctinomus australis</i>	White-striped mastiff bat	X	X	-
<i>Miniopterus oceanensis</i>	Eastern bent-wing bat		X	-

\* exotic species

#### Mammals of Lambert Park

Species	Common name	2006	2016	Maximum (2016)
<i>Trichosurus vulpecula</i>	Brush-tailed possum	X	X	3
<i>Pseudocheirus peregrinus</i>	Ring-tailed possum		X	7
<i>Rattus rattus</i>	Black rat*	X	X	5

Species	Common name	2006	2016	Maximum (2016)
<i>Canis lupis familiaris</i>	Domestic dog*	X	X	1
<i>Felis catus</i>	Domestic cat*	X	X	1
<i>Oryctolagus cuniculus</i>	European rabbit*		X	1
<i>Pteropus poliocephalus</i>	Grey-headed flying fox	X	X	5
<i>Chalinolobus gouldii</i>	Gould's wattled bat	X		-

\* exotic species

#### Mammals of Darvall Park

Species	Common name	2006	2016	Maximum (2016)
<i>Trichosurus vulpecula</i>	Brush-tailed possum	X	X	3
<i>Pseudocheirus peregrinus</i>	Ring-tailed possum		X	15
<i>Rattus rattus</i>	Black rat*	X	X	2
<i>Mus musculus</i>	House mouse*	X		-
<i>Canis lupis familiaris</i>	Domestic dog*	X	X	3
<i>Felis catus</i>	Domestic cat*	X		-
<i>Oryctolagus cuniculus</i>	European rabbit*		X	1
<i>Pteropus poliocephalus</i>	Grey-headed flying fox	X	X	2
<i>Chalinolobus gouldii</i>	Gould's wattled bat	X	X	-
<i>Nyctinomus australis</i>	White-striped mastiff bat	X		-
<i>Miniopterus oceanensis</i>	Eastern bent-wing bat		X	-

\* exotic species

#### Mammals of Field of Mars Reserve

Species	Common name	2006	2016	Maximum (2016)
<i>Trichosurus vulpecula</i>	Brush-tailed possum	X	X	5
<i>Pseudocheirus peregrinus</i>	Ring-tailed possum	X	X	15
<i>Petards crevices</i>	Sugar glider	X		-
<i>Perameles nasuta</i>	Long-nosed bandicoot		X	2
<i>Wallabia bicolor</i>	Swamp Wallaby		X	1
<i>Ornithorhynchus anatinus</i>	Short-beaked echidna	X		-
<i>Rattus rattus</i>	Black rat*	X	X	6
<i>Mus musculus</i>	House mouse*	X		-
<i>Canis lupis familiaris</i>	Domestic dog*			3
<i>Vulpes vulpes</i>	European red fox*	X	X	1
<i>Felis catus</i>	Domestic cat*	X	X	1
<i>Oryctolagus cuniculus</i>	European rabbit*	X	X	2
<i>Pteropus poliocephalus</i>	Grey-headed flying fox	X	X	2
<i>Chalinolobus gouldii</i>	Gould's wattled bat	X	X	-
<i>Chalinolobus morio</i>	Chocolate wattled bat		X	-
<i>Nyctophilus geoffroyi</i>	Lesser long-eared bat	X		-
<i>Nyctinomus australis</i>	White-striped mastiff bat	X	X	-
<i>Miniopterus oceanensis</i>	Eastern bent-wing bat		X	-
<i>Myotis adversus</i>	Mouse-eared fishing bat	X		-
<i>Vespadelus regulus</i>	Southern forest bat*		X	-
<i>Vespadelus vulturinus</i>	Little forest bat*		X	-

\* exotic species

#### Mammals of Bell Park

Species	Common name	2006	2016	Maximum (2016)
<i>Trichosurus vulpecula</i>	Brush-tailed possum	N/A	X	3

Species	Common name	2006	2016	Maximum (2016)
<i>Pseudocheirus peregrinus</i>	Ring-tailed possum	N/A	X	16
<i>Rattus rattus</i>	Black rat*	N/A	X	1
<i>Canis lupis familiaris</i>	Domestic dog*	N/A	X	2
<i>Felis catus</i>	Domestic cat*	N/A	X	1
<i>Pteropus poliocephalus</i>	Grey-headed flying fox	N/A	X	2
<i>Chalinolobus gouldii</i>	Gould's wattled bat	N/A	X	-
<i>Miniopterus oceanensis</i>	Eastern bent-wing bat	N/A	X	-
<i>Nyctinomus australis</i>	White-striped mastiff bat	N/A	X	-

\* exotic species

#### Mammals of Outlook Park

Species	Common name	2006	2016	Maximum (2016)
<i>Trichosurus vulpecula</i>	Brush-tailed possum	N/A	X	2
<i>Pseudocheirus peregrinus</i>	Ring-tailed possum	N/A	X	6
<i>Rattus rattus</i>	Black rat*	N/A	X	5
<i>Canis lupis familiaris</i>	Domestic dog*	N/A	X	2
<i>Pteropus poliocephalus</i>	Grey-headed flying fox	N/A	X	1
<i>Chalinolobus gouldii</i>	Gould's wattled bat	N/A	X	-
<i>Miniopterus oceanensis</i>	Eastern bent-wing bat	N/A	X	-
<i>Nyctinomus australis</i>	White-striped mastiff bat	N/A	X	-

\* exotic species

#### C2.4 Birds

Over the past ten years the bird diversity and density of the Ryde reserves has seen some noticeable changes, but overall have increased. A total of 76 bird species were detected across all parks in 2016. This is an increase from the 51 species detected during surveys in 2006. Individual birds may be permanent residents of each park, or may move across the parks depending on the size of the park, the habitat structure and flora species, and particular species of bird in question. Although it is safe to assume that almost every bird species detected across the entire area would use each park at some point in the year, with the exception of short-range habitat specialists such as whipbirds and wrens.

There are particular species which well and truly may be gone from the area, such as crested shrike-tits, white-plumed honeyeaters, and the smaller cuckoo species.

There is a definite increase in the number of resident bird species with some permanent new additions. The possible absence of certain species could be due to a change in habitat structure over ten years due to bush regeneration, or more vegetation growth and thus becoming denser (in particular the Field of Mars Reserve), or the presence of two now-abundant aggressive species; the rainbow lorikeet, and the noisy miner. These two species populations have grown significantly in the past couple of decades due to open woodland habitat preference (noisy miner) and increase in urban planting of flowering trees (Smith & Lill 2008; Oldland & Clarke 2007; Hastings & Beattie 2006; White et al. 2005; Grey et al. 1998; Veerman 1991). They live communally, are hyper-aggressive to other small bird species and lorikeets exclude other hollow-nesting birds from all available hollows.

Although the birds detected probably use each park across the area, certain species for preferred particular parks. Darvall Park is full of large old growth Eucalypts offering a wide range

of hollows, thus the canopy is dominated by an abundance of parrots and contains species not detected in the other parks. The southern end of Lambert Park contains a variety of fruiting rainforest trees which provide a food source to support the local bowerbirds, and migrating figbirds, orioles and topknot pigeons. The Field of Mars Reserve depicts a similar trend with bird diversity as it does with all other fauna taxa, due to its size and variety of habitats it supports the most bird diversity and density, as well as providing a corridor for migratory summer species moving across the state.

Another significant trend across the parks is also happening across all of North Sydney; the increase in brush-turkeys. These large birds are slowly recolonising their former historic range due to an increase of fox management, improved bushland and a transition from exotic to native urban gardens (Hoh 2016; Meacham 2015).

Besides the increase in a variety of native species there has also been a noticeable decrease in feral species. Common starlings and house sparrows were not detected at all, and only one Indian myna was detected on the outskirts of a single park. In addition to the sighting of a single blackbird and several red-whiskered bulbuls, feral birds were only found in the smallest and most disturbed parks, as they prefer to live in the urban habitat as opposed to bushland.

Finally, five predatory birds were detected across the reserves. The three diurnal raptors were the brown goshawk, collared sparrowhawk and pacific baza. The south-eastern region of Ryde can easily support these species as the goshawk and sparrowhawk are bird specialists (the dominant food source in the area) and the baza's diet consists primarily of large invertebrates, another well represented food source. The two nocturnal owls were the boobook and the powerful owl. Powerful owls are a threatened species and their population size in Sydney is unknown despite an increase in recent records. The loss of large breeding hollows is a main reason for their decline, yet perhaps with an increase of their primary food source the ring-tailed possum, their offspring survival and population recruitment is increasing around Sydney.

From comparison of number of species recorded in Autumn and Spring, there was a total of 92 species record, with:

- more recorded in Spring (83) than in Autumn (65);
- 57 recorded in Autumn and Spring;
- 9 recorded in Autumn and not Spring; and
- 26 recorded in Spring and not in Autumn.

#### Comparison of species detected in Autumn and Spring

		Spring		
		Absent	Present	Total
Autumn	Absent	0	26	26
	Present	9	57	65
	Total	9	83	92

Comparison of number of species recorded in 2006 and 2016, of the 92 species recorded, there were:

- more recorded in 2016 (83) than in Autumn (65);
- 37 recorded in 2006 and 2016;
- 17 recorded in 2006 and not in 2016; and
- 38 recorded in 2016 and not in 2006.



### Comparison of species detected in 2006 and 2016

		2016		
		Absent	Present	Total
2006	Absent	0	38	38
	Present	17	37	65
	Total	17	83	92

### Birds of Brush Farm Park

Species	Common name	2006	2016	Maximum (2016)
<i>Alectura lathamii</i>	Australian brush-turkey		X	3
<i>Cracticus tibicen tibicen</i>	Australian magpie	X	X	3
<i>Corvus coronoides</i>	Australian raven	X	X	4
<i>Coracina novaehollandiae</i>	Black-faced cuckoo-shrike	X	X	2
<i>Ninox novaeseelandiae boobook</i>	Boobook	X	X	1
<i>Gerygone mouki</i>	Brown gerygone		X	3
<i>Acanthiza pusilla</i>	Brown thornbill	X		-
<i>Scythrops novaehollandiae</i>	Channel-billed cuckoo		X	1
<i>Accipiter cirrocephalus</i>	Collared sparrowhawk		X	1
<i>Falcunculus frontatus</i>	Crested Shrike-tit	X		-
<i>Platycercus elegans</i>	Crimson rosella	X		-
<i>Eopsaltria australis</i>	Eastern yellow robin	X		-
<i>Eudynamis orientalis</i>	Eastern koel	X	X	1
<i>Pachycephala pectoralis</i>	Golden whistler		X	2
<i>Cracticus torquatus</i>	Grey butcherbird	X	X	2
<i>Rhipidura albiscapa</i>	Grey fantail		X	2
<i>Chrysococcyx basalus</i>	Horsfield's bronze-cuckoo	X		-
<i>Microeca fascinans</i>	Jacky winter	X		-
<i>Alisterus scapularis</i>	King parrot		X	2
<i>Dacelo novaeguineae</i>	Laughing kookaburra	X	X	3
<i>Meliphaga lewinii</i>	Lewin's honeyeater		X	1
<i>Anthochaera chrysoptera</i>	Little wattlebird		X	1
<i>Glossopsitta concinna</i>	Musk lorikeet		X	4
<i>Manorina melanocephala melanocephala</i>	Noisy miner	X	X	11
<i>Phylidonyris novaehollandiae</i>	New-Holland honeyeater	X		-
<i>Oriolus sagittatus</i>	Olive-backed oriole	X		-
<i>Grallina cyanoleuca</i>	Peewee		X	2
<i>Strepera graculina</i>	Pied currawong	X	X	3
<i>Ninox strenua</i>	Powerful owl	X	X	2
<i>Trichoglossus haematodus</i>	Rainbow lorikeet		X	10
<i>Pachycephala rufiventris</i>	Rufous whistler	X		-
<i>Anthochaera carunculata</i>	Red wattlebird	X	X	2
<i>Rhipidura rufifrons</i>	Rufous fantail		X	2
<i>Ptilonorhynchus violaceus</i>	Satin bowerbird		X	1
<i>Zosterops lateralis westernensis</i>	Silvereye		X	6
<i>Pardalotus punctatus</i>	Spotted pardalote	X	X	5
<i>Acanthiza lineata</i>	Striated thornbill	X		-
<i>Malurus cyaneus cyanocephalus</i>	Superb fairy-wren		X	4
<i>Podargus strigoides strigoides</i>	Tawny frogmouth		X	1
<i>Hirundo neoxena</i>	Welcome swallow		X	2

Species	Common name	2006	2016	Maximum (2016)
<i>Rhipidura leucophrys</i>	Willy wagtail	X		-
<i>Sericornis frontalis frontalis</i>	White-browed scrubwren	X	X	8

### Birds of Lambert Park

Species	Common name	2006	2016	Maximum (2016)
<i>Alectura lathamii</i>	Australian brush-turkey		X	5
<i>Cracticus tibicen tibicen</i>	Australian magpie	X	X	4
<i>Corvus coronoides</i>	Australian raven	X	X	3
<i>Sphecotheres vieilloti</i>	Australasian figbird		X	2
<i>Scythrops novaehollandiae</i>	Channel-billed cuckoo		X	1
<i>Ocyphaps lophotes</i>	Crested pigeon	X		-
<i>Sturnus vulgaris</i>	Common starling*	X		-
<i>Gallus gallus</i>	Domestic chicken*		X	4
<i>Eopsaltria australis</i>	Eastern yellow robin			-
<i>Platycercus eximius</i>	Eastern rosella	X		-
<i>Eudynamis orientalis</i>	Eastern Koel		X	1
<i>Eolophus roseicapillus</i>	Galah		X	2
<i>Pachycephala pectoralis</i>	Golden whistler		X	1
<i>Cracticus torquatus</i>	Grey butcherbird	X	X	2
<i>Passer domesticus</i>	House sparrow*	X		-
<i>Sturnus tristis</i>	Indian myna*	X		-
<i>Alisterus scapularis</i>	King parrot		X	3
<i>Dacelo novaeguineae</i>	Laughing kookaburra	X	X	2
<i>Glossopsitta concinna</i>	Musk lorikeet		X	2
<i>Manorina melanocephala melanocephala</i>	Noisy miner	X	X	5
<i>Oriolus sagittatus</i>	Olive-backed oriole		X	5
<i>Grallina cyanoleuca</i>	Peewee		X	1
<i>Strepera graculina</i>	Pied currawong	X		4
<i>Trichoglossus haematodus</i>	Rainbow lorikeet	X		7
<i>Anthochaera carunculata</i>	Red wattlebird	X	X	1
<i>Pycnonotus jocosus</i>	Red-whiskered bulbul*	X		-
<i>Rhipidura rufifrons</i>	Rufous fantail		X	2
<i>Ptilonorhynchus violaceus</i>	Satin bowerbird		X	1
<i>Zosterops lateralis westernensis</i>	Silvereye		X	2
<i>Pardalotus punctatus</i>	Spotted pardalote		X	2
<i>Cacatua galerita</i>	Sulphur-crested cockatoo		X	1
<i>Podargus strigoides strigoides</i>	Tawny frogmouth	X	X	2
<i>Lopholaimus antarcticus</i>	Topknot pigeon		X	1
<i>Rhipidura leucophrys</i>	Willy wagtail		X	1
<i>Sericornis frontalis frontalis</i>	White-browed scrubwren		X	4
<i>Ptilotula penicillata</i>	White-plumed honeyeater	X		-

\* feral species

### Birds of Darvall Park

Species	Common name	2006	2016	Maximum (2016)
<i>Corvus coronoides</i>	Australian raven	X	X	3
<i>Cracticus tibicen tibicen</i>	Australian magpie	X	X	3
<i>Threskiornis molucca</i>	Australian white ibis		X	1
<i>Turdus merula</i>	Common blackbird*		X	1
<i>Platycercus elegans</i>	Crimson rosella	X	X	2
<i>Sturnus vulgaris</i>	Common starling*	X		-
<i>Chalcophaps indica</i>	Emerald dove		X	1
<i>Psophodes olivaceus</i>	Eastern whipbird	X	X	1
<i>Platycercus eximius</i>	Eastern rosella	X	X	4
<i>Eudynamys orientalis</i>	Eastern koel		X	1
<i>Acanthorhynchus tenuirostris</i>	Eastern spinebill	X		-
<i>Eolophus roseicapillus</i>	Galah		X	4
<i>Cracticus torquatus</i>	Grey butcherbird		X	4
<i>Alisterus scapularis</i>	King parrot		X	2
<i>Dacelo novaeguineae</i>	Laughing kookaburra		X	4
<i>Cacatua sanguinea</i>	Little corella		X	3
<i>Manorina melanocephala melanocephala</i>	Noisy miner	X	X	25
<i>Oriolus sagittatus</i>	Olive-backed oriole		X	1
<i>Anas superciliosa</i>	Pacific black duck		X	2
<i>Strepera graculina</i>	Pied currawong	X	X	3
<i>Trichoglossus haematodus</i>	Rainbow lorikeet	X	X	50
<i>Zosterops lateralis westernensis</i>	Silvereye		X	5
<i>Cacatua galerita</i>	Sulphur-crested cockatoo	X	X	15
<i>Malurus cyaneus cyanochlamys</i>	Superb fairy-wren		X	4
<i>Podargus strigoides strigoides</i>	Tawny frogmouth		X	2
<i>Sericornis frontalis frontalis</i>	White-browed scrubwren		X	6

\* feral species

### Birds of Field of Mars Reserve

Species	Common name	2006	2016	Maximum (2016)
<i>Alectura lathami</i>	Australian brush-turkey		X	1
<i>Corvus coronoides</i>	Australian raven	X	X	4
<i>Threskiornis molucca</i>	Australian white ibis	X	X	10
<i>Anhinga novaehollandiae</i>	Australian darter	X		-
<i>Coracina novaehollandiae</i>	Black-faced cuckoo-shrike	X	X	2
<i>Monarcha melanopsis</i>	Black-faced monarch		X	4
<i>Ninox novaeseelandiae boobook</i>	Boobook	X	X	1
<i>Gerygone mouki</i>	Brown gerygone		X	2
<i>Accipiter fasciatus</i>	Brown goshawk		X	1
<i>Acanthiza pusilla</i>	Brown thornbill		X	4
<i>Scythrops novaehollandiae</i>	Channel-billed cuckoo	X	X	3
<i>Anas castanea</i>	Chestnut teal	X	X	2
<i>Platycercus elegans</i>	Crimson rosella		X	2
<i>Ocyphaps lophotes</i>	Crested pigeon		X	3
<i>Sturnus vulgaris</i>	Common starling*	X		-
<i>Gallinula tenebrosa</i>	Dusky moorhen		X	2
<i>Eopsaltria australis</i>	Eastern yellow robin	X	X	1

Species	Common name	2006	2016	Maximum (2016)
<i>Acanthorhynchus tenuirostris</i>	Eastern spinebill	X	X	3
<i>Eudynamys orientalis</i>	Eastern koel		X	1
<i>Cacomantis flabelliformis</i>	Fan-tailed cuckoo	X	X	1
<i>Pachycephala pectoralis</i>	Golden whistler		X	2
<i>Cracticus torquatus</i>	Grey butcherbird	X	X	2
<i>Rhipidura albiscapa</i>	Grey fantail	X	X	5
<i>Chrysococcyx basalis</i>	Horsfield's bronze-cuckoo			-
<i>Sturnus tristis</i>	Indian myna*	X		-
<i>Ardea intermedia</i>	Intermediate egret	X		-
<i>Microeca fascinans</i>	Jacky winter	X		-
<i>Alisterus scapularis</i>	King parrot		X	2
<i>Dacelo novaeguineae</i>	Laughing kookaburra	🌐	X	3
<i>Myiagra rubecula</i>	Leaden flycatcher		X	1
<i>Cacatua sanguinea</i>	Little corella		X	2
<i>Anthochaera chrysoptera</i>	Little wattlebird		X	1
<i>Phalacrocorax sulcirostris</i>	Little black cormorant	X		-
<i>Microcarbo melanoleucos</i>	Little pied cormorant	X		-
<i>Cracticus tibicen tibicen</i>	Australian magpie	X	X	3
<i>Glossopsitta concinna</i>	Musk lorikeet		X	4
<i>Nycticorax caledonicus</i>	Nankeen night-heron	X		-
<i>Manorina melanocephala melanocephala</i>	Noisy miner	X	X	25
<i>Philemon corniculatus</i>	Noisy Friarbird	X		-
<i>Phylidonyris novaehollandiae</i>	New-Holland honeyeater	X		-
<i>Oriolus sagittatus</i>	Olive-backed oriole		X	1
<i>Aegotheles cristatus</i>	Owlet nightjar	X		-
<i>Grallina cyanoleuca</i>	Peewee	X	X	1
<i>Aviceda subcristata</i>	Pacific baza		X	1
<i>Anas superciliosa</i>	Pacific black duck	X	X	4
<i>Strepera graculina</i>	Pied currawong	X	X	3
<i>Ninox strenua</i>	Powerful owl	X	X	1
<i>Trichoglossus haematodus</i>	Rainbow lorikeet	X	X	8
<i>Pachycephala rufiventris</i>	Rufous whistler	X		-
<i>Rhipidura rufifrons</i>	Rufous fantail		X	1
<i>Anthochaera carunculata</i>	Red wattlebird	X	X	4
<i>Neochmia temporalis</i>	Red-browed finch	X	X	3
<i>Pycnonotus jocosus</i>	Red-whiskered bulbul	X	X	3
<i>Todiramphus sanctus</i>	Sacred kingfisher	X	X	1
<i>Myzomela sanguinolenta</i>	Scarlet honeyeater		X	1
<i>Zosterops lateralis westernensis</i>	Silvereye		X	20
<i>Ptilonorhynchus violaceus</i>	Satin bowerbird		X	2
<i>Pardalotus punctatus</i>	Spotted pardalote		X	8
<i>Streptopelia chinensis</i>	Spotted turtledove*	X		-
<i>Acanthiza lineata</i>	Striated thornbill		X	3
<i>Malurus cyaneus cyanochlamys</i>	Superb fairy-wren	X	X	5
<i>Cacatua galerita</i>	Sulphur-crested cockatoo	X	X	2
<i>Podargus strigoides strigoides</i>	Tawny frogmouth		X	2
<i>Zosterops lateralis lateralis</i>	Tasmanian silvereye		X	6
<i>Lopholaimus antarcticus</i>	Topknot pigeon		X	3
<i>Malurus lamberti lamberti</i>	Variiegated fairy-wren		X	8

Species	Common name	2006	2016	Maximum (2016)
<i>Hirundo neoxena</i>	Welcome swallow	X	X	1
<i>Rhipidura leucophrys</i>	Willy wagtail	X		-
<i>Sericornis frontalis frontalis</i>	White-browed scrubwren	X	X	4
<i>Ptilotula penicillata</i>	White-plumed honeyeater	X		-
<i>Egretta novaehollandiae</i>	White-faced heron	X		-
<i>Phylidonyris niger</i>	White-cheeked honeyeater		X	2
<i>Melithreptus lunatus</i>	White-naped honeyeater		X	2
<i>Cormobates leucophaea</i>	White-throated treecreeper		X	2
<i>Chenonetta jubata</i>	Wood duck		X	20
<i>Acanthiza nana</i>	Yellow thornbill		X	3
<i>Caligavis chrysops</i>	Yellow-faced honeyeater		X	3
<i>Calyptorhynchus funereus</i>	Yellow-tailed black cockatoo		X	1

\* exotic species

#### Birds of Bell Park

Species	Common name	2006	2016	Maximum (2016)
<i>Alectura lathami</i>	Australian brush-turkey	N/A	X	1
<i>Cracticus tibicen tibicen</i>	Australian magpie	N/A	X	2
<i>Corvus coronoides</i>	Australian raven	N/A	X	4
<i>Gallus gallus</i>	Domestic chicken*	N/A	X	3
<i>Eudynamys orientalis</i>	Eastern koel	N/A	X	1
<i>Cracticus torquatus</i>	Grey butcherbird	N/A	X	5
<i>Sturnus tristis</i>	Indian myna*	N/A	X	2
<i>Alisterus scapularis</i>	King parrot	N/A	X	2
<i>Dacelo novaeguineae</i>	Laughing kookaburra	N/A	X	2
<i>Glossopsitta concinna</i>	Musk lorikeet	N/A	X	2
<i>Manorina melanocephala melanocephala</i>	Noisy miner	N/A	X	12
<i>Grallina cyanoleuca</i>	Peewee	N/A	X	2
<i>Strepera graculina</i>	Pied currawong	N/A	X	3
<i>Ninox strenua</i>	Powerful owl	N/A	X	1
<i>Trichoglossus haematodus</i>	Rainbow lorikeet	N/A	X	8
<i>Anthochaera carunculata</i>	Red wattlebird	N/A	X	1
<i>Columba livia</i>	Rock dove*	N/A	X	10
<i>Zosterops lateralis westernensis</i>	Silvereye	N/A	X	3
<i>Cacatua galerita</i>	Sulphur-crested cockatoo	N/A	X	3
<i>Podargus strigoides strigoides</i>	Tawny frogmouth	N/A	X	1
<i>Sericornis frontalis frontalis</i>	White-browed scrubwren	N/A	X	3
<i>Egretta novaehollandiae</i>	White-faced heron	N/A	X	1

\* exotic species

#### Birds of Outlook Park

Species	Common name	2006	2016	Maximum (2016)
<i>Alectura lathami</i>	Australian brush-turkey	N/A	X	1

Species	Common name	2006	2016	Maximum (2016)
<i>Corvus coronoides</i>	Australian raven	N/A	X	1
<i>Cracticus tibicen tibicen</i>	Australian magpie	N/A	X	1
<i>Eolophus roseicapillus</i>	Galah	N/A	X	3
<i>Cracticus torquatus</i>	Grey butcherbird	N/A	X	1
<i>Alisterus scapularis</i>	King parrot	N/A	X	2
<i>Dacelo novaeguineae</i>	Laughing kookaburra	N/A	X	2
<i>Manorina melanocephala melanocephala</i>	Noisy miner	N/A	X	5
<i>Strepera graculina</i>	Pied currawong	N/A	X	2
<i>Trichoglossus haematodus</i>	Rainbow lorikeet	N/A	X	12
<i>Columba livia</i>	Rock dove*	N/A	X	3
<i>Cacatua galerita</i>	Sulphur-crested cockatoo	N/A	X	2
<i>Malurus cyaneus cyanochlamys</i>	Superb fairy-wren	N/A	X	1
<i>Podargus strigoides strigoides</i>	Tawny frogmouth	N/A	X	1
<i>Sericornis frontalis frontalis</i>	White-browed scrubwren	N/A	X	2

\* exotic species

## C2.5 Fish

Fish diversity of the parks has slightly increased compared to ten years ago. All the same fresh and brackish water fish are still present within the Field of Mars Reserve, and a new species the common galaxias was detected. The Field of Mars Reserve is the only site surveyed which has a permanent freshwater and estuary ecosystem, thus supporting stable populations of fish. The only other site with adequate water bodies is Darvall Park, yet no fish were detected at all. The introduced plague minnow was previously the only fish detected in Darvall Park, yet now the park may not support any fish species. The reason for this is unknown as the water quality is good enough to support healthy frog populations and turtles.

The most significant detections from the fish surveys were the presence of short-finned eels in Brush Farm Park and Lambert Park. Two individuals in each park, both sub-adult and juvenile stage were detected in the only water bodies within the parks, next to the eastern concrete culvert of Brush Farm Park and the southern culvert of Lambert Park. Although relatively common in freshwater habitats of East-coast Australia, the presence of a predator in these water bodies indicates a supportable ecosystem and adequate food source.

### Fish of Brush Farm Park

Species	Common name	2006	2016	Maximum (2016)
<i>Anguilla australis</i>	Short-finned eel		X	2 elvers

### Fish of Lambert Park

Species	Common name	2006	2016	Maximum (2016)
<i>Anguilla australis</i>	Short-finned eel		X	2 sub-adults

### Fish of Darvall Park

Species	Common name	2006	2016	Maximum (2016)
<i>Gambusia holbrooki</i>	Plague minnow*	X		-

\* exotic species

### Fish of the Field of Mars Reserve

Species	Common name	2006	2016	Maximum (2016)
<i>Gambusia holbrooki</i>	Plague minnow*	X	X	7
<i>Gobiomorphus australis</i>	Striped gudgeon	X	X	2
<i>Hypseleotris galii</i>	Firetail gudgeon	X	X	2
<i>Philypnodon macrostomus</i>	Dwarf flathead gudgeon	X	X	2
<i>Trachystoma petardi</i>	Freshwater mullet	X	X	6
<i>Galaxias maculatus</i>	Common galaxias		X	4
<i>Anguilla australis</i>	Short-finned eel	X	X	1

\* exotic species

## C2.6 Invertebrate fauna

### C2.6.1 Comparability with previous (Biosphere 2006) results

While a variety of invertebrate fauna were detected across the reserves in the 2006 survey, the level of taxonomic resolution provided in this previous survey (identification only to family or higher levels of taxonomic hierarchy) means that comparison between survey periods is difficult. Additionally, the identification of each group as either occurring or not occurring provides little information on the diversity of species within each group. As such, making conclusions regarding the extent or trajectory of change over time since 2006 is not possible.

It should be noted that the total diversity of invertebrates within a reserve will in all cases be much larger than can be detected in a single survey. It is likely that a very large number of invertebrate groups are represented within all of the reserves, and that the total number of species within each group is much larger than what was detected in both the 2006 and 2016 surveys. Detecting a greater portion of these species would require employing a range of group-specific survey techniques, for example pitfall trapping for ant, spider or other ground-dwelling invertebrate species or yellow pan trapping for diurnal flying insects. While these additional techniques would yield a greater diversity of detected species, they are beyond the scope of the previous (Biosphere 2006) investigation upon which this current survey is based.

### C2.6.2 Habitat availability in Ryde reserves

Overall, the greatest diversity of invertebrate species was found in Brush Farm Park and Field of Mars reserves, largely due to the broad range of different habitats available for invertebrate species, and the overall larger areas of these reserves. Invertebrate species generally show strong reliance on the particular narrow suite of microhabitat traits that they are adapted for, and a broad range of these specific habitats are available within the surveyed reserves. The specific habitat values of some of these different microhabitat habitat elements are discussed below.

#### C2.6.2.1 Ground-layer vegetation

A variety of low habitats were present across the reserves. Of these, native grasslands of Basket Grass (*Opismenus aemulus* and *O. imbecilis*) and Weeping Grass (*Microlaena stipoides*) supported a great diversity of species. These areas provide ideal habitat for active hunting spiders such as Lynx Spiders (family Oxyopidae) and Jumping Spiders (family Salticidae). Also, a large diversity of moth species was present within these low grasslands, even during diurnal sampling. This grassland habitat was available in all reserves; however the greatest extents were found in Brush Farm Park, Darvall Park and the Field of Mars reserve.

Other low vegetation provides habitat for invertebrate species. For example, flowering shrubs and forbs provide nectar resources for butterflies and other nectar feeding insect species such as bees (family Apidae) and hover flies (family Syrphidae). In turn, these floral resources provide ideal locations for sit-and-wait predators such as Crab Spiders and Flower Spiders (family Thomisidae).

#### **C2.6.2.2 Shrubs and mid-storey vegetation**

Habitat complexity in the mid-storey and lower canopy provides locations for web-building spiders. Particularly, a variety of Orb-weaving Spiders in the family Araneidae (particularly *Araneus* sp. and *Eriophora* sp.), Leaf-curling Spiders (*Phonognatha* sp.) and Net-casting spiders (family Deinopidae) were abundant in these habitats across all Ryde reserves at the time of survey.

#### **C2.6.2.3 Leaf litter and debris**

Leaf litter and other ground-layer debris provide a range of habitat, particularly wetter areas under dense Eucalyptus canopy. Species supported by this microhabitat type include Springtails (families Isotomidae and Sminthuridae), Earthworms (Order Megadrilaceae), Scolopendran Centipedes (family Scolopendridae), Slaters (family Oniscidea), Amphipods (family Talpidae), Mites (family Trombididae), Ground beetles (family Carabidae), Termites (family Termitidae), and Cockroaches (Family Blattidae). A range of more generalised ground-dwelling groups such as ants (family Formicidae), Wolf Spiders (family Lycosidae) and Ant-hunting Spiders (family Zodariidae) are also abundant within leaf litter. All Ryde reserves sampled provided some areas of leaf litter habitat.

#### **C2.6.2.4 Aquatic habitats**

The availability of aquatic habitat within the reserves during the current investigation was low overall, with dry conditions prior to the survey reducing the extent of available sampling sites. However, a number of invertebrate groups were still detected, including larvae of Dragonflies and/or Damselflies (Odonata), Water Treaders (Hemiptera – Veliidae) and Back Swimmers (Hemiptera – Notonectidae).

#### **C2.6.2.5 *Eucalyptus* and *Acacia* spp. foliage**

The availability of low foliage of *Acacia* and *Eucalyptus* species provided habitat for a diverse range of herbivorous and sap-sucking insects. Particularly, a variety of Leaf Beetles (family Chrysomelidae), Lady Beetles (*Coccinella* sp.), Leaf Hoppers (family Cicadellidae), Aphids (family Aphidae) and a variety of Scale Insects (superfamily Coccoidea) are abundant in this microhabitat type, particularly when fresh growth is available. Other groups associated with these habitats include 'Wattle Pig' (*Leptopius* sp.) and other Weevils in the family Curculionidae), Scarab Beetles (family Scarabidae), Clerid Beetles (family Cleridae) and Long Horned Beetles (family Cerambycidae). Additionally, ant species in the subfamily Dolichoderinae (*Anonychomyrma* sp. and *Iridomyrmex* sp.) are associated with this habitat as they provide tending services to Scale Insects and Aphids.

The presence of low *Acacia* sp. and *Eucalyptus* sp. allowed for easy hand collection of a variety of the above mentioned groups. While all reserves provided *Eucalyptus* spp. habitat, this was not always readily sampled as it was in some cases above the reach of the surveyor. Overall, these habitats were most abundant in Brush Farm Park, Darvall Park and Field of Mars reserve.



### C2.6.2.6 Tree trunks

Tree species, particularly those with flaking bark such as Sydney Blue Gum (*Eucalyptus saligna*), Flooded Gum (*Eucalyptus grandis*) or Blackbutt (*E. pilularis*), provide important habitat for a range of species. Particularly, Bark Hoppers (family Eurybrachyidae, especially *Platybachrys* sp.), Bark Cockroaches (family Blaberidae), arboreal ants such as *Crematogaster* sp., and a range of predatory spiders are associated with these habitats.

### C2.6.3 Invertebrates recorded in the parks

#### Invertebrates recorded in Brush Farm Park

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
Annelida - Clitellata						
		<i>Hirudinea</i>		Leeches	X	
Annelida - Oligochaeta						
		<i>Megadrilacea</i>		Earthworms	X	1
Arthropoda - Arachnida						
		<i>Araneae</i>				
		Araneidae	<i>Argiope keyserlingi</i>	St Andrew's Cross Spider		1
		Araneidae	<i>Demadiana</i> sp.	Orb-weaving Spiders		1
		Araneidae	<i>Eriophora</i> sp.	Orb-weaving Spiders		1
		Araneidae	<i>Phonognatha</i> sp.	Leaf Curling Spiders		1
		Araneidae		orb-weavers	X	6
		Lycosidae		wolf spiders	X	
		Nephidae	<i>Nephila plumipes</i>	Golden Orb-weaving Spiders		1
		Oxyopidae	<i>Oxyopes</i> sp.	Lynx Spiders		3
		Salticidae		Jumping Spiders		3
		Tetragnathidae		Long-jawed spiders	X	
		Theridiidae		Scaffold Web Weavers		1
		Thomsidae	<i>Diaea</i> sp.	Flower Spiders		1
		Thomsidae	<i>Diaea</i> sp.	Flower Spiders		1
		Thomsidae	<i>Tmarus</i> sp.	Crab Spiders		1
		Thomsidae		Crab Spiders		4
		<i>Trombidiformes</i>				
		Trombidiidae		Mites	X	1
Arthropoda - Chilopoda						
		<i>Scolopendromorpha</i>				
		Scolopendridae		Centipedes	X	
		Scolopendridae	<i>Scolopendra</i> sp.	Centipedes		1
Arthropoda - Diplopoda						
				Millipedes	X	
		<i>Polydesmida</i>		Polydesmid Millipedes		1

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
Arthropoda - Entognatha						
				Springtails	X	
		<i>Entomobryomorpha</i>				
		Isotomidae		Springtails		1
		<i>Symphyleona</i>				
		Smithuridae		Springtails		1
Arthropoda - Insecta						
		<i>Blattodea</i>				
		Blattidae		Cockroaches	X	1
		Blaberidae	<i>Laxta</i> sp.	Bark Cockroaches		1
		Blaberidae - Panesthiinae		Wood Cockroaches		1
		Blattellidae	<i>Ellipsidon</i> sp.	Bush Cockroaches		1
		Blattidae	<i>Polyzosteria</i> sp.	Coackroaches		1
		Termitidae		Termites	X	
		<i>Coleoptera</i>				
		Buprestidae		Jewel Beetles	X	
		Carabidae		Ground Beetles	X	1
		Cerambycidae		Long-horned beetles	X	
		Chrysomelidae		Leaf Blisters	X	
		Chrysomelidae - Chrysomelinae	<i>Calomera ruficeps</i>	Metallic Green Acacia Beetle		1
		Chrysomelidae - Chrysomelinae	<i>Dicranosterna</i> sp.	Acacia Leaf Beetle		1
		Chrysomelidae - Chrysomelinae	<i>Lamprolina</i> sp.	Pittosporum Beetles		1
		Chrysomelidae - Criocerinae		Narrow-necked Leaf Beetles		1
		Chrysomelidae - Cryptocephalinae		Leaf Cylinder Beetles		2
		Cincindelidae		Tiger Beetles	X	
		Circulionidae	<i>Leptopius</i> sp.	Wattle Pig Weevils		1
		Cleridae		Clerid Beetles		1
		Coccinellidae		Labybirds	X	
		Coccinellidae	<i>Cocinella</i> sp.	Ladybirds		1
		Dytiscidae		Diving Beetles	X	
		Elateridae		Click Beetles	X	
		Scarabaeidae		Scarab Beetles	X	
		Staphylinidae		Rove Beetles	X	
		<i>Dermaptera</i>				
				Earwigs	X	
		<i>Diptera</i>				
		Assilidae		Robber Flies	X	
		Calliphoridae		Blow Flies	X	1

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
		Culicidae		Mosquitoes	X	
		Culicidae	<i>Culex</i> sp.	Mosquitoes		3
		Lauxaniidaae		Lauxaniid Flies		1
		Muscidae		House Flies	X	1
		Mycetophilidae		Fungus Gnats		1
		Sciaridae		Fungus Gnats		1
		Syrphidae		Hover Flies	X	1
		Tabanidae		March Flies	X	
		Tipulidae	<i>Tipula</i> sp.	Crane Flies		2
		Tipulidae		Crane Flies	X	
				Flies		2
	<i>Diptera - Nematocera</i>					
				Fungus Gnats and Midges		3
	<i>Ephemeroptera</i>					
				Mayflies	X	
	<i>Hemiptera</i>					
		Alydidae	<i>Melanacanthus scutellaris</i>	Small Brown Bean Bug		1
		Aphididae		Aphids	X	2
		Cicadellidae		Leaf Hoppers	X	
		Cicadellidae		Leafhoppers		3
		Cicadidae		Cicadas	X	
		Coccoidea	<i>Ceroplastes</i> sp.	Scale Insects		1
		Coccoidea		Scale Insects	X	3
		Eurybrachyidae	<i>Platybrachys</i> sp.	Bark Hoppers		1
		Flatidae		Plant Hoppers		1
		Gerridae		Water Striders	X	
		Lygaeidae		Ground Bugs	X	
		Miridae		Mirid Bugs		1
		Notonectidae		Backswimmers	X	
		Pentatomidae		Shield Bugs	X	
		Pentatomidae	<i>Poecilometis patruelis</i>	Common Gum Tree Shield Bug		1
		Reduviidae		Assassin Bugs	X	
		Reduviidae	<i>Pristhesancus plagipennis</i>	Common Assassin Bug		1
	<i>Hymenoptera</i>					
		Apidae		Bees	X	
		Formicidae		Ants	X	
		Formicidae - Dolichoderinae	<i>Iridomyrmex</i> sp.	Tyrant Ants		1
		Formicidae - Ectatominae	<i>Rhytidoponera metallica</i>	Greed-headed Ant		1

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
		Formicidae - Formicinae	<i>Polyrachis</i> sp.	Ants		1
		Formicidae - Myrmicinae	<i>Creumatogaster</i> sp.	Ants		2
		Formicidae - Myrmicinae	<i>Monomorium</i> sp.	Ants		1
		Formicidae - Myrmicinae	<i>Myrmecia</i> sp.	Bull Ants		1
		Ichneumonidae		Ichneumonid Wasps		2
		Ichneumonoidea (superfamily)		Parasitic Wasps	X	
		Mymaridae		Fairy Flies		3
		Pergidae		Sawflies		2
		Sphecidae		Sphecid Wasps	X	
		Vespidae		Vespid Wasps	X	
	<i>Lepidoptera</i>					
		Arctidae		Tiger Moths	X	
		Danainae (subfamily)		Milkweed Butterflies	X	
		Geometridae		Geometer Moths	X	
		Hesperiidae		Skippers	X	
		Lycaenidae		Ant Blue Butterflies	X	
		Nymphalidae		Swallowtail Butterflies	X	
		Psyshidae		Case Moths		1
		Sesiidae		Clearwing Moths	X	
		Sphingidae		Hawk Moths	X	
				Moths		11
	<i>Mantodea</i>					
		Mantidae		Praying Mantis	X	1
	<i>Neuroptera</i>					
				Lace Wings	X	1
	<i>Odonata</i>					
		Anisoptera (suborder)		Dragonflies	X	
		Zygoptera		Damselflies	X	
	<i>Orthoptera</i>					
		Acrididae		Grasshoppers	X	1
		Gryllacridae		Tree Crickets	X	
		Gryllidae		Crickets	X	
		Gryllotalpidae		Mole Crickets	X	
		Tettigoniidae		Katydid	X	
	<i>Plecoptera</i>					
		Plecoptera (order)		Stoneflies	X	

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
	<i>Trichoptera</i>			Caddisflies		1
Arthropoda - Malacostraca						
	<i>Amphipoda</i>					
		Taltridae		Amphipods		1
	<i>Isopoda</i>					
		Oniscidea (suborder)		Slaters	X	1
		Oniscidea	<i>Armadillidium vulgare</i>	Slaters		1
		Oniscidea	<i>Porcellio scaber</i>	Slaters		1
Molusca - Gastropoda						
	<i>Helicoidea</i>					
		Helicidae		Snails	X	
	<i>Parmacelloidea</i>					
		Milacidae		Slugs	X	
	<i>Pulmonata</i>					
				Land Snails		1

#### Invertebrates recorded in Lambert Park

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
Annelida - Oligochaeta						
	<i>Megadrilacea</i>					
				Earthworms	X	1
Arthropoda - Arachnida						
	<i>Araneae</i>					
		Araneidae		Orb-weavers	X	
		Araneidae	<i>Araneus</i> sp.	Angulate Orb-weaving Spiders		4
		Araneidae	<i>Argiope keyserlingi</i>	St Andrew's Cross Spider		1
		Araneidae	<i>Eriophora</i> sp.	Orb-weaving Spiders		1
		Araneidae	<i>Phonognatha</i> sp.	Leaf Curling Spiders		1
		Deinopidae	<i>Deinopis</i> sp.	Net-casting Spiders		1
		Lycosidae		Wolf spiders	X	
		Miturgidae		Prowling Spiders		2
		Oxyopidae	<i>Oxyopes</i> sp.	Lynx Spiders		1
		Salticidae		Jumping Spiders		2
		Theridiidae	<i>Argyroides</i> sp.	Jewdrop Spiders		1

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
		Thomisidae	<i>Diaea</i> sp.	Crap Spiders		1
		Thomisidae	<i>Sidymella</i> sp.	Crap Spiders		1
				Spiders		2
	<i>Trombidiformes</i>					
		Trombidiidae		Mites, ticks	X	
Arthropoda - Chilopoda						
	<i>Scolopendromorpha</i>					
		Scolopendridae		Centipedes	X	
		Scolopendridae	<i>Scolopendra</i> sp.	Centipedes		1
Arthropoda - Diplopoda						
	<i>Polydesmida</i>					
				Polydesmid Millipedes		1
Arthropoda - Entognatha						
				Springtails	X	
	<i>Entomobryomorpha</i>					
		Isotomidae		Springtails		1
	<i>Symphyleona</i>					
		Smithuridae		Springtails		1
Arthropoda - Insecta						
	<i>Blattodea</i>					
		Blattidae		Cockroaches	X	1
		Termitidae		Termites	X	
	<i>Coleoptera</i>					
		Carabidae		Ground Beetles	X	
		Cerambycidae		Long-horned beetles	X	
		Chrysomelidae - Chrysomelinae		Leaf Beetles		1
		Cleridae		Clerid Beetles		1
		Coccinellidae		Ladybirds	X	
		Coccinellidae	<i>Coccinella</i> sp.	Ladybirds		1
		Cryptophagidae		Fungus Beetles		1
		Dytiscidae		Diving Beetles	X	
		Elateridae		Click Beetles	X	
		Scarabaeidae		Scarab Beetles	X	
		Staphylinidae		Rove Beetles		1
	<i>Diptera</i>					
		Calliphoridae		Blow Flies	X	1
		Culicidae		Mosquitoes	X	
		Culicidae	<i>Culex</i> sp.	Mosquitoes		2
		Muscidae		House Flies	X	

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
		Mycetophilidae		Fungus Gnats		1
		Phoridae		Phorid Flies		1
		Pipunculidae		Big-headed Flies		1
		Syrphidae		Hover Flies	X	
		Tipulidae		Crane Flies	X	
		Tipulidae	<i>Tipula sp.</i>	Crane Flies		1
				Flies		3
	<i>Hemiptera</i>					
		Aphididae		Aphids	X	1
		Cicadellidae		Leaf Hoppers	X	
		Cicadidae		Cicadas	X	
		Coccoidea		Scale Bugs	X	1
		Coccoidea	<i>Ceroplastes sp.</i>	Scale Insects		1
		Derbidae		Plant Hoppers		1
		Eurybrachyidae	<i>Platybrachys sp.</i>	Bark Hoppers		1
		Flatidae		Plant Hoppers		1
		Gerridae		Water Striders	X	
		Pentatomidae		Shield Bugs	X	
		Pentatomidae	<i>Alcaeus sp.</i>	Shield Bugs		1
		Pentatomidae	<i>Theseus sp.</i>	Shield Bugs		1
		Reduviidae		Assassin Bugs	X	
		Reduviidae	<i>Pristhesancus plagipennis</i>	Common Assassin Bug		1
	<i>Hymenoptera</i>					
		Apidae		Bees	X	
		Formicidae		Ants	X	
		Formicidae - Dolichoderinae	<i>Ochetellus sp.</i>	Ants		1
		Formicidae - Formicinae	<i>Notoncus capitatus</i>	Ants		1
		Formicidae - Formicinae	<i>Nylanderia sp.</i>	Ants		1
		Formicidae - Formicinae	<i>Polyrachis sp.</i>	Ants		1
		Formicidae - Myrmicinae	<i>Crematogaster sp.</i>	Ants		1
		Ichneumonidae		Ichneumonid Wasps		1
		Vespidae		Vespid Wasps	X	1
	<i>Lepidoptera</i>					
		Nymphalidae - Danainae		Milkweed Butterflies	X	
		Hesperiidae		Skippers	X	
		Lycaenidae		Ant Blue	X	

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
				Butterflies		
		Notodontidae - Thaumetopoeinae		Processionary Caterpillars		1
		Nymphalidae	<i>Euploea core</i>	Common Crow Butterfly		1
		Nymphalidae	<i>Vanessa kershawi</i>	Australian Painted Lady Butterfly		1
		Psyshidae		Case Moths		1
		Sphingigae		Hawk Moths	X	
				Moths		6
	<i>Mantodea</i>					
		Mantidae		Praying mantis		1
	<i>Neuroptera</i>					
				Lacewings		1
	<i>Odonata</i>					
		Anisoptera (suborder)		Dragonflies	X	
	<i>Orthoptera</i>					
		Acrididae		Grasshoppers	X	1
		Gryllacridae		Tree Crickets	X	
		Gryllidae		Crickets	X	
		Gryllidae		True Crickets		1
		Gryllotalpidae		Mole Crickets	X	
		Tettigoniidae		Katydids		1
	<i>Trichoptera</i>					
				Caddisflies		2
Arthropoda - Malacostraca						
	<i>Amphipoda</i>					
		Taltridae		Amphipods		1
	<i>Isopoda</i>					
		Oniscidea	<i>Armadillidium vulgare</i>	Slaters		1
				Slaters		1
Molusca - Gastropoda						
	<i>Helicoidea</i>					
		Helicidae		Snails	X	1
	<i>Parmacelloidea</i>					
		Milacidae		Slugs	X	

#### Invertebrates recorded in Darvall Park

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
Annelida - Oligochaeta						



Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
		<i>Megadrilacea</i>		Earthworms	X	1
Arthropoda - Arachnida						
		<i>Araneae</i>				
		Araneidae		orb-weavers	X	
		Araneidae	<i>Araneus</i> sp.	Orb-weaving Spiders		1
		Araneidae	<i>Argiope keyserlingi</i>	St Andrew's Cross Spider		1
		Araneidae	<i>Phonognatha</i> sp.	Leaf Curling Spiders		1
		Deinopidae	<i>Deinopis</i> sp.	Net-casting Spiders		1
		Lycosidae		wolf spiders	X	
		Oxyopidae	<i>Oxyopes</i> sp.	Lynx Spiders		3
		Salticidae		Jumping Spiders		3
		Tetragnathidae		Stretch Spiders		1
		Theridiidae		Scaffold-web Weavers		1
		Thomisidae		Crab Spiders		1
		Zodariidae	<i>Storena</i> sp.	Ant-hunting Spiders		1
		<i>Trombidiformes</i>				
		Trombidiidae		Mites	X	1
Arthropoda - Chilopoda						
		<i>Scolopendromorpha</i>				
		Scolopendridae		Centipedes	X	
		Scolopendridae	<i>Scolopendra</i> sp.	Centipedes		1
Arthropoda - Entognatha						
				Springtails	X	
		<i>Symphyleona</i>				
		Smithuridae		Springtails		1
Arthropoda - Insecta						
		<i>Blattodea</i>				
		Blattidae		Coackroaches	X	1
		Termitidae		Termites	X	
		<i>Coleoptera</i>				
		Carabidae		Ground Beetles	X	1
		Cerambycidae		Long-horned beetles	X	
		Chrysomelidae - Chrysomelinae	<i>Calomera ruficeps</i>	Metallic Green Acacia Beetle		1
		Coccinellidae		Labybirds	X	
		Coccinellidae	<i>Cocinella</i> sp.	Ladybirds		1

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
		Dytiscidae		Diving Beetles	X	
		Elateridae		Click Beetles	X	
		Scarabaeidae		Scarab Beetles	X	
		Staphylinidae		Rove Beetles	X	1
	<i>Dermaptera</i>					
				Earwigs	X	
	<i>Diptera</i>					
		Assilidae		Robber Flies	X	
		Calliphoridae		Blow Flies	X	
		Culicidae	<i>Culex</i> sp.	Mosquitoes		1
		Culicidae	<i>Culex</i> sp.	Mosquitoes		1
		Culicidae	<i>Culex</i> sp.	Mosquitoes		1
		Culicidae	<i>Culex</i> sp.	Mosquitoes		1
		Culicidae		Mosquitoes	X	
		Muscidae		House Flies	X	
		Syrphidae		Hover Flies	X	
		Tabanidae		March Flies	X	
		Tipulidae (larvae)		Crane Flies		1
	<i>Diptera - Nematocera</i>					
				Fungus Gnats and Midges		5
	<i>Hemiptera</i>					
		Alydidae	<i>Noliphus</i> sp.	Board-headed Bugs		1
		Aphididae		Aphids	X	1
		Cicadellidae		Leaf Hoppers	X	
		Cicadellidae		Leafhoppers		2
		Cicadellidae		Treehoppers		2
		Cicadidae		Cicadas	X	
		Cixiidae		Cixiid Planthoppers		1
		Coccoidea		Scale Insects	X	3
		Eurybrachyidae	<i>Platybrachys</i> sp.	Bark Hoppers		1
		Gerridae		Water Striders	X	
		Lygaeidae		Ground Bugs	X	
		Membracidae	<i>Sextius</i> sp.	Horned Treehoppers		1
		Miridae		Mirid Bugs		1
		Notonectidae		Backswimmers	X	
		Pentatomidae		Shield Bugs	X	
		Pentatomidae	<i>Poecilometis patruelis</i>	Common Gum Tree Shield Bug		1

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
		Reduviidae		Assassin Bugs	X	
		Reduviidae	<i>Pristhesancus plagipennis</i>	Common Assassin Bug		1
		Scutelleridae		Shield Backed Bugs		1
		Veliidae		Water Treaders		1
	<i>Hymenoptera</i>					
		Apidae		Bees	X	
		Apidae - Apinae	<i>Apis mellifera</i>	European Honey Bee		1
		Braconidae		Braconid Wasps		1
		Formicidae		Ants	X	
		Formicidae - Dolichoderinae	<i>Anonychomyrma sp.</i>	Ants		1
		Formicidae - Dolichoderinae	<i>Iridomyrmex chasei</i>	Tyrant Ants		1
		Formicidae - Dolichoderinae	<i>Leptomymex sp.</i>	Spider Ants		1
		Formicidae - Ectatominae	<i>Rhytidoponera metallica</i>	Greed-headed Ant		1
		Formicidae - Formicinae	<i>Notoncus capitatus</i>	Ants		1
		Formicidae - Formicinae	<i>Nylanderia sp.</i>	Ants		1
		Formicidae - Formicinae	<i>Polyrachis sp.</i>	Ants		1
		Formicidae - Myrmiicinae	<i>Myrmecia sp.</i>	Jumping Ants		1
		Ichneumonoidea (superfamily)		Parasitic Wasps	X	
		Ichneumonidae - Ichneumoninae		Ichneumon Wasps		1
		Mymaridae		Fairy Flies		2
		Pergidae		Sawflies		2
		Vespididae		Vespid Wasps	X	
	<i>Lepidoptera</i>					
		Danainae (subfamily)		Milkweed Butterflies	X	
		Geometridae		Geometer Moths	X	
		Hesperiidae		Skippers	X	1
		Nymphalidae		Swallowtail Butterflies	X	

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
		Nymphalidae	<i>Euploea core</i>	Common Crow Butterfly		1
		Psyshidae		Case Moths		1
		Sesiidae		Clearwing Moths	X	
		Sphingigae		Hawk Moths	X	
				Moths		18
	<i>Mantodea</i>					
		Mantidae		Praying Mantis	X	
	<i>Neuroptera</i>					
				Lacewings		1
	<i>Odonata - Anisoptera</i>					
				Dragonflies	X	1
	<i>Orthoptera</i>					
		Acrididae		Grasshoppers	X	
		Gryllacridae		Tree Crickets	X	
		Gryllidae		True Crickets	X	1
		Gryllotalpidae		Mole Crickets	X	
		Tettigoniidae		Katydids	X	1
	<i>Plecoptera</i>					
				Stoneflies	X	
	<i>Trichoptera</i>					
				Caddisflies		1
Arthropoda - Malacostraca						
	<i>Amphipoda</i>					
		Taltridae		Amphipods		1
	<i>Isopoda</i>					
		Oniscidea	<i>Armadillidium vulgare</i>	Slaters		1
Molusca - Gastropoda						
	<i>Helicoidea</i>					
		Helicidae		Snails	X	
	<i>Parmacelloidea</i>					
		Milacidae		Slugs	X	
	<i>Pulmonata</i>					
		Planorbidae		Planorbid Snails		2

**Invertebrates recorded in the Field of Mars Reserve**

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
Annelida - Clitellata						
	<i>Hirudinea</i>					
				Leeches	X	
Annelida - Oligochaeta						

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
		<i>Megadrilacea</i>		Earthworms	X	1
Arthropoda - Arachnida						
		<i>Araneae</i>				
		Araneidae		Orb-weaving Spiders	X	5
		Araneidae	<i>Araneus sp.</i>	Orb-weaving Spiders		3
		Araneidae	<i>Argiope keyserlingi</i>	St Andrew's Cross Spider		1
		Araneidae	<i>Eriophora sp.</i>	Orb-weaving Spiders		1
		Araneidae	<i>Phonognatha sp.</i>	Leaf Curling Spiders		1
		Deinopidae	<i>Deinopis sp.</i>	Net-casting Spiders		1
		Lycosidae		Wolf spiders	X	1
		Oxyopidae	<i>Oxyopes sp.</i>	Lynx Spiders		3
		Salticidae		Jumping Spiders		7
		Tetragnathidae	<i>Leucage granulata</i>	Silver Orb Spider		1
		Tetragnathidae		Stretch Spiders	X	2
		Theridiidae	<i>Argyrodes sp.</i>	Scaffold-web Weavers		1
		Theridiidae		Scaffold-web Weavers		1
		Thomisidae	<i>Phrynarachne sp.</i>	Crab Spiders		1
		Thomisidae	<i>Sidymella sp.</i>	Crab Spiders		1
		Thomisidae		Crab Spiders		3
		Zodariidae	<i>Storena sp.</i>	Ant-hunting Spiders		1
		Zodariidae		Ant-hunting Spiders		2
		<i>Araneae - Mygalophae</i>				
		Idiopidae		Trapdoor Spiders		1
		<i>Trombidiformes</i>				
		Trombidiidae		Mites	X	2
Arthropoda - Chilopoda						
		<i>Scolopendromorpha</i>				
		Scolopendridae		Centipedes	X	
		Scolopendridae	<i>Scolopendra sp.</i>	Centipedes		1
Arthropoda - Diplopoda						
				Millipedes	X	

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
	<i>Polydesmida</i>					
				Polydesmid Millipedes		1
	<i>Spirobolida</i>					
				Spirobold millipede		1
Arthropoda - Entognatha						
				Springtails	X	
	<i>Entomobryomorpha</i>					
		Isotomidae		Springtails		3
	<i>Symphyleona</i>					
		Smithuridae		Springtails		4
Arthropoda - Insecta						
	<i>Blattodea</i>					
		Blaberidae		Bark Cockroaches		1
		Blattidae		Cockroaches	X	4
		Termitidae	<i>Nasutitermes sp.</i>	Termites		1
		Termitidae		Termites	X	
	<i>Coleoptera</i>					
		Buprestidae		Jewel Beetles	X	
		Carabidae		Ground Beetles	X	2
		Cerambycidae		Long-horned beetles	X	
		Chrysomelidae		Leaf Blisters	X	
		Chrysomelidae - Cryptocephalinae		Leaf Cylinder Beetles		2
		Cincindelidae		Tiger Beetles	X	
		Cleridae		Clerid Beetles		1
		Coccinellidae		Labybirds	X	
		Coccinellidae	<i>Cocinella sp.</i>	Ladybirds		3
		Curculionidae		Weevils	X	
		Dytiscidae		Diving Beetles	X	
		Elateridae		Click Beetles	X	
		Gyrrinidae		Whirligig beetles	X	
		Scarabaeidae		Scarab Beetles	X	
		Scarabaeidae		Cockchafers		3
		Silphidae		Carrion Beetles	X	
		Staphylinidae		Rove Beetles	X	
		Zophoridae		Ironbark beetles	X	
	<i>Dermaptera</i>					
				Earwigs	X	1

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
	<i>Diptera</i>					
		Asilidae		Robber Flies	X	
		Asilidae - Ommatiinae	<i>Ommatius sp.</i>	Feathery Antennae Robber Flies		1
		Athericidae		Water Snipe Flies		1
		Calliphoridae		Blow Flies	X	
		Calliphoridae	<i>Calliphora sp.</i>	Blowflies		1
		Ceratopogonidae		Biting Midges		4
		Chironomidae		Midges	X	
		Culicidae		Mosquitoes	X	
		Empididae		Balloon Flies		2
		Muscidae		Bush Flies	X	3
		Mycetophilidae		Fungus Gnats		4
		Stratiomyidae		Soldier Flies		2
		Syrphidae		Hover Flies	X	
		Tabanidae		March Flies	X	
		Tephritidae		Fruit Flies		1
		Tipulidae		Crane Flies	X	1
				Flies		1
	<i>Diptera - Nematocera</i>					
				Fungus Gnats and Midges		9
	<i>Ephemeroptera</i>					
				Mayflies	X	
	<i>Hemiptera</i>					
		Alydidae	<i>Noliphus sp.</i>	Board-headed Bugs		1
		Aphididae		Aphids	X	
		Cicadellidae		Leaf Hoppers	X	
		Cicadellidae		Leafhoppers and Treehoppers		8
		Cicadellidae		Leafhoppers		1
		Cicadidae - Cicadinae		Cicadas	X	1
		Cixiidae		Cixiid Planthoppers		2
		Coccoidea		Scale Bugs	X	
		Eurybrachyidae	<i>Platybrachys sp.</i>	Bark Hoppers		1
		Flatidae		Plant Hoppers		1
		Fulgoridae	<i>Eurinopsyche sp.</i>	Fulorid Planthopper		1
		Gerridae		Water Striders	X	
		Lygaeidae		Ground Bugs	X	

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
		Margarodidae		Mealy Bugs		1
		Membracidae	<i>Sextius</i> sp.	Horned Treehoppers		1
		Notonectidae		Backswimmers	X	2
		Pentatomidae		Stink Bugs	X	
		Reduviidae		Assassin Bugs	X	
		Reduviidae	<i>Pristhesancus plagipennis</i>	Common Assassin Bug		1
		Ricaniidae		Ricaniid Planthopper		1
	<i>Hymenoptera</i>					
		Apidae		Bees	X	
		Apidae	<i>Xylocopa bombylans</i>	Metallic Carpenter Bee		1
		Apidae	<i>Amegilla</i> sp.	Blue-banded Bees		1
		Braconidae		Braconid Wasps		1
		Formicidae		Ants	X	
		Formicidae - Dolichoderinae	<i>Anonychomyrma</i> sp.	Ants		1
		Formicidae - Dolichoderinae	<i>Iridomyrmex chasei</i>	Tyrant Ants		1
		Formicidae - Ectatominae	<i>Rhytidoponera metallica</i>	Greed-headed Ant		1
		Formicidae - Formicinae	<i>Camponotus consobrinus</i>	Ants		1
		Formicidae - Formicinae	<i>Camponotus nigriceps</i>	Ants		1
		Formicidae - Formicinae	<i>Notoncus capitatus</i>	Ants		1
		Formicidae - Formicinae	<i>Nylanderia</i> sp.	Ants		2
		Formicidae - Formicinae	<i>Paraparatrechina</i> sp.	Ants		1
		Formicidae - Formicinae	<i>Polyrachis</i> sp.	Ants		2
		Formicidae - Myrmicinae	<i>Aphenogaster longiceps</i>	Ants		1
		Formicidae - Myrmicinae	<i>CreMATogaster</i> sp.	Ants		2
		Formicidae - Myrmicinae	<i>Monomorium</i> sp.	Ants		1
		Formicidae - Myrmicinae	<i>Pheidole</i> sp.	Ants		1



Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
		Formicidae - Myrmicinae	<i>Tetramorium</i> sp.	Ants		1
		Formicidae - Myrmicinae	<i>Myrmecia</i> sp.	Bull Ants		1
		Formicidae - Myrmicinae	<i>Myrmecia</i> sp.	Jumping Ants		1
		Ichneumonoidea (superfamily)		Parasitic Wasps	X	
		Ichneumonidae - Banchinae		Mottled Ichneumon Wasps		1
		Ichneumonidae - Ichneumoninae		Orange Ichneumon Wasps		1
		Sphecidae		Sphecid Wasps	X	
		Tiphiidae		Tiphiid Wasps		1
		Vespidae		Vespid Wasps	X	
		Vespidae	<i>Polistes</i> sp.	Paper Nest Wasps		1
	<i>Lepidoptera</i>					
		Arctidae		Tiger Moths	X	
		Danainae (subfamily)		Milkweed Butterflies	X	
		Geometridae		Geometer Moths	X	2
		Hesperiidae		Skippers	X	1
		Lycaenidae		Ant Blue Butterflies	X	
		Notodontidae - Thaumetopoeinae		Processionary Caterpillars		1
		Nymphalidae	<i>Euploea core</i>	Common Crow Butterfly		1
		Nymphalidae		Swallowtail Butterflies	X	
		Sesiidae		Clearwing Moths	X	
		Sphingidae		Hawk Moths	X	
				Moths		31
	<i>Mantodea</i>					
		Mantidae		Praying Mantis	X	2
	<i>Neuroptera</i>					
				Lace Wings	X	
	<i>Odonata</i>					
		Lestidae		Damselflies		3
		Zygoptera		Damselflies	X	
		Anisoptera		Dragonflies	X	4

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	Biosphere 2006 (pres/abs)	ACA 2016 (# species)
	<i>Orthoptera</i>					
		Acrididae		Grasshoppers	X	3
		Gryllacridae		Tree Crickets	X	
		Gryllidae		True Crickets	X	2
		Gryllotalpidae		Mole Crickets	X	
		Tettigoniidae		Katydid	X	
	<i>Plecoptera</i>					
				Stoneflies	X	
	<i>Trichoptera</i>					
				Caddisflies		1
Arthropoda - Malacostraca						
	<i>Amphipoda</i>					
		Taltridae		Amphipods		1
Arthropoda - Malacostraca						
	<i>Isopoda</i>					
		Oniscidea	<i>Armadillidium vulgare</i>	Slaters		1
		Oniscidea	<i>Porcellio scaber</i>	Slaters		1
Molusca - Gastropoda						
	<i>Helicoidea</i>					
		Helicidae		Snails	X	
	<i>Parmacelloidea</i>					
		Milacidae		Slugs	X	
	<i>Pulmonata</i>					
		Planorbidae		Planorbid Snails		1
				Land Snails		1
	<i>Sigmurethra</i>					
		Milacidae		Slugs		1

#### Invertebrates recorded in Bell Park

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	ACA 2016 (# species)
Annelida - Oligochaeta					
		<i>Megadrilacea</i>		Earthworms	1
Arthropoda - Arachnida					
	<i>Araneae</i>				
		Araneidae	<i>Araneus</i> sp.	Orb-weaving Spiders	1
		Araneidae	<i>Dolophones</i> sp.	Orb-weaving Spiders	1
		Araneidae	<i>Eriophora</i> sp.	Orb-weaving Spiders	1
		Araneidae		Orb-weaving Spiders	4
		Miturgidae		Prowling Spiders	1
		Oxyopidae	<i>Oxyopes</i> sp.	Lynx Spiders	1
		Salticidae		Jumping Spiders	1

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	ACA 2016 (# species)
		Sparassidae		Huntsman Spiders	1
		Tetragnathidae	<i>Leucage granulata</i>	Silver Orb Spider	1
		Uloboridae		Uloborid Spiders	1
	<i>Trombidiformes</i>				
		Trombidiidae		Mites	1
Arthropoda - Diplopoda					
	<i>Polydesmida</i>			Polydesmid Millipedes	2
Arthropoda - Entognatha					
	<i>Entomobryomorpha</i>				
		Isotomidae		Springtails	1
	<i>Symphyleona</i>				
		Smithuridae		Springtails	2
Arthropoda - Insecta					
	<i>Blattodea</i>				
		Blaberidae		Bark Cockroaches	1
	<i>Coleoptera</i>				
		Carabidae		Ground Beetles	1
		Chrysomelidae - Chrysomelinae		Leaf Beetles	1
		Staphylinidae		Rove Beetles	1
	<i>Diptera</i>				
		Culicidae	<i>Culex sp.</i>	Mosquitoes	3
		Muscidae		Bush Flies	2
		Mycetophilidae		Fungus Gnats	2
		Sciaridae		Fungus Gnats	2
		Stratiomyidae		Soldier Flies	1
		Syrphidae		Hover Flies	1
		Tipulidae		Crane Flies	1
	<i>Diptera - Nematocera</i>				
				Fungus Gnats and Midges	2
	<i>Hemiptera</i>				
		Aphididae		Aphids	2
		Cicadellidae		Leafhoppers and Treehoppers	2
		Cicadellidae		Leafhoppers	1
		Coccoidea		Scale Insects	3
		Eurybrachyidae	<i>Platybrachys sp.</i>	Bark Hoppers	1
		Miridae	<i>Trilaccus sp.</i>	Mirid Bugs	1
		Miridae		Mirid Bugs	1
		Reduviidae		Assassin Bug	1

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	ACA 2016 (# species)
	<i>Hymenoptera</i>				
		Apidae - Apinae	<i>Apis mellifera</i>	European Honey Bee	1
		Formicidae - Dolichoderinae	<i>Anonychomyrma</i> sp.	Ants	1
		Formicidae - Dolichoderinae	<i>Iridomyrmex chasei</i>	Tyrant Ants	1
		Formicidae - Dolichoderinae	<i>Iridomyrmex</i> sp.	Tyrant Ants	1
		Formicidae - Ectatominae	<i>Rhytidoponera metallica</i>	Greed-headed Ant	1
		Formicidae - Formicinae	<i>Notoncus capitatus</i>	Ants	1
		Formicidae - Myrmicinae	<i>Crematogaster</i> sp.	Ants	2
		Formicidae - Myrmicinae	<i>Pheidole</i> sp.	Ants	1
		Ichneumonidae - Ichneumoninae		Orange Ichneumon Wasps	1
	<i>Lepidoptera</i>				
		Psyshidae		Case Moths	1
				Moths	21
	<i>Neuroptera</i>			Lacewings	1
	<i>Trichoptera</i>			Caddisflies	2
Arthropoda - Malacostraca					
	<i>Amphipoda</i>				
		Taltridae		Amphipods	2
	<i>Isopoda</i>				
		Oniscidea	<i>Armadillidium vulgare</i>	Slaters	1
		Oniscidea	<i>Porcellio scaber</i>	Slaters	1

#### Invertebrates recorded in Outlook Park

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	ACA 2016 (# species)
Arthropoda - Arachnida					
	<i>Araneae</i>				
		Araneidae	<i>Argiope keyserlingi</i>	St Andrew's Cross Spider	1
		Araneidae	<i>Phonognatha</i> sp.	Leaf Curling Spiders	1
		Araneidae		Orb-weaving Spiders	3
		Clubionidae	<i>Clubionia</i> sp.	Sac Spiders	1
		Lycosidae		Wolf Spiders	1
		Oxyopidae	<i>Oxyopes</i> sp.	Lynx Spiders	2
		Salticidae		Jumping Spiders	2

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	ACA 2016 (# species)
		Sparassidae		Huntsman Spiders	1
		Tetragnathidae	<i>Leucage granulata</i>	Silver Orb Spider	1
		Theridiidae		Scaffold-web Weavers	3
		Thomsidae	<i>Diaea sp.</i>	Flower Spiders	2
		Thomsidae		Crab Spiders	1
Arthropoda - Diplopoda					
	<i>Polydesmida</i>				
				Polydesmid Millipedes	2
	<i>Spirostreptida</i>				
				Spirostrepid Millipedes	1
Arthropoda - Entognatha					
	<i>Entomobryomorpha</i>				
		Isotomidae		Springtails	2
Arthropoda - Insecta					
	<i>Blattodea</i>				
		Blaberidae		Bark Cockroaches	1
		Blattidae	<i>Polyzosteria sp.</i>	Coackroaches	1
		Blattidae		Coackroaches	2
	<i>Coleoptera</i>				
		Carabidae		Ground Beetles	1
		Chrysomelidae - Chrysomelinae		Leaf Beetles	1
		Elateridae		Click Beetles	1
		Scarabidae		Cockchafers	2
	<i>Diptera</i>				
		Calliphoridae	<i>Calliphora sp.</i>	Blowflies	1
		Culicidae	<i>Culex sp.</i>	Mosquitoes	5
		Muscidae		Bush Flies	1
		Mycetophilidae		Fungus Gnats	2
		Sciaridae		Fungus Gnats	3
		Stratiomyidae		Soldier Flies	1
		Tipulidae		Crane Flies	2
				Flies	2
	<i>Diptera - Nematocera</i>				
				Fungus Gnats and Midges	6
	<i>Hemiptera</i>				
		Cicadellidae			
				Leafhoppers and Treehoppers	2
		Cicadellidae		Leafhoppers	1
		Cicadidae - Cicadinae		Cicadas	1

Phylum - Class	Order	Family - Subfamily	Genus/species	Common name	ACA 2016 (# species)
		Cixiidae		Cixiid Planthoppers	1
		Eurybrachyidae	<i>Platybrachys</i> sp.	Bark Hoppers	1
		Reduviidae		Assassin Bug	1
	<i>Hymenoptera</i>				
		Formicidae - Dolichoderinae	<i>Iridomyrmex</i> sp.	Tyrant Ants	1
		Formicidae - Ectatominae	<i>Rhytidoponera metallica</i>	Greed-headed Ant	1
		Formicidae - Formicinae	<i>Paraparatrechina</i> sp.	Ants	1
		Formicidae - Myrmicinae	<i>CreMATogaster</i> sp.	Ants	2
		Formicidae - Myrmicinae	<i>Myrmecia</i> sp.	Bull Ants	1
		Ichneumonidae		Ichneumon Wasps	1
	<i>Lepidoptera</i>				
		Nymphalidae	<i>Euploea core</i>	Common Crow Butterfly	1
		Psyshidae		Case Moths	1
				Moths	19
	<i>Psocoptera</i>				
				Booklice	3
	<i>Trichoptera</i>				
				Caddisflies	3
Arthropoda - Malacostraca					
	<i>Amphipoda</i>				
		Taltridae		Amphipods	2
	<i>Isopoda</i>				

## Reference for Section C

- Chen, T. 2013. Fox baiting program in Ku-ring-gai causes rise of bandicoots and ticks. *North Shore Times*. Web: <http://www.dailytelegraph.com.au/newslocal/north-shore/fox-baiting-program-causes-rise-of-bandicoots-and-ticks/story-fngr8h9d-1226684782853>, accessed 19 Oct. 2016.
- Claridge, A.W., Paull, D.J. and Barry, S.C., 2010. Detection of medium-sized ground-dwelling mammals using infrared digital cameras: an alternative way forward?. *Australian Mammalogy*, 32(2), pp.165-171.
- Clarke, M.F. and Oldland, J.M., 2007. Penetration of remnant edges by noisy miners (*Manorina melanocephala*) and implications for habitat restoration. *Wildlife Research*, 34(4), pp.253-261.
- Dexter, N. and Murray, A., 2009. The impact of fox control on the relative abundance of forest mammals in East Gippsland, Victoria. *Wildlife Research*, 36(3), pp.252-261.
- Friedlander, J. 2007. Possum Magic Is Fading Away. *Sydney Morning Herald*. Web: <http://www.smh.com.au/news/environment/possum-magic-is-fading-away/2007/07/16/1184559705364.html>, accessed 19 Oct. 2016.
- Grey, Marilyn J., Michael F. Clarke, and Richard H. Loyn. "Influence of the Noisy Miner *Manorina melanocephala* on avian diversity and abundance in remnant Grey Box woodland." *Pacific Conservation Biology* 4.1 (1998): 55-69.
- Hastings, R.A. and Beattie, A.J., 2006. Stop the bullying in the corridors: can including shrubs make your revegetation more Noisy Miner free?. *Ecological Management & Restoration*, 7(2), pp.105-112.
- Hoh, A., 2016. 2016. Brush turkeys invading suburban Sydney backyards. *702 ABC Sydney*. Web: <http://www.abc.net.au/news/2016-03-31/brush-turkeys-haunting-sydney-backyards/7287518>, accessed 19 Oct. 2016.
- Hoskin, C.J., Grigg, G.C., Stewart, D. A. & Macdonald, S.L., 2015. Frogs of Australia (1.0.2 / 4210) [Mobile application software]. Retrieved from <http://www.ugmedia.com.au>.
- Inions, G.B., Tanton, M.T. and Davey, S.M., 1989. Effect of Fire on the Availability of Hollows in Trees Used by the Common Brushtail Possum, *Trichosurus Vulpecula* Kerr, 1792, and the Ringtail Possum, *Pseudocheirus Peregrinus* Boddaerts, 1785. *Wildlife Research*, 16(4), pp.449-458.
- Kennett, R., Roe, J., Hodges, K. and Georges, A., 2009. *Chelodina longicollis* (Shaw 1784)– eastern long-necked turtle, common long-necked turtle, common snake-necked turtle. *Chelonian Research Monographs*, 5, pp.31-1.
- Koenig, J., Shine, R. and Shea, G., 2001. The ecology of an Australian reptile icon: how do blue-tongued lizards (*Tiliqua scincoides*) survive in suburbia?. *Wildlife Research*, 28(3), pp.214-227.
- Lunney, D., Cullis, B. and Eby, P., 1987. Effects of logging and fire on small mammals in Mumbulla State Forest, near Bega, New South Wales. *Wildlife Research*, 14(2), pp.163-181.

Matthews, A., Lunney, D., Waples, K., and Hardy, J., 2003. Brushtail Possums: “champions of the suburbs” or “our tormentors”? In *Urban Wildlife: More than meets the eye*. Edited by D. Lunney and S. Burgin. Proceedings of the Royal Zoological Society of NSW, Sydney.

Meacham, S., 2015. The Turkey army that’s invading suburban Sydney. *The New Daily*. Web: <http://thenewdaily.com.au/news/state/nsw/2015/10/06/brush-turkeys-invading-sydney/>, accessed 19 Oct. 2016.

Mo, M., 2014. Habitat selection of the Broad-tailed Gecko *Phyllurus platurus* in an urban Sydney bushland remnant. *Australian Zoologist*, 37(1), pp.95-101.

Paull, D.J., Claridge, A.W. and Barry, S.C., 2011. There's no accounting for taste: bait attractants and infrared digital cameras for detecting small to medium ground-dwelling mammals. *Wildlife Research*, 38(3), pp.188-195.

Scott, L.K., Hume, I.D. and Dickman, C.R., 1999. Ecology and population biology of long-nosed bandicoots (*Perameles nasuta*) at North Head, Sydney Harbour National Park. *Wildlife Research*, 26(6), pp.805-821.

Shine, R. and Koenig, J., 2001. Snakes in the garden: an analysis of reptiles “rescued” by community-based wildlife carers. *Biological Conservation*, 102(3), pp.271-283.

Smith, J. and Lill, A., 2008. Importance of eucalypts in exploitation of urban parks by Rainbow and Musk Lorikeets. *Emu*, 108(2), pp.187-195.

Stokeld, D., Hamer, A.J., van der Ree, R., Pettigrove, V. and Gillespie, G., 2014. Factors influencing occurrence of a freshwater turtle in an urban landscape: a resilient species?. *Wildlife Research*, 41(2), pp.163-171.

Veerman, P.A., 2016. The changing status of the Rainbow Lorikeet *Trichoglossus haematodus* in south-east Australia: the role of wild and escaped birds. *Australian Field Ornithology*, 14(1).

White, A., 2006. Ryde flora and fauna study 2006. *Biosphere Environmental Consultants Pty Ltd*. Ryde City Council. Available at: [fieldofmarseec.nsw.edu.au/wp-content/uploads/2013/02/FloraFauna\\_Study\\_2006.pdf](http://fieldofmarseec.nsw.edu.au/wp-content/uploads/2013/02/FloraFauna_Study_2006.pdf).

White, J.G., Antos, M.J., Fitzsimons, J.A. and Palmer, G.C., 2005. Non-uniform bird assemblages in urban environments: the influence of streetscape vegetation. *Landscape and urban planning*, 71(2), pp.123-135.



## Appendix C1. Method details

### Spotlighting

The head torches used for nocturnal surveys were Ledlenser H14.

### Reptile identification

Reptiles were identified visually using:

Wilson, S. and Swan, G., 2013. A complete guide to Reptiles of Australia. 4<sup>th</sup> edition.

### Frog identification

Frogs were identified visually and aurally using the mobile application:

Hoskin, C.J., Grigg, G.C., Stewart, D. A. & Macdonald, S.L., 2015. Frogs of Australia (1.0.2 / 4210) [Mobile application software]. Retrieved from <http://www.ugmedia.com.au>.

### Mammals

Mammals were identified in the field visually using:

Menkhorst, P. and Knight, F., 2010. A field guide to the mammals of Australia. 3<sup>rd</sup> edition.

Tracks, diggings and scats were visually identified using:

Triggs, B., 2004. Tracks, scats and other traces: a field guide to Australian mammals. Revised edition.

Hair tubes were used for small mammals in preference to cage traps as they are less stressful on fauna, require less handling time, cause less concern with the public and usually aren't interfered with. This technique proved very successful for surveys conducted in other council areas by Biosphere (Kogarah Bushland Reserves: 1997; Rockdale LGA: 1999, Ryde City Council: 2006). This method was used as a repeat from 2006, as per council requirements. The hair tubes used were standard single entrance baited PVC tubes hired from Bernview Environmental Consulting (<http://bernviewenviro.com.au/>) the same as 2006.

Although hair tubing was once a successful non-invasive sampling technique for small mammals, it has its limitations (Claridge et al. 2010; Paull 2011). Tube placement can influence success rates, and in open public parks there are restrictions to where tubes can be installed. Regardless of how well hidden the tubes were we still found that 12 tubes were moved from their original placement and a further nine were completely missing, presumably stolen by the public or pet dogs. We also found that they required a large amount of handling time, and their detections weren't very substantial. Only black rat (*Rattus rattus*) and brush-tailed possum (*Trichosurus vulpecula*) hair was 'trapped' and both these species were easily detected during active searches. Our recommendations for future small mammal surveys would be a number of infrared digital camera traps with baits. If well hidden and chained to trees/stakes baited cameras require a small amount of handling time, are less installation-site specific, and images can be easily identified in the office.

### Birds

Birds were surveyed for using a standard point-count method with flora quadrats as the survey points, details of the method can be found:

<http://www.environment.nsw.gov.au/resources/howyoucanhelp/09BirdSurveysBaseline.pdf>

Birds were identified visually and aurally using the mobile application:

Morcombe, M. and Stewart, D., 2016. eGuide to the birds of Australia (v1.4.2) [mobile application software].

Owl calls were amplified using a Braven BRV-X Bluetooth speaker, on maximum volume.

### **Fish**

Fish were identified in the hand using:

Schmida, G., 2015. A wild Australia guide to freshwater fishes. 2<sup>nd</sup> Edition.

### **Invertebrates**

Invertebrates were identified in the hand or under a microscope using:


Nocturnal invertebrates were collected using UV light traps of the following design:


<http://www.theskepticalmoth.com/techniques/light-traps/>

## **Acknowledgements**

We wish to thank the City of Ryde Environment Unit, Bushcare Volunteers and bush regeneration contractors for their assistance, photographs and anecdotal sightings.

## Figures



 Site boundary (surveyed in 2006 and 2016)



 Site boundary (surveyed in 2016 only)


Figure 1a.  
Boundaries of the six parks / reserve overlaid  
on the SIX Maps 1:25 000 topographic map



Site boundary (surveyed in 2006 and 2016)

Site boundary (surveyed in 2016 only)


Figure 1a-1.  
Boundaries of Bell Park, Brush Farm Park, and Lambert Park,  
overlaid on the SIX Maps 1:25,000 topographic map



Site boundary (surveyed in 2006 and 2016)

Site boundary (surveyed in 2016 only)


Figure 1a-2.  
Boundaries of Darvall Park, and Outlook Park,  
overlaid on the SIX Maps 1:25,000 topographic map



Site boundary (surveyed in 2006 and 2016)

Figure 1a-3.  
Boundary of the Field of Mars Reserve overlaid on the SIX Maps 1:25,000 topographic map






Site boundary (surveyed in 2006 and 2016)

Site boundary (surveyed in 2016 only)



Figure 1b.  
 Boundaries of the six parks / reserve overlaid on the  
 NearMap aerial photograph (dated 19 January 2016)



**Brush Farm Park**

**Lambert Park**

**Bell Park**

-  Site boundary (surveyed in 2006 and 2016)
-  Site boundary (surveyed in 2016 only)







Figure 1b-1.  
Boundaries of Brush Farm, Bell Park and Lambert Park overlaid  
on the NearMap aerial photograph (dated 19 January 2016)



-  Site boundary (surveyed in 2006 and 2016)
-  Site boundary (surveyed in 2016 only)




Figure 1b-2.  
Boundaries of Darvall and Outlook Park overlaid  
on the NearMap aerial photograph (dated 19 January 2016)



# Field of Mars Reserve



 Site boundary (surveyed in 2006 and 2016)



Figure 1b-3.  
Boundary of the Field of Mars Reserve overlaid on the  
NearMap aerial photograph (dated 19 January 2016)



- Site boundary (surveyed 2006 and 2016)
- Site boundary (surveyed 2016 only)
- R2 Low Density Residential
- E2 Environmental Conservation
- RE1 Public Recreation
- SP2 Infrastructure
- RE2 Private Recreation





Figure 2a.  
Boundaries of Bell Park, Brush Farm Park and Lambert Park overlaid on the Land Zoning map sheet LZN\_002 (Ryde LEP 2014)



- Site boundary (surveyed 2006 and 2016)
- Site boundary (surveyed 2016 only)
- R2 Low Density Residential
- R4 High Density Residential
- E2 Environmental Conservation
- RE1 Public Recreation
- RE2 Private Recreation
- SP2 Infrastructure
- B4 Mixed Use





Figure 2b.  
Boundaries of Darvall Park and Outlook Park  
overlaid on the Land Zoning map sheet LZN\_002 (Ryde LEP 2014)



- Site boundary (surveyed 2006 and 2016)
- R2 Low Density Residential
- E2 Environmental Conservation
- RE1 Public Recreation
- IN2 Light Industrial
- SP2 Infrastructure
- SP1 Special Activities
- B1 Neighbourhood Centre





Figure 2c.  
Boundary of the Field of Mars Reserve, overlaid on the  
Land Zoning map sheets LZN\_009 and LZN\_010 (Ryde LEP 2014)








Figure 3a-2.  
 Boundary for the Field of Mars Reserve overlaid on the  
 Sydney 1:100,000 Geological Map (Herbert and West 1983)





Figure 3a-3.  
 Boundary for the Field of Mars Reserve overlaid on the  
 Sydney 1:100,000 Geological Map (Herbert and West 1983) with contour lines



- Site boundary (surveyed in 2006 and 2016)
- Rwa = Ashfield Shale
- Rh = Hawkesbury Sandstone
- Qha = Quaternary deposits
- Transect
- Soil sample location




Figure 3a-4.  
Soil transects overlaid on the Sydney 1:100,000  
Geological Map (Herbert and West 1983)




Figure 3b-1.  
 Boundaries for Bell Park, Brush Farm Park, Darvall Park, Lambert Park, and Outlook Park overlaid on the Sydney 1:100,000 Soil Landscape Map (Chapman et al. 1989)






Figure 3b-2.  
 Boundary of the Field of Mars Reserve overlaid on the  
 Sydney 1:100,000 Soil Landscape Map (Chapman et al. 1989)











Figure 4b.  
 Boundary of the Field of Mars Reserve, overlaid on the  
 Heritage Map sheets HER\_006, HER\_010 (Ryde LEP 2014)





-  Site boundary (surveyed in 2006 and 2016)
-  Site boundary (surveyed in 2016 only)







Figure 5a.  
Boundaries of Bell Park, Brush Farm Park, and Lambert Park overlaid on SIX Maps historical aerial photograph (dated 1943)



-  Site boundary (surveyed in 2006 and 2016)
-  Site boundary (surveyed in 2006 only)





Figure 5b.  
Boundaries of Darvall Park and Outlook Park  
overlaid on SIX Maps historical aerial photograph (dated 1943)



Site boundary (surveyed in 2006 and 2016)

Figure 5c.  
Boundary of the Field of Mars Reserve overlaid on the  
SIX Maps historical aerial photograph (dated 1943)




Figure 6a-1.  
Boundaries of Bell Park, Brush Farm Park and Lambert Park, overlaid on the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)




Figure 6a-2.  
 Boundaries of Darvall Park and Outlook Park, overlaid on the  
 Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)




Figure 6a-3.  
 Boundary of the Field of Mars Reserve overlaid on the  
 Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)

APPENDIX 5:

VEGETATION MAPS

Map 5A: Brush Farm Park and Lambert Park




Figure 6b-1. Native vegetation mapping of Brush Farm Park and Lambert Park (Map 5A, Appendix 5, Biosphere 2006)

### MAP 5B: Darvall Park Vegetation Community




Figure 6b-2. Native vegetation mapping of Darvall Park (Map 5B, Appendix 5, Biosphere 2006)



MAP 5C: Field of Mars Vegetation Communities





Figure 6b-3. Native vegetation of the Field of Mars Reserve (Map 5C, Appendix 5, Biosphere 2006)



- Site boundary (surveyed in 2006 and 2016)
- Site boundary (surveyed in 2016)
- Site boundary (unspecified park/recreation area and road reserve)
- Quadrats from previous study, 2006
- Quadrats from current study, 2016





Figure 7a.  
 Quadrats surveyed in 2006 and 2016 in Bell Park, Brush Farm Park, and Lambert Park  
 overlaid on the NearMap aerial photograph (dated 19 January 2016)



- Site boundary (previously studied)
- Site boundary (unstudied)
- Quadrats from previous study, 2006
- Quadrats from current study, 2016




Figure 7b.  
 Quadrats surveyed in 2006 and 2016 in Darvall Park and Outlook Park overlaid on the NearMap aerial photograph (dated 19 January 2016)



Figure 7c.  
 Quadrats surveyed in 2006 and 2016 in the Field of Mars Reserve overlaid on the NearMap aerial photograph (dated 19 January 2016)






Figure 8a. Ground truthing locations surveyed in 2016 in Bell Park, Brush Farm Park, and Lambert Park overlaid on the NearMap aerial photograph (dated 19 January 2016) and the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)



- Site boundary (surveyed in 2006 and 2016)
- Site boundary (surveyed in 2016)
- Quadrats from previous study, 2006
- Quadrats from current study, 2016
- Ground truthing from 2016 study




Figure 8b-1. Ground truthing locations surveyed in 2016 in Darvall Park overlaid on the NearMap aerial photograph (dated 19 January 2016) and the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)



Sydney Metropolitan Mapping

- S\_WSF01: Blue Gum High Forest
- Urban\_E/N: Urban Exotic/Native

- Site boundary (surveyed in 2006 and 2016)
- Site boundary (surveyed in 2016)
- Quadrats from previous study, 2006
- Quadrats from current study, 2016
- Ground truthing from 2016 study




Figure 8b-2. Ground truthing locations surveyed in 2016 in Outlook Park overlaid on the NearMap aerial photograph (dated 19 January 2016) and the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)





Figure 8c.  
 Ground truthing locations surveyed in 2016 in the Field of Mars Reserve overlaid on the NearMap aerial photograph (dated 19 January 2016) and the Native Vegetation of the Sydney Metropolitan Area mapping (OEH 2013, VIS\_ID 3817)





 Site boundary (surveyed in 2006 and 2016)  
 = tracking (GPS) for targeted threatened species search

Figure 9a.  
GPS tracking for the targeted threatened species search in the Field of Mars Reserve  
on the 4 November 2016, overlaid on the NearMap aerial photograph (dated 19 January 2016)




Figure 9b. GPS tracking for the targeted threatened species search in the Field of Mars Reserve on the 4 November 2016, with ground truthing locations surveyed in 2016, overlaid on the NearMap aerial photograph (dated 19 January 2016)

## Tables

**Table 1. Species recorded at the 11 re-surveyed quadrats (Brush Farm Park, Darvall Park, Field of Mars Reserve, Lambert Park) and 2 previously unsurveyed quadrats (Bell Park, Outlook Park)**

Note: 1. Asterisk (\*) before botanical name signifies exotic species. Hash symbol (#) signifies a non-local native, planted or naturalised.  
 2. Families are grouped under headings 1. Pteridophytes, 2. Gymnosperms, 3. Dicotyledons, 4. Monocotyledons. One or more of these plant groups may be absent from these sites.

3. The numbers in the columns denote average percent (%) projected foliage cover for each quadrat, from the percent (%) projected foliage cover recorded in the four sub-quadrats.

Botanical name	Common name	Quadrat												
		BF#1	BF#1	BF#2	BF#3	BF#4	DP#1	FoM#1	FoM#2	FoM#3	FoM#4	FoM#5	LP#1	OP#1
<b>1. Pteridophytes</b>														
<b>Adiantaceae</b>														
<i>Adiantum hispidulum</i>	Rough Maidenhair Fern			22.5									0.025	
<i>Pellaea nana</i>	Small Sickle Fern								0.025					
<b>Aspleniaceae</b>														
<i>Asplenium australasicum</i>	Birds-nest Fern								0.05					
<i>Asplenium flabellifolium</i>	Necklace Spleenwort, Necklace Fern								0.025					
<b>Blechnaceae</b>														
<i>Doodia aspera</i>	Prickly Rasp Fern									0.025				
<i>Doodia caudata</i>	Small Rasp Fern			0.025					0.025					
<b>Cyatheaceae</b>														
# <i>Cyathea cooperi</i>	Straw Tree-fern, Scaly Tree-Fern												0.25	
<b>Dennstaedtiaceae</b>														
<i>Pteridium esculentum</i>	Bracken								0.025	0.75	1.75			
<b>Dicksoniaceae</b>														
<i>Calochlaena dubia</i>	Rainbow Fern, False Bracken								0.525					
<b>Lindsaeaceae</b>														
<i>Lindsaea linearis</i>	Screw Fern											0.025		
<b>Pteridaceae</b>														
<i>Pteris tremula</i>	Tender Brake							0.075						

Botanical name	Common name	Quadrat											
		BF#1	BF#1	BF#2	BF#3	BF#4	DF#1	FoM#1	FoM#2	FoM#3	FoM#4	FoM#5	LP#1
<b>Thelypteridaceae</b>													
<i>Cyclosorus dentatus</i>			0.25									0.05	
<b>2. Gymnosperms</b>													
<b>Araucariaceae</b>													
<i>Araucaria heterophylla</i>	Norfolk Island Pine												0.025
<b>3. Dicotyledons</b>													
<b>Acanthaceae</b>													
<i>Avicennia marina</i> subsp. <i>australasica</i>	Grey Mangrove						31.25						
* <i>Justicia</i> sp.													0.025
<i>Pseuderanthemum variable</i>	Pastel Flower	0.05	0.025	0.1	1.75	0.025	0.075						0.55
* <i>Thunbergia alata</i>	Black-eyed Susan					0.05							
<b>Aceraceae</b>													
* <i>Acer negundo</i>	Box-elder Maple, Box Elder											0.025	
<b>Aizoaceae</b>													
<i>Tetragonia tetragonioides</i>	New Zealand Spinach, Native Spinach, Warrigal Cabbage						6.275						
<b>Amaranthaceae</b>													
<i>Alternanthera denticulata</i>	Common Joyweed						0.025						
<b>Anacardiaceae</b>													
* <i>Toxicodendron succedaneum</i>	Rhus Tree, Wax Tree					0.5							
<b>Aphanopetalaceae</b>													
<i>Aphanopetalum resinosum</i>	Gum Vine		4.5	0.025								0.75	
<b>Apiaceae</b>													
<i>Apium prostratum</i>	Sea Celery						1.55						
<i>Centella asiatica</i>	Indian Pennywort					0.025							
* <i>Cyclospermum leptophyllum</i>	Slender Celery					0.025							
<i>Platysace lanceolata</i>	Lance-leaf Platysace, Shrubby Platysace											0.025	
<i>Xanthosia pilosa</i>									0.025				

Botanical name	Common name	Quadrat											
		BF#1	BF#1	BF#2	BF#3	BF#4	DF#1	FoM#1	FoM#2	FoM#3	FoM#4	LP#1	OP#1
Xanthosia tridentata	Rock Xanthosia											0.025	
<b>Apocynaceae</b>													
* Araujia sericifera	Moth Vine, Cruel Plant		0.05			0.025			0.025				
Parsonia straminea	Common Silkpod, Monkey Rope									0.05			
<b>Araliaceae</b>													
* Hedera helix	Ivy, English Ivy	0.025		1.25							0.05	0.05	0.025
Polyscias sambucifolia	Elderberry Panax					0.025				2	0.05		
<b>Asteraceae</b>													
* Aster subulatus	Wild Aster						0.05						
* Bidens pilosa	Cobbler's Pegs	0.25			0.075	0.025						0.025	
* Coreopsis lanceolata	Calliopsis, Coreopsis								0.025				
* Delairea odorata	Cape Ivy	1											
* Hypochaeris radicata	Catsear, False Dandelion										0.025		
Ozothamnus diosmifolius	White Dogwood										0.5		
Sigesbeckia orientalis	Indian Weed				0.025	0.3							0.025
* Sonchus oleraceus	Common Sow-thistle, Milk-thistle						0.025						
<b>Basellaceae</b>													
* Anredera cordifolia	Madeira Vine, Lamb's Tail	0.5		0.05			0.025					0.075	
<b>Bignoniaceae</b>													
* Jacaranda mimosifolia	Jacaranda	1.25											0.5
Pandorea pandorana	Wonga Vine				0.55			1.525	0.25		0.075		
<b>Casuarinaceae</b>													
Allocauarina littoralis	Black She-oak							1.25		2	0.5		
Allocauarina torulosa	Forest She-oak												0.5
# Casuarina cunninghamiana	River Oak, River She-oak											2.5	
Casuarina glauca	Swamp Oak, Swamp She-oak						7.5						
<b>Celastraceae</b>													
Celastrus australis									0.025				
Denhamia silvestris	Narrow-leaved Orangebark, Orange Bush, Orange Bark								0.25				
Elaeodendron australe	Red Olive-plum		2.75		7.5	0.5							











Botanical name	Common name	Quadrat												
		BF#1	BF#1	BF#2	BF#3	BF#4	DF#1	FoM#1	FoM#2	FoM#3	FoM#4	FoM#5	LP#1	OP#1
<b>Passifloraceae</b>														
<i>Passiflora herbertiana</i>	Native Passionfruit			0.025		4							2.5	
* <i>Passiflora suberosa</i>	Corky Passionflower				1.775	0.75								0.025
* <i>Passiflora subpeltata</i>	White Passionfruit	0.25	0.025	0.025	0.275								0.5	
<b>Pittosporaceae</b>														
<i>Billardiera scandens</i>	Hairy Apple Berry, Dumplings										0.075	0.025		
<i>Bursaria spinosa</i>	Australian Boxthorn		0.25											
<i>Pittosporum multiflorum</i>	Orange Thorn		0.525		1.525									
<i>Pittosporum revolutum</i>	Yellow Pittosporum					1.25							0.775	0.25
<i>Pittosporum undulatum</i>	Pittosporum	1.25	14.5	5.75	12	5.75		10	1.275	1.775	1	7.75	5.5	
<b>Plantaginaceae</b>														
* <i>Plantago major</i>	Large Plantain									0.025				
<b>Polygonaceae</b>														
* <i>Acetosa sagittata</i>	Rambling Dock, Turkey Rhubarb									0.275				
<i>Persicaria</i> sp.	Knotweed									0.025				
<i>Rumex brownii</i>	Slender Dock									0.3				
<b>Primulaceae</b>														
<i>Samolus repens</i>	Creeping Brookweed									21.75				
<b>Proteaceae</b>														
<i>Banksia spinulosa</i>	Hairpin Banksia										1.025			
# <i>Grevillea robusta</i>	Silky Oak	1.25			0.525							0.525		2.5
<i>Grevillea sericea</i>														
<i>Hakea laevipes</i>											0.25			
<i>Hakea salicifolia</i>	Willow Hakea											1		
<i>Lambertia formosa</i>	Mountain Devil								0.25					
<i>Lomatia silaifolia</i>	Crinkle Bush							0.025	0.025	0.325	0.025			
<i>Persoonia levis</i>	Broad-leaved Geebung											0.25		
# <i>Stenocarpus sinuatus</i>	Firewheel Tree			0.025	0.05									0.025
<b>Ranunculaceae</b>														
<i>Clematis glycinoides</i>	Headache Vine, Traveller's Joy, Old Man's Beard		2			0.325	0.75		2					0.25

Botanical name	Common name	Quadrat												
		BF#1	BF#1	BF#2	BF#3	BF#4	DF#1	FoM#1	FoM#2	FoM#3	FoM#4	FoM#5	LP#1	OP#1
<b>Rosaceae</b>														
* Eriobotrya japonica	Loquat												0.05	
* Rubus laudatus	Plains Blackberry					1.025								
Rubus rosifolius	Native Raspberry, Rose-leaf Bramble												0.5	
<b>Rubiaceae</b>														
* Galium aparine	Cleavers, Goose-grass, Bedstraw						0.025							
Morinda jasminoides	Morinda	0.25	1.75										0.55	
<b>Rutaceae</b>														
Correa reflexa	Native Fuchsia										0.025			
Melicope micrococca	Hairy-leaved Doughwood	0.25	6.5									0.775	0.025	
# Murraya paniculata	Jasmine-orange, Cosmetic Bark												0.025	
Zieria smithii	Sandfly Zieria, Stinkwood							0.775						
<b>Sapindaceae</b>														
Alectryon subcinerus	Native Quince	10	1.25											
* Cardiospermum grandiflorum	Balloon Vine					1.25	0.05							
Dodonaea triquetra	Hopbush							0.25		0.05	3.275		0.025	
Guioa semiglauca	Guioa	0.5	0.25											
<b>Solanaceae</b>														
* Cestrum nocturnum	Night-scented Jessamine, Lady of the Night					0.25							0.775	
* Physalis peruviana	Cape Gooseberry												0.025	
Solanum aviculare	Kangaroo Apple					0.025								
* Solanum chenopodioides	White-tip Nightshade							0.025						
* Solanum nigrum	Blackberry Nightshade								0.025					
Solanum prinophyllum	Forest Nightshade					0.025								
* Solanum pseudocapsicum	Jerusalem Cherry	0.025	0.25			0.075							0.05	
* Solanum seaforthianum	Brazilian Nightshade							0.775						
* Solanum sp.								0.25						
<b>Sterculiaceae</b>														
# Brachychiton acerifolius	Illawarra Flame-tree, Flame Kurrajong	0.5	2	2.25	1.275								1.5	
<b>Stylidiaceae</b>														
Stylidium graminifolium	Grass-leaf Triggerplant												0.025	



Botanical name	Common name	Quadrat												
		BF#1	BF#1	BF#2	BF#3	BF#4	DF#1	FoM#1	FoM#2	FoM#3	FoM#4	FoM#5	LP#1	OP#1
<b>Asphodelaceae</b>														
* Aloe arborescens			0.025											
<b>Commelinaceae</b>														
Commelina cyanea	Blue Spiderwort		3	0.025			2.525	0.05						1
* Tradescantia fluminensis	Trad, Wandering Jew		45	7.525		0.5	42.5	0.3		0.025				8
<b>Cyperaceae</b>														
* Cyperus eragrostis	Drain Flat-sedge, Umbrella Sedge							0.025						
Cyperus imbecillis														0.025
Cyperus tetraphyllus				0.25									0.3	0.275
Gahnia clarkei	Tall Saw-sedge								0.025					
Lepidosperma elatius	Tall Sword-sedge								0.025	0.8				
Lepidosperma laterale	Variable Sword-sedge							0.05			0.05			
Schoenus melanostachys	Black Bog-rush											1		
<b>Iridaceae</b>														
* Dietes grandiflora	Butterfly Iris													0.25
* Dietes iridioides	Butterfly Iris									0.025				
<b>Juncaceae</b>														
Juncus continuus	Rush											0.025		
Juncus kraussii	Sea Rush						8.25							
<b>Lomandraceae</b>														
Lomandra cylindrica	Needle Mat-rush											1		
Lomandra gracilis												0.25		
Lomandra longifolia	Honey Reed, Spike Mat-rush		1			0.05	2.525		1.75	1.25	9.75	10	0.25	0.75
Lomandra multiflora	Many-flowered Mat-rush								0.075		0.25			
Lomandra obliqua											0.025	0.025		
<b>Orchidaceae</b>														
Cryptostylis erecta	Tartan Tongue Orchid											0.025		
Cryptostylis subulata	Large Tongue-orchid, Cow Orchid									0.025				
<b>Philesiaceae</b>														
Eustrephus latifolius	Wombat Berry		0.275	0.55	0.275	0.5			0.5					

Botanical name	Common name	Quadrat												
		BF#1	BF#1	BF#2	BF#3	BF#4	DF#1	FoM#1	FoM#2	FoM#3	FoM#4	FoM#5	LP#1	OP#1
<i>Geitonoplesium cymosum</i>	Scrambling Lily												0.025	
<b>Phormiaceae</b>														
<i>Dianella caerulea</i>	Blue Flax-lily					0.5	0.025		0.3	0.025	0.025	1.25	0.525	0.025
<i>Dianella revoluta</i>	Blue Flax-lily, Spreading Flax-lily									0.075				
<b>Poaceae</b>														
* <i>Andropogon virginicus</i>	Whisky Grass											0.025		
<i>Anisopogon avenaceus</i>	Oat Spear Grass									0.025				
<i>Aristida vagans</i>	Threeawn Speargrass									0.025				
<i>Austrostipa pubescens</i>	Speargrass										0.05	0.025		
<i>Cymbopogon refractus</i>	Barbed-wire Grass							0.025						
<i>Digitaria parviflora</i>	Smallflower Finger Grass											0.05		
* <i>Echinochloa crus-galli</i>	Barnyard Grass						0.25							
<i>Echinopogon caespitosus</i>	Tufted Hedgehog Grass											0.075		
<i>Echinopogon ovatus</i>	Forest Hedgehog Grass							0.05		0.025				
* <i>Ehrharta erecta</i>	Panic Veld-grass	15.025	1.275	2	47.5	9.25		0.5				2.025	0.075	
<i>Entolasia marginata</i>	Bordered Panic			0.025	0.05				0.025					0.05
<i>Entolasia stricta</i>	Wiry Panic				0.025				1	0.025	27.5	6.5		
<i>Eragrostis brownii</i>	Brown's Lovegrass											0.025		
<i>Hemarthra uncinata</i>	Mat Grass						16.775							
<i>Imperata cylindrica</i>	Blady Grass							3.5		6.25	10.75			
* <i>Lolium perenne</i>	Perennial Ryegrass							0.275						
* <i>Megathyrsus maximus</i>	Guinea Grass						1.25							
<i>Microlaena stipoides</i>	Weeping Grass, Meadow Rice-grass		0.025		2.525	4		3.25		1.75	2	0.25		
<i>Oplismenus aemulus</i>	Australian Basket Grass, Wavy Beard Grass	0.05		0.05		10	0.025	3.25			0.025	0.05	5.75	
<i>Oplismenus imbecillis</i>	Narrow-leaved Basket Grass		0.275		0.3	0.5			0.025			0.05		
* <i>Paspalum dilatatum</i>	Paspalum					0.025				0.025				
<i>Poa affinis</i>										0.5		0.25		
* <i>Setaria parviflora</i>								0.275						
<i>Themeda triandra</i>	Kangaroo Grass							8.25		4.75				
<b>Smilacaceae</b>														
<i>Smilax australis</i>	Lawyer Vine, Wait-a-while, Barbwire Vine			0.275	0.775									
<i>Smilax glyciphylla</i>	Sweet Sarsaparilla							0.25	0.25		0.25	1		





Table 2: Species recorded greater than 2 m in height in the 11 re-surveyed quadrats (Brush Farm Park, Darvall Park, Field of Mars Reserve, Lambert Park) and 2 previously unsurveyed quadrats (Bell Park, Outlook Park)

**Bell Park**

Species	1-1		1-2		1-3		1-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
* <i>Celtis sinensis</i>					1	3		
# <i>Corymbia citriodora</i>			1	27				
# <i>Eucalyptus botryoides</i>							1	22
# <i>Eucalyptus grandis</i>	2	30			2	20	1	18
# <i>Eucalyptus scoparia</i>			1	20				
# <i>Grevillea robusta</i>					1	14		
* <i>Ligustrum lucidum</i>			3	3	1	3	5	4
<i>Melaleuca armillaris</i>	2	12						
<i>Pittosporum undulatum</i>							2	5
<i>Syzygium australe</i>			1	8				

**Brush Farm Park**

Species	1-1		1-2		1-3		1-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Acacia parramattensis</i>			2	12	1	2		
<i>Acmena smithii</i>					1	4		
<i>Alectryon subcinereus</i>					1	5	2	8
# <i>Brachychiton acerifolius</i>							1	6
<i>Breynia oblongifolia</i>			3	2			1	2
<i>Bursaria spinosa</i>			1	2			1	2
<i>Callistemon salignus</i>					1	6		
* <i>Celtis sinensis</i>			1	4	1	2		
* <i>Cinnamomum camphora</i>							1	2
<i>Clerodendrum tomentosum</i>	1	2						
# <i>Eucalyptus grandis</i>	2	15	1	12	1	12		
# <i>Eucalyptus microcorys</i>	3	10						
<i>Eucalyptus saligna</i>					1	28		
<i>Ficus coronata</i>							1	4
<i>Ficus rubiginosa</i>			1	10				
* <i>Jacaranda mimosifolia</i>							1	18
# <i>Melia azedarach</i>			1	3				
<i>Notelaea longifolia</i>			1	2				
<i>Pittosporum undulatum</i>	2	2	6	6	4	6	4	3
<i>Senna pendula</i> var. <i>glabrata</i>					1	3		
<i>Syncarpia glomulifera</i>	1	3	1	3	1	12	1	20

**Brush Farm Park**

Species	2-1		2-2		2-3		2-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Acmena smithii</i>			4	10	2	12		
<i>Alectryon subcinereus</i>	3	4					1	5
# <i>Brachychiton acerifolius</i>							3	6
<i>Breynia oblongifolia</i>					2	3		
<i>Elaeodendron australe</i>	5	8	1	2				
<i>Eupomatia laurina</i>	1	7	5	6				
<i>Ficus coronata</i>	6	8	5	8	2	4	7	7
<i>Melicope micrococca</i>	1	3	4	7	1	10		
<i>Myrsine variabilis</i>							5	4
<i>Pittosporum undulatum</i>					3	10		
<i>Schizomeria ovata</i>			1	13				
<i>Syncarpia glomulifera</i>					2	6	1	6

Species	3-1		3-2		3-3		3-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
# <i>Brachychiton acerifolius</i>	1	6			1	7		
* <i>Celtis occidentalis</i>	1	3						
* <i>Cinnamomum camphora</i>	2	5						
<i>Eucalyptus acmenoides</i>	1	22						
<i>Eucalyptus paniculata</i>	1	23						
<i>Eucalyptus saligna</i>							2	14
<i>Ficus coronata</i>			1	8	4	8	1	6
<i>Ficus rubiginosa</i>			1	6				
<i>Grevillea robusta</i>			1	10			1	12
<i>Pittosporum undulatum</i>	8	6	7	6	1	3	4	4
<i>Syncarpia glomulifera</i>					1	16		

**Brush Farm Park**

	4-1		4-2		4-3		4-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Acacia parramattensis</i>	7	10	4	10			1	5
# <i>Brachychiton acerifolius</i>	1	5						
<i>Breynia oblongifolia</i>	1	2						
<i>Elaeodendron australe</i>			3	10			10	6
<i>Eucalyptus</i> sp.			2	6				
* <i>Ligustrum lucidum</i>					2	5	1	5
# <i>Melaleuca styphelioides</i>			1	2				
<i>Notelaea longifolia</i>	1	2	3	3			2	3
<i>Pittosporum undulatum</i>			1	2	2	3	2	4
<i>Syncarpia glomulifera</i>	1	17	2	18	4	19		
* <i>Tecomaria capensis</i>					1	7		

**Darvall Park**

Species	1-1		1-2		1-3		1-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Acmena smithii</i>							1	6
<i>Callistemon salignus</i>					1	4	1	?
<i>Elaeodendron australe</i>							1	5
# <i>Eucalyptus grandis</i>	1	20						
# <i>Eucalyptus microcorys</i>					2	13		
<i>Eucalyptus paniculata</i>	1	20					1	12
<i>Eucalyptus pilularis</i>	2	20						
<i>Eucalyptus saligna</i>			4	20	2	25	1	?
<i>Homalanthus populifolius</i>	1	2						
# <i>Melaleuca styphelioides</i>			4	6				
<i>Pittosporum revolutum</i>			1	3				
<i>Syncarpia glomulifera</i>			2	6			2	7

**Field of Mars Reserve**

Species	1-1		1-2		1-3		1-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Aegiceras corniculatum</i>			1	3				
<i>Avicennia marina</i> subsp. <i>australasica</i>	1	10			2	10		
<i>Casuarina glauca</i>	11	14	6	15			9	12

Species	2-1		2-2		2-3		2-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Angophora costata</i>	5	13	1	13		1	18	
<i>Allocasuarina littoralis</i>	2	4	3	10			2	6
<i>Breynia oblongifolia</i>			1	2				
<i>Elaeocarpus reticulatus</i>	1	3						
<i>Eucalyptus punctata</i>					1	25		
<i>Eucalyptus resinifera</i> subsp. <i>resinifera</i>	2	16	1	20	1	25		
<i>Pittosporum undulatum</i>	1	3	3	5			5	4
<i>Syncarpia glomulifera</i>	1	15	5	15	4	10		

Species	3-1		3-2		3-3		3-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Ceratopetalum apetalum</i>	5	11	3	12	10	15	5	4
* <i>Ligustrum lucidum</i>					1	6	4	4
* <i>Ligustrum sinense</i>					3	6		
<i>Pittosporum undulatum</i>	1	3	2	10	3	12		
<i>Tristaniopsis laurina</i>	1	14	2	13	2	10	3	8

**Field of Mars Reserve**

Species	4-1		4-2		4-3		4-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Allocasuarina littoralis</i>	1	4	3	8	2	4	4	4
<i>Angophora costata</i>	3	14	1	10	3	18	3	20
<i>Corymbia gummifera</i>			1	9	2	16	2	18
<i>Eucalyptus resinifera</i> subsp. <i>resinifera</i>	3	18	3	18	1	20		
<i>Eucalyptus racemosa</i>	2	18	4	18	3	20	2	20
<i>Glochidion ferlinianidi</i>					1	4		
<i>Kunzea ambigua</i>	3	5	2	5				
<i>Pittosporum undulatum</i>	1	2	1	3			1	2
<i>Polyscias sambucifolia</i>	1	3			1	2		

Species	5-1		5-2		5-3		5-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Acacia linifolia</i>	2	2			1	2		
<i>Acacia longifolia</i>					1	2		
<i>Acacia terminalis</i> subsp. <i>angustifolia</i>	2	2	1	2	3	2	1	2
<i>Allocasuarina littoralis</i>					1	6	1	9
<i>Angophora costata</i>			4	20			1	12
<i>Corymbia gummifera</i>							2	16
<i>Dodonaea triquetra</i>	1	2	1	2	12	3		
<i>Elaeocarpus reticulatus</i>	3	5	8	5	3	4	12	6
<i>Eucalyptus piperita</i>	1	18						
<i>Eucalyptus resinifera</i> subsp. <i>resinifera</i>	1	6	1	16	2	20		
<i>Grevillea sericea</i>					1	2		
<i>Hakea salicifolia</i>			1	8				
<i>Kunzea ambigua</i>	7	5	13	5	4	6	11	5
<i>Melaleuca linariifolia</i>	1	7						
<i>Ozothamnus diosmifolius</i>					1	2		
<i>Persoonia levis</i>					1	2		

Lambert Park

Species	1-1		1-2		1-3		1-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Acmena smithii</i>	1	20					3	12
# <i>Casuarina cunninghamiana</i>			2	25				
<i>Cestrum nocturnum</i>			1	3				
# <i>Eucalyptus grandis</i>							1	25
<i>Eucalyptus saligna</i>			1	20			1	25
<i>Ficus coronata</i>	1	2			1	2	2	8
<i>Ficus rubiginosa</i>			4	20	1	20	1	15
<i>Homalanthus populifolius</i>	5	8						
<i>Melicope micrococca</i>			1	5				
<i>Pittosporum revolutum</i>			20	3				
<i>Pittosporum undulatum</i>	1	4		1	2			



## Outlook Park

Species	1-1		1-2		1-3		1-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Acmena smithii</i>	1	2	1	4				
<i>Allocasuarina torulosa</i>					3	13		
* <i>Araucaria heterophylla</i>			1	5				
# <i>Brachychiton acerifolius</i>	3	6	1	3	1	2	2	2
<i>Ceratopetalum apetalum</i>	1	13						
<i>Cryptocarya glaucescens</i>	1	2			1	2		
<i>Elaeocarpus reticulatus</i>								
<i>Eucalyptus saligna</i>	1	35	1	25			2	8
<i>Ficus coronata</i>	2	6					1	35
<i>Ficus rubiginosa</i>	1	14					1	6
<i>Glochidion ferdinandii</i>	1	3						
# <i>Grevillea robusta</i>					1	18		
<i>Homalanthus populifolius</i>			1	2				
* <i>Jacaranda mimosifolia</i>			1	7				
<i>Melaleuca styphelioides</i>			1	3	2	11		
<i>Pittosporum undulatum</i>			2	9	1	2	2	8
# <i>Stenocarpus sinuatus</i>					1	3		
<i>Syncarpia glomulifera</i>			1	25			1	20
# <i>Syzygium paniculatum</i>			1	5				



Botanical name	Quadrats													
	BP#1	BF#1	BF#2	BF#3	BF#4	DP#1	FoM#1	FoM#2	FoM#3	FoM#4	FoM#5	LP#1	OP#1	
<b>Apiaceae</b>														
Apium prostratum							4							
Centella asiatica						1								
* Cyclospermum leptophyllum							1							
Platysace lanceolata												1		
Xanthosia tridentata												1		
<b>Apocynaceae</b>														
* Araujia sericifera			1			1								
Parsonsia straminea										1				
<b>Araliaceae</b>														
* Hedera helix		1		2									1	1
Polyscias sambucifolia										3	2			
<b>Asteraceae</b>														
* Aster subulatus							2							
* Bidens pilosa		1				1							1	
* Conyza sumatrensis			1											
* Coreopsis lanceolata										1				
* Delairea odorata	3													
* Hypochaeris radicata												1		
Ozothamnus diosmifolius												2		
Sigesbeckia orientalis						2								1
* Sonchus oleraceus							1							
<b>Basellaceae</b>														
* Anredera cordifolia		3		1			1						3	
<b>Bignoniaceae</b>														
* Jacaranda mimosifolia		1												1
Pandorea pandorana											1			
<b>Casuarinaceae</b>														
Allocasuarina littoralis										3	3			
Allocasuarina torulosa														1
# Casuarina cunninghamiana													4	
Casuarina glauca							4							
<b>Celastraceae</b>														
Elaeodendron australe						1								
<b>Chenopodiaceae</b>														
Atriplex australasica							2							
<b>Convolvulaceae</b>														
Calystegia marginata		1											1	
Dichondra sp. A														1
* Ipomoea indica	4	3												
<b>Cunoniaceae</b>														
Ceratopetalum apetalum														2
<b>Elaeocarpaceae</b>														
Elaeocarpus reticulatus												4		1
<b>Ericaceae Styphelioideae</b>														
Epacris pulchella										1	1			
Leucopogon ericoides												1		
Leucopogon juniperinus										1	1			
Leucopogon lanceolatus												1		
<b>Euphorbiaceae</b>														
Breynia oblongifolia		2	3									1		
Glochidion ferdinandi										1				1

Botanical name	Quadrats													
	BP#1	BF#1	BF#2	BF#3	BF#4	DP#1	FoM#1	FoM#2	FoM#3	FoM#4	FoM#5	LP#1	OP#1	
Homalanthus populifolius						1	1			1		4	1	
Micrantheum ericoides										3	3			
Phyllanthus gunnii						3								
Phyllanthus hirtellus										1	2			
<b>Fabaceae Caesalpinioideae</b>														
* Senna pendula var. glabrata	2	1					1							
<b>Fabaceae Faboideae</b>														
* Crotalaria sp.							1							
Glycine clandestina						2					1			
Glycine tabacina						1								
Kennedia rubicunda											1			
Platylobium formosum											2			
Pultenaea tuberculata											1			
<b>Fabaceae Mimosoideae</b>														
Acacia decurrens		1	1											
Acacia fimbriata			1											
Acacia linifolia									1	3				
Acacia longifolia										3				
Acacia parramattensis		2				1								
Acacia suaveolens											1			
Acacia terminalis subsp. angustifolia											3			
<b>Geraniaceae</b>														
Geranium homeanum						4								
<b>Goodeniaceae</b>														
Goodenia hederacea										2	2			
<b>Haloragaceae</b>														
Gonocarpus teucroides										1	1			
<b>Lamiaceae</b>														
Clerodendrum tomentosum		1		1										
Plectranthus parviflorus		2	1			3							1	
<b>Lauraceae</b>														
Cassytha pubescens										2	1			
* Cinnamomum camphora		2		1						1		1	1	
Cryptocarya glaucescens													2	
<b>Lobeliaceae</b>														
Lobelia anceps							3							
Pratia purpurascens						1				2	1			
<b>Malvaceae</b>														
Hibiscus heterophyllus			1											
* Sida rhombifolia						2								
<b>Meliaceae</b>														
# Melia azedarach		1										1	1	
# Toona ciliata													1	
<b>Menispermaceae</b>														
Sarcopetalum harveyanum		1	1	2										
Stephania japonica var. discolor		2	2	3		2						3		
<b>Moraceae</b>														
Ficus coronata		1	4	5								3	4	
Ficus rubiginosa		4										5	2	
* Morus alba		1												
<b>Myrsinaceae</b>														
Aegiceras corniculatum							2							

Botanical name	Quadrats												
	BP#1	BF#1	BF#2	BF#3	BF#4	DP#1	FoM#1	FoM#2	FoM#3	FoM#4	FoM#5	LP#1	OP#1
Myrsine variabilis			2										
<b>Myrtaceae</b>													
Acmena smithii		1				1						6	2
Angophora costata									4	4			
Backhousia myrtifolia													1
Callistemon salignus		1	1			1							
# Corymbia citriodora	4												
Corymbia gummifera									4				
# Eucalyptus botryoides	5												
Eucalyptus globoidea										1			
# Eucalyptus grandis	5	4				5						4	
# Eucalyptus microcorys		3				1							
Eucalyptus paniculata				4		4							
Eucalyptus pilularis						5							
Eucalyptus piperita										5			
Eucalyptus racemosa									4				
Eucalyptus resinifera subsp. resinifera									4	5			
Eucalyptus saligna		4				4						4	4
# Eucalyptus scoparia	4												
Kunzea ambigua									3	4			
Leptospermum trinervium									1				
Melaleuca linariifolia										1			
Melaleuca styphelioides						2							2
Syncarpia glomulifera		5	4	5		4							5
Syzygium australe	4												1
# Syzygium paniculatum													1
<b>Nandinaceae</b>													
* Nandina domestica												1	
<b>Ochnaceae</b>													
* Ochna serrulata			2							1	2	2	
<b>Oleaceae</b>													
* Jasminum polyanthum	4												
* Ligustrum lucidum	4	1										2	
* Ligustrum sinense									3	2	1		
Notelaea longifolia		1				1							
* Olea europaea subsp. cuspidata												1	
<b>Oxalidaceae</b>													
* Oxalis latifolia													2
* Oxalis pes-caprae		1											
<b>Passifloraceae</b>													
Passiflora herbertiana			2									4	
* Passiflora suberosa						1							1
* Passiflora subpeltata	1	1	2									3	
<b>Pittosporaceae</b>													
Billardiera scandens									2	2			
Bursaria spinosa		1											
Pittosporum multiflorum		1											
Pittosporum revolutum						1						3	1
Pittosporum undulatum	1	5	4	5					3	3	4	4	
<b>Plantaginaceae</b>													
* Plantago major							1						
<b>Polygonaceae</b>													
* Acetosa sagittata							2						
Persicaria sp.							1						
Rumex brownii							3						

Botanical name	Quadrats													
	BP#1	BF#1	BF#2	BF#3	BF#4	DP#1	FoM#1	FoM#2	FoM#3	FoM#4	FoM#5	LP#1	OP#1	
<b>Primulaceae</b>														
Samolus repens							5							
<b>Proteaceae</b>														
Banksia spinulosa										3				
# Grevillea robusta	1			2										1
Grevillea sericea											3			
Hakea laevipes										1				
Hakea salicifolia											3			
Lomatia silaifolia										2	3			
Persoonia levis											1			
# Stenocarpus sinuatus				1										1
<b>Ranunculaceae</b>														
Clematis glycinoides		3	1			3							2	
<b>Rosaceae</b>														
* Eriobotrya japonica													1	
* Rubus laudatus						3								
Rubus rosifolius														1
<b>Rubiaceae</b>														
* Galium aparine							1							
Morinda jasminoides		2	2										4	
<b>Rutaceae</b>														
Correa reflexa											1			
Melicope micrococca		1	4										1	1
# Murraya paniculata													1	
<b>Sapindaceae</b>														
Alectryon subcinereus		4	2											
* Cardiospermum grandiflorum						4	2							
Dodonaea triquetra									2	4				1
Guioa semiglauca		1	1											
<b>Solanaceae</b>														
* Cestrum nocturnum						2							3	
* Physalis peruviana													1	
* Solanum chenopodioides						1								
* Solanum nigrum							1							
* Solanum pseudocapsicum		1												1
* Solanum seaforthianum						2								
* Solanum sp.						1								
<b>Sterculiaceae</b>														
# Brachychiton acerifolius		1	3	2										2
<b>Thymelaeaceae</b>														
Pimelea linifolia										2	2			
<b>Ulmaceae</b>														
* Celtis occidentalis				1										1
* Celtis sinensis	1	1												
Trema tomentosa var. aspera		2	1									2		1
<b>Urticaceae</b>														
* Parietaria judaica		3												
<b>Verbenaceae</b>														
* Lantana camara	7									1	1			
<b>Vitaceae</b>														
Cayratia clematidea		3	1	2		4								3
Cissus antarctica		1	4	2		5						1		



Botanical name	Quadrats													
	BP#1	BF#1	BF#2	BF#3	BF#4	DP#1	FoM#1	FoM#2	FoM#3	FoM#4	FoM#5	LP#1	OP#1	
Entolasia marginata														1
Entolasia stricta										5		4		
Eragrostis brownii												2		
Hemarthria uncinata							5							
Imperata cylindrica										4		4		
* Lolium perenne							2							
* Megathyrsus maximus						3								
Microlaena stipoides			1			4				3		3	3	
Oplismenus aemulus		2		2		4	1					1	3	4
Oplismenus imbecillis			1			1							3	
* Paspalum dilatatum						2				1				
Poa affinis												1		
Themeda triandra										4				
<b>Smilacaceae</b>														
Smilax australis				2										
Smilax glyciophylla												3		
<b>Xanthorrhoeaceae</b>														
Xanthorrhoea media										3				





Botanical name	Common name	Sub-quadrats				Spot locs.		GT locs.	
		1-1	1-2	1-3	1-4	A	B	GTa	GTb
<b>Fagaceae</b>									
* <i>Quercus robur</i>	English Oak, Pedunculate Oak								X
<b>Lauraceae</b>									
* <i>Cinnamomum camphora</i>	Camphor Laurel					X	X		
<b>Meliaceae</b>									
# <i>Toona ciliata</i>	Australian Red Cedar							X	
<b>Moraceae</b>									
<i>Ficus rubiginosa</i>	Port Jackson Fig, Rusty Fig					X			
<b>Myrtaceae</b>									
<i>Acmena smithii</i>	Lilly-pilly							X	
<i>Angophora costata</i>	Sydney Red Gum, Smooth-barked App							X	
<i>Angophora floribunda</i>	Rough-barked Apple							X	
# <i>Corymbia citriodora</i>	Lemon-scented Gum		20						
# <i>Eucalyptus botryoides</i>	Bangalay				25				
# <i>Eucalyptus grandis</i>	Flooded Gum	35		25	10				X
# <i>Eucalyptus microcorys</i>	Tallowwood					X	X	X	X
<i>Eucalyptus paniculata</i>	Grey Ironbark								X
<i>Eucalyptus pilularis</i>	Blackbutt						X	X	X
# <i>Eucalyptus robusta</i>	Swamp Mahogany						X		
<i>Eucalyptus saligna</i>	Sydney Blue Gum								X
# <i>Eucalyptus scoparia</i>	Wallangarra White Gum		10					X	
# <i>Melaleuca armillaris</i>	Bracelet Honey-myrtle	5							
<i>Melaleuca styphelioides</i>	Prickly Paperbark						X		
<i>Syzygium australe</i>	Brush Cherry		15						
# <i>Syzygium paniculatum</i>	Brush Cherry							X	
<i>Tristaniopsis laurina</i>	Water Gum, Kanuka Box							X	
<b>Ochnaceae</b>									
* <i>Ochna serrulata</i>	Mickey Mouse Plant					X			
<b>Oleaceae</b>									
* <i>Jasminum polyanthum</i>	Jasmine	2	1	5	20		X		
* <i>Ligustrum lucidum</i>	Broad-leaved Privet, Glossy Privet		2		10		X	X	
* <i>Ligustrum sinense</i>	Small-Leaved Privet, Chinese Privet							X	
* <i>Olea europaea</i> subsp. <i>cuspidata</i>	African Olive					X			
<b>Passifloraceae</b>									
* <i>Passiflora suberosa</i>	Corky Passionflower								X
* <i>Passiflora subpeltata</i>	White Passionfruit				1				
<b>Pittosporaceae</b>									
<i>Pittosporum undulatum</i>	Pittosporum				5		X	X	
<b>Proteaceae</b>									
# <i>Grevillea robusta</i>	Silky Oak			5					
<b>Rubiaceae</b>									
* <i>Rothmannia globosa</i>	Tree Gardenia								X
<b>Solanaceae</b>									
<i>Solanum aviculare</i>	Kangaroo Apple					X			
<b>Sterculiaceae</b>									
# <i>Brachychiton acerifolius</i>	Illawarra Flame-tree, Flame Kurrajong						X		
<b>Ulmaceae</b>									
* <i>Celtis occidentalis</i>	American Hackberry					X			
* <i>Celtis sinensis</i>	Chinese Hackberry, Chinese Nettle-tre			1					
<b>Verbenaceae</b>									
* <i>Lantana camara</i>	Lantana	60	90	80	80		X	X	

Botanical name	Common name	Sub-quadrats				Spot locs.		GT locs.	
		1-1	1-2	1-3	1-4	A	B	GTa	GTb
<b>Vitaceae</b>									
Cissus antarctica	Kangaroo Vine					X		X	
<b>4. Monocotyledons</b>									
<b>Agavaceae</b>									
* Yucca sp.	Yucca				0.1				
<b>Arecaceae</b>									
* Phoenix canariensis	Canary Island Date								X
<b>Asparagaceae</b>									
* Asparagus aethiopicus	Asparagus Fern			1	1				X
* Asparagus plumosus	Climbing Asparagus Fern			1				X	
<b>Commelinaceae</b>									
* Tradescantia fluminensis	Trad, Wandering Jew						X	X	X
<b>Lomandraceae</b>									
Lomandra longifolia	Honey Reed, Spike Mat-rush							X	
<b>Poaceae</b>									
* Ehrharta erecta	Panic Veld-grass					X			

**Table 4b - Species recorded in sub-quadrats and at supplementary sampling locations in Brush Farm Park**

Note 1. Asterisk (\*) before botanical name signifies exotic species. Hash symbol (#) signifies a non-local native, planted or naturalised.

2. Families are grouped under headings 1. Pteridophytes, 2. Gymnosperms, 3. Dicotyledons, 4. Monocotyledons. One or more of these plant groups may be absent from this site.

3. The numbers in the columns for sub-quadrats denote percent (%) projected foliage cover in each of the sub-quadrats within the 20 m x 20 m or 10 m x 40 m quadrats. For the spot locations and ground truthing locations, presence only is indicated (X).

4. 'GT locs.' = Ground truthing locations.

Botanical name	Sub-quadrats												GT locs.						
	1-1	1-2	1-3	1-4	2-1	2-2	2-3	2-4	3-1	3-2	3-3	3-4	4-1	4-2	4-3	4-4	GTa	GTb	GTc
<b>1. Pteridophytes</b>																			
<b>Adiantaceae</b>																			
<i>Adiantum hispidulum</i>					15	20	15	40									X		X
<i>Pellaea falcata</i>																			X
<b>Blechnaceae</b>																			
<i>Doodia aspera</i>																	X		X
<i>Doodia caudata</i>						0.1													X
<b>Cyatheaceae</b>																			
# <i>Cyathea cooperi</i>																	X		X
<b>Dicksoniaceae</b>																			
<i>Dicksonia antarctica</i>																			X.
<b>Pteridaceae</b>																			
<i>Pteris tremula</i>										0.1	0.1	0.1					X	X	X
<b>Thelypteridaceae</b>																			
<i>Cyclosorus dentatus</i>					1														X
<b>3. Dicotyledons</b>																			
<b>Acanthaceae</b>																			
<i>Pseuderanthemum variabile</i>		0.1	0.1		0.1				0.1	0.1	0.1	0.1	0.1	1	5	1			X

Botanical name	Sub-quadrats													GT locs.					
	1-1	1-2	1-3	1-4	2-1	2-2	2-3	2-4	3-1	3-2	3-3	3-4	4-1	4-2	4-3	4-4	GTa	GTb	GTc
<b>Aphanopetalaceae</b>																			
Aphanopetalum resinosum	3		5	10							0.1								
<b>Apocynaceae</b>																			
* Araujia sericifera						0.1	0.1												
<b>Araliaceae</b>																			
* Hedera helix		0.1							5										
<b>Asteraceae</b>																			
* Bidens pilosa	1											0.1	0.1	0.1					
Sigesbeckia orientalis													0.1						
<b>Basellaceae</b>																			
* Anredera cordifolia	1	1						0.1	0.1									X	
<b>Bignoniaceae</b>																			
* Jacaranda mimosifolia				5															
Pandorea pandorana												0.1	2	0.1		X			
<b>Celastraceae</b>																			
Celastrus australis														0.1		X			
Denhamia silvestris															1				
Elaeodendron australe					10	1													X
<b>Convolvulaceae</b>																			
Calystegia marginata			0.1															X	
* Ipomoea indica	2		0.1															X	
<b>Cunoniaceae</b>																			
Schizomeria ovata									50										X
<b>Euphorbiaceae</b>																			
Breynia oblongifolia		4		1			5						0.1	0.1		X			
Glochidion ferdinandi					1	1													X
Homalanthus populifolius																X		X	
Phyllanthus gunnii																		X	

Botanical name	Sub-quadrats													GT locs.					
	1-1	1-2	1-3	1-4	2-1	2-2	2-3	2-4	3-1	3-2	3-3	3-4	4-1	4-2	4-3	4-4	GTa	GTb	GTc
<b>Eupomatiaceae</b>																			
Eupomatia laurina					2	15													X
<b>Fabaceae Caesalpinoideae</b>																			
* Senna pendula var. glabrata			1																
<b>Fabaceae Faboideae</b>																			
Glycine clandestina												0.1							
Glycine tabacina												0.1	0.1						
Indigofera australis												0.1	0.1						
<b>Fabaceae Mimosoideae</b>																			
Acacia decurrens		3																	
Acacia parramattensis		5	1									8	10	0.1	5				X
<b>Lamiaceae</b>																			
Clerodendrum tomentosum	1							0.1											
Plectranthus parviflorus		1	0.1									1	1		0.1				X
<b>Lauraceae</b>																			
* Cinnamomum camphora	0.1			0.1					2								X		X
Cryptocarya glaucescens																			
<b>Malvaceae</b>																			
Hibiscus heterophyllus																			X
* Sida rhombifolia												0.1							
<b>Meliaceae</b>																			
# Melia azedarach		1																	X
Synoum glandulosum																		X	
# Toona ciliata																			X
<b>Menispermaceae</b>																			
Sarcopetalum harveyanum				0.1	0.1				1		0.1	0.1							X
Stephania japonica var. discolor		0.1				2			2		3	2	0.1	0.1					
<b>Moraceae</b>																			
Ficus coronata				5	30	10	10	10	10	20	40	20					X	X	X

Botanical name	Sub-quadrats																GT locs.		
	1-1	1-2	1-3	1-4	2-1	2-2	2-3	2-4	3-1	3-2	3-3	3-4	4-1	4-2	4-3	4-4	GTa	GTb	GTc
<i>Ficus rubiginosa</i>	15	3						1	10										
* <i>Morus alba</i>		0.1																	
<b>Myrsinaceae</b>																			
<i>Myrsine variabilis</i>					5	2	3								1				X
<b>Myrtaceae</b>																			
<i>Acmena smithii</i>			3			3										X			X
<i>Backhousia myrtifolia</i>																			X
<i>Callistemon salignus</i>	1		4																X
<i>Eucalyptus acmenoides</i>								30											
# <i>Eucalyptus grandis</i>	5																		X
# <i>Eucalyptus microcorys</i>	10		10																
<i>Eucalyptus paniculata</i>								30								X			
<i>Eucalyptus saligna</i>	3	3	20							10									
<i>Melaleuca styphelioides</i>												0.1	0.1						
<i>Syncarpia glomulifera</i>	1	1		20		10	5			20			20	20	60			X	
<i>Syzygium australe</i>																		X	
<b>Ochnaceae</b>																			
* <i>Ochna serrulata</i>					1	0.1	3	2							0.1				
<b>Oleaceae</b>																			
* <i>Ligustrum lucidum</i>			0.1									0.1	0.1			2	X		
<i>Notelaea longifolia</i>	1			2								1	5		2				
* <i>Olea europaea</i> subsp. <i>cuspidata</i>														0.1	0.1		X		
<b>Oxalidaceae</b>																			
* <i>Oxalis pes-caprae</i>			0.1																
<b>Passifloraceae</b>																			
<i>Passiflora herbertiana</i>						0.1						3	10	2	1	X			
* <i>Passiflora suberosa</i>												0.1	5		2				
* <i>Passiflora subpellata</i>			0.1			0.1						0.1		1					
<b>Pittosporaceae</b>																			
<i>Bursaria spinosa</i>		1																	
<i>Pittosporum multiflorum</i>		0.1		2								3	3	0.1					

Botanical name	Sub-quadrats																GT locs.		
	1-1	1-2	1-3	1-4	2-1	2-2	2-3	2-4	3-1	3-2	3-3	3-4	4-1	4-2	4-3	4-4	GTa	GTb	GTc
<i>Pittosporum undulatum</i>	3	40	10	5	5	15	3	20	3	5	20	3	3	5	10	X	X	X	
<b>Proteaceae</b>																			
# <i>Grevillea robusta</i>								0.1		2									
# <i>Stenocarpus sinuatus</i>					0.1			0.1	0.1									X	
<b>Ranunculaceae</b>																			
<i>Clematis glycinoides</i>	3	2	3									0.1	0.1	0.1	1				
<b>Rosaceae</b>																			
<i>Rubus rosifolius</i>																X			
* <i>Rubus</i> sp.																	X		
<b>Rubiaceae</b>																			
<i>Morinda jasminoides</i>		1			2	2	3											X	
<b>Rutaceae</b>																			
<i>Melicope micrococca</i>		1			1	15	10									X			
<b>Sapindaceae</b>																			
<i>Alectryon subcinerus</i>				40	2		3											X	
<i>Guioa semiglauc</i>				2			1												
<b>Solanaceae</b>																			
<i>Solanum aviculare</i>															0.1				
* <i>Solanum mauritianum</i>																X			
<i>Solanum prinophyllum</i>															0.1				
* <i>Solanum pseudocapsicum</i>				0.1		1						0.1		0.1	0.1	X			
<b>Sterculiaceae</b>																			
# <i>Brachychiton acerifolius</i>				2		3	5	4		5		2	2	1	0.1	X			
<b>Ulmaceae</b>																			
* <i>Celtis occidentalis</i>								2											
* <i>Celtis sinensis</i>																			
<i>Trema tomentosa</i> var. <i>aspera</i>			2	0.1												X	X		
<b>Urticaceae</b>																			
* <i>Parietaria judaica</i>		3																	



Botanical name	Sub-quadrats													GT locs.					
	1-1	1-2	1-3	1-4	2-1	2-2	2-3	2-4	3-1	3-2	3-3	3-4	4-1	4-2	4-3	4-4	GTa	GTb	GTc
<b>Verbenaceae</b>																			
* Lantana camara															4	15			
<b>Vitaceae</b>																			
Cayratia clematidea	0.1	0.1				0.1			0.1	0.1	1	0.1	1	2	2	1			X
Cissus antarctica			0.1			2	10	5	2		2		0.1	0.1			X		X
<b>4. Monocotyledons</b>																			
<b>Anthericaceae</b>																			
* Chlorophytum comosum														2	5	40			
<b>Araceae</b>																			
Alocasia brisbanensis																	X		
<b>Areaceae</b>																			
Livistona australis									1			5					X	X	X
* Phoenix canariensis		0.1																	
* Syagrus romanzoffiana															0.1				
<b>Asparagaceae</b>																			
* Asparagus aethiopicus									0.1				0.1	1	1	1			
* Asparagus plumosus						0.1													
<b>Asphodelaceae</b>																			
* Aloe arborescens																			
<b>Commelinaceae</b>																			
Commelina cyanea	2	10					0.1												X
* Tradescantia fluminensis	30	40	80	30			0.1	30							2		X	X	
<b>Cyperaceae</b>																			
Cyperus tetraphyllus										1									X
Gahnia melanocarpa																	X		
<b>Lomandraceae</b>																			
# Lomandra hystrix																			X
Lomandra longifolia													0.1	0.1					

Botanical name	Sub-quadrats													GT locs.					
	1-1	1-2	1-3	1-4	2-1	2-2	2-3	2-4	3-1	3-2	3-3	3-4	4-1	4-2	4-3	4-4	GTa	GTb	GTc
<b>Philesiaceae</b>																			
<i>Eustrephus latifolius</i>	1			0.1	0.1	1	0.1	1	1		0.1		1		1		X		X
<b>Phormiaceae</b>																			
<i>Dianella caerulea</i>														2			X		
<b>Poaceae</b>																			
* <i>Ehrharta erecta</i>	20	30	0.1	10		0.1		5	5	1	1	1	30	80	50	30	X	X	
<i>Entolasia marginata</i>									0.1				0.1	0.1					
<i>Entolasia stricta</i>																0.1			
<i>Microlaena stipoides</i>						0.1							0.1	10					
<i>Opismenus aemulus</i>		0.1	0.1						0.1		0.1								
<i>Opismenus imbecillis</i>						1	0.1						0.1	0.1		1			X
<b>Smilacaceae</b>																			
<i>Smilax australis</i>					0.1	1					3	0.1					X		X

**Table 4c - Species recorded in sub-quadrats and at supplementary sampling locations in Darvall Park**

Notes: 1. Asterisk (\*) before botanical name signifies exotic species. Hash symbol (#) signifies a non-local native, planted or naturalised.

2. Families are grouped under headings 1. Pteridophytes, 2. Gymnosperms, 3. Dicotyledons, 4. Monocotyledons. One or more of these plant groups may be absent from this site.

3. The numbers in the columns for sub-quadrats denote percent (%) projected foliage cover in each of the sub-quadrats within the 20 m x 20 m or 10 m x 40 m quadrats. For the spot locations and ground truthing locations, presence only is indicated (X).

4. 'GT locs.' = Ground truthing locations.

Botanical name	Common name	Sub-quadrats							GT locs.																		
		1-1	1-2	1-3	1-4	GTa	GTb	GTc	GTd	GTe	GTf	GTg															
<b>1. Pteridophytes</b>																											
<b>Blechnaceae</b>																											
Doodia aspera	Prickly Rasp Fern																		X								
<b>3. Dicotyledons</b>																											
<b>Acanthaceae</b>																											
Pseuderanthemum variabile	Pastel Flower		0.1																	X							
* Thunbergia alata	Black-eyed Susan			0.1	0.1																						
<b>Anacardiaceae</b>																											
* Toxicodendron succedaneum	Rhus Tree, Wax Tree		2																								
<b>Apiaceae</b>																											
Centella asiatica	Indian Pennywort		0.1																								
Hydrocotyle hirta	Hairy Pennywort																				X						
<b>Apocynaceae</b>																											
* Araujia sericifera	Moth Vine, Cruel Plant				0.1																						
<b>Araliaceae</b>																											
Polyscias sambucifolia	Elderberry Panax																				X						
# Schefflera actinophylla	Queensland Umbrella Tree								X																		

Botanical name	Common name	Sub-quadrats							GT locs.																				
		1-1	1-2	1-3	1-4	GTa	GTb	GTc	GTd	GTe	GTf	GTg																	
<b>Asteraceae</b>																													
* Bidens pilosa	Cobbler's Pegs		0.1																										
Ozothamnus diosmifolius	White Dogwood																							X					
Sigesbeckia orientalis	Indian Weed	0.1		0.1	1																								
<b>Basellaceae</b>																													
* Anredera cordifolia	Madeira Vine, Lamb's Tail																							X					
<b>Bignoniaceae</b>																													
Pandorea pandorana	Wonga Vine																						X						
<b>Celastraceae</b>																													
Denhamia silvestris	Narrow-leaved Orangebark, Orange Bush,																						X						
Elaeodendron australe	Red Olive-plum				2																	X							
<b>Convolvulaceae</b>																													
Dichondra sp. A	Hairy Kidney Weed																							X					
* Ipomoea indica	Blue Morning Glory										X											X							
<b>Euphorbiaceae</b>																													
Glochidion ferdinandi	Cheese Tree									X																			
Homalanthus populifolius	Bleeding heart, Native Poplar	1																					X	X					
Phyllanthus gunnii	Scrubby Spurge		2		0.1																								
<b>Fabaceae Faboideae</b>																													
Glycine clandestina	Twining Glycine				1																		X						
Glycine tabacina					1																								
Indigofera australis	Native Indigo																						X						
Kennedia rubicunda	Dusky Coral-pea									X																			
<b>Fabaceae Mimosoideae</b>																													
Acacia floribunda	White Sally Wattle																							X					
Acacia parramattensis	Parramatta Green Wattle	0.1			0.1																	X	X						
<b>Geraniaceae</b>																													
Geranium homeanum	Rainforest Cranesbill		5	2	15	3							X																

Botanical name	Common name	Sub-quadrats							GT locs.																					
		1-1	1-2	1-3	1-4	GTa	GTb	GTc	GTd	GTe	GTf	GTg																		
<b>Lamiaceae</b>																														
<i>Plectranthus parviflorus</i>	Cockspur Flower	2	5	1	1																									
<b>Lobeliaceae</b>																														
<i>Pratia purpurascens</i>	Whiteroot			1	0.1																									
<b>Malvaceae</b>																														
* <i>Pavonia hastata</i>																							X							
* <i>Sida rhombifolia</i>	Paddy's Lucerne		0.1	0.1	1																									
<b>Menispermaceae</b>																														
<i>Sarcopetalum harveyanum</i>	Pearl Vine																						X							
<i>Stephania japonica</i> var. <i>discolor</i>	Snake Vine	0.1	0.1		1																		X							
<b>Moraceae</b>																														
<i>Ficus coronata</i>	Creek Sandpaper Fig						X																							
<b>Myrtaceae</b>																														
<i>Acmena smithii</i>	Lilly-pilly				2					X			X										X							
<i>Backhousia myrtifolia</i>	Grey Myrtle, Ironwood						X																X							
<i>Callistemon salignus</i>	White Bottlebrush, Pink-tips			1	2																									
<i>Eucalyptus acmenoides</i>	White Mahogany										X												X							
# <i>Eucalyptus grandis</i>	Flooded Gum	10						X															X							
# <i>Eucalyptus microcorys</i>	Tallowwood			5				X																						
<i>Eucalyptus paniculata</i>	Grey Ironbark	10							10																					
<i>Eucalyptus pilularis</i>	Blackbutt	10																												
<i>Eucalyptus saligna</i>	Sydney Blue Gum		15	30	10					X			X										X							
# <i>Eucalyptus tereticornis</i>	Forest Red Gum																						X							
# <i>Lophostemon confertus</i>	Brush Box									X																				
<i>Melaleuca styphelioides</i>	Prickly Paperbark		5	0.1						X			X										X							
<i>Syncarpia glomulifera</i>	Turpentine		5		5					X			X										X							
<b>Oleaceae</b>																														
* <i>Ligustrum lucidum</i>	Broad-leaved Privet, Glossy Privet																						X							
<i>Notelaea longifolia</i>	Large Mock-olive	1	0.1										X										X							

Botanical name	Common name	Sub-quadrats					GT locs.						
		1-1	1-2	1-3	1-4	GTa	GTb	GTc	GTd	GTe	GTf	GTg	
<b>Passifloraceae</b>													
* <i>Passiflora suberosa</i>	Corky Passionflower		2	1									
<b>Pittosporaceae</b>													
<i>Pittosporum revolutum</i>	Yellow Pittosporum		5			X	X				X		
<i>Pittosporum undulatum</i>	Pittosporum						X					X	
<b>Proteaceae</b>													
# <i>Macadamia tetraphylla</i>	Bopple Nut, Rough-shelled Macadamia										X		
<b>Ranunculaceae</b>													
<i>Clematis glycinoides</i>	Headache Vine, Traveller's Joy, Old Man's			1	2					X	X		
<b>Rosaceae</b>													
* <i>Rubus laudatus</i>	Plains Blackberry		0.1	2	2								
<i>Rubus rosifolius</i>	Native Raspberry, Rose-leaf Bramble					X	X					X	
<b>Rubiaceae</b>													
* <i>Galium aparine</i>	Cleavers, Goose-grass, Bedstraw											X	
<b>Sapindaceae</b>													
* <i>Cardiospermum grandiflorum</i>	Balloon Vine		5										
<i>Dodonaea triquetra</i>	Hopbush											X	
<b>Solanaceae</b>													
* <i>Cestrum nocturnum</i>	Night-scented Jessamine, Lady of the Nigh			1									
<i>Solanum aviculare</i>	Kangaroo Apple										X		
* <i>Solanum chenopodioides</i>	White-tip Nightshade				0.1								
* <i>Solanum mauritianum</i>	Tree Tobacco, Wild Tobacco											X	
* <i>Solanum seaforthianum</i>	Brazilian Nightshade		2	1	0.1								
* <i>Solanum sp.</i>			1										
<b>Sterculiaceae</b>													
# <i>Brachychiton acerifolius</i>	Illawarra Flame-tree, Flame Kurrajong					X					X		
<b>Ulmaceae</b>													
<i>Trema tomentosa</i> var. <i>aspera</i>	Poison Peach, Peach-leaf Poison Bush										X		

Botanical name	Common name	Sub-quadrats							GT locs.				
		1-1	1-2	1-3	1-4	GTa	GTb	GTc	GTd	GTe	GTf	GTg	
<b>Verbenaceae</b>													
* Lantana camara	Lantana							X	X	X		X	
<b>Vitaceae</b>													
Cayratia clematidea	Slender Grape	5	10	2	1	X				X			
Cissus antarctica	Kangaroo Vine	30	40	3	2		X			X	X		
Cissus hypoglauca	Native Grape, Water Vine									X	X		
<b>4. Monocotyledons</b>													
<b>Alstroemeriaceae</b>													
* Alstroemeria pulchella	Parrot Alstroemeria						X						
<b>Araceae</b>													
Alocasia brisbanensis	Cunjevoi Lily						X						
* Colocasia esculenta	Taro, Dasheen											X	
<b>Asparagaceae</b>													
* Asparagus aethiopicus	Asparagus Fern						0.1						
* Asparagus officinalis	Asparagus	1											
<b>Commelinaceae</b>													
Commelina cyanea	Blue Spiderwort	5	2	3	0.1					X			
* Tradescantia fluminensis	Trad, Wandering Jew	60	70	30	10	X	X	X	X	X	X	X	
<b>Cyperaceae</b>													
Cyperus tetrathyllus											X		
Gahnia melanocarpa	Black-fruit Saw-sedge							X					
<b>Lomandraceae</b>													
Lomandra longifolia	Honey Reed, Spike Mat-rush	0.1	5	5	5						X		
<b>Musaceae</b>													
* Musa acuminata	Banana											X	
<b>Philesiaceae</b>													
Eustrephus latifolius	Wombat Berry										X		

Botanical name	Common name	Sub-quadrats							GT locs.																			
		1-1	1-2	1-3	1-4	GTa	GTb	GTc	GTd	GTe	GTf	GTg																
<b>Phormiaceae</b>																												
<i>Dianella caerulea</i>	Blue Flax-lily		0.1								X												X					
<b>Poaceae</b>																												
* <i>Ehrharta erecta</i>	Panic Veld-grass	10	10	15	2	X					X		X	X	X	X	X	X	X	X	X	X	X	X				
* <i>Megathyrsus maximus</i>	Guinea Grass		5																									
<i>Microlaena stipoides</i>	Weeping Grass, Meadow Rice-grass	10		1	5						X		X	X	X	X	X	X	X	X	X	X	X	X				
<i>Oplismenus aemulus</i>	Australian Basket Grass, Wavy Beard Gra	5	15	5	15	X																						
<i>Oplismenus imbecillis</i>	Narrow-leaved Basket Grass	2																										
* <i>Paspalum dilatatum</i>	Paspalum				0.1																							























**Table 4d-2 - Species recorded at supplementary ground truthing quadrats in Field of Mars Reserve**

Notes: 1. Asterisk (\*) before botanical name signifies exotic species. Hash symbol (#) signifies a non-local native, planted or naturalised.

2. Families are grouped under headings 1. Pteridophytes, 2. Gymnosperms, 3. Dicotyledons, 4. Monocotyledons. One or more of these plant groups may be absent from this site.

3. The numbers in the columns for sub-quadrats denote percent (%) projected foliage cover in each of the 10 m x 10 m sub-quadrats within the 20 m x 20 m or 10 m x 40 m quadrats. The numbers in the columns for GT Quadrats denote average percent (%) projected foliage cover for each quadrat, from the percent (%) projected foliage cover recorded in the four sub-quadrats.

4. 'GT Quadrats' = Ground truthing quadrats.

Botanical name	Sub-quadrats												GT Quadrats							
	A-1	A-2	A-3	A-4	B-1	B-2	B-3	B-4	C-1	C-2	C-3	C-4	D-1	D-2	D-3	D-4	A	B	C	D
<b>1. Pteridophytes</b>																				
<b>Adiantaceae</b>																				
Cheilanthes sieberi													0.1							0.025
<b>Cyatheaceae</b>																				
# Cyathea cooperi			5	2													1.75			
<b>Davalliaceae</b>																				
# Nephrolepis cordifolia		2															0.5			
<b>Dennstaedtiaceae</b>																				
Hypolepis muelleri			1														0.25			
Pteridium esculentum			0.1					2	3	5	5	1	1			1	0.025	3.75	0.5	
<b>Dicksoniaceae</b>																				
Calochlaena dubia												5							1.25	
<b>3. Dicotyledons</b>																				
<b>Acanthaceae</b>																				
Brunoniella pumilio																	0.1			0.025
Pseuderanthemum variabile													0.1	1	0.1	0.1				0.325

Botanical name	Sub-quadrats												GT Quadrats							
	A-1	A-2	A-3	A-4	B-1	B-2	B-3	B-4	C-1	C-2	C-3	C-4	D-1	D-2	D-3	D-4	A	B	C	D
<b>Apiaceae</b>																				
Centella asiatica			0.1														0.025			
Platsace lanceolata											1								0.25	
Xanthosia pilosa								0.1	0.1	1	0.1								0.325	
<b>Araliaceae</b>																				
Polyscias sambucifolia	3	3	1	1				0.1	1	1	1	1	1	1	1	2	0.025	1	0.75	
<b>Asteraceae</b>																				
Ozothamnus diosmifolius											0.1								0.025	
<b>Bignoniaceae</b>																				
Pandorea pandorana	1	1	2						0.1	0.1		2	1	2	1	1		0.05	1.5	
<b>Casuarinaceae</b>																				
Allocasuarina littoralis	2	3	1	3	30	15	15	2	5	5	5	5	15	2		2.25	22.5	4.25	5.5	
Allocasuarina torulosa													2		3				1.25	
<b>Dilleniaceae</b>																				
Hibbertia aspera													1	2						0.75
Hibbertia riparia					1	1	1	1										1		
<b>Elaeocarpaceae</b>																				
Elaeocarpus reticulatus	3	2	1					0.1	0.1	2			0.1				1.5	0.025	0.525	0.025
<b>Ericaceae Styphelioideae</b>																				
Epactis pulchella					1	2	3	1									1.75			
Leucopogon juniperinus												2	2	3	1					2
Leucopogon lanceolatus										1									0.25	
Monotoca scoparia								1											0.25	
Woolisia pungens										0.1									0.025	
<b>Euphorbiaceae</b>																				
Breynia oblongifolia	2	2	2	3									1			1	2.25			0.5
Glochidion ferdinandi	1	5	1									0.1	1	1	1	1.75			0.775	
Homalanthus populifolius	1	2		0.1						0.1		0.1				0.775			0.025	0.025
Micranthemum ericoides								0.1		0.1									0.025	0.025
Phyllanthus hirtellus					1			1		0.1	0.1	1	1	1			0.5	0.05	0.5	

Botanical name	Sub-quadrats												GT Quadrats							
	A-1	A-2	A-3	A-4	B-1	B-2	B-3	B-4	C-1	C-2	C-3	C-4	D-1	D-2	D-3	D-4	A	B	C	D
<b>Fabaceae Caesalpinioideae</b>																				
* <i>Senna pendula</i> var. <i>glabrata</i>	1	1	1	1												2	0.75			0.5
<b>Fabaceae Faboideae</b>																				
<i>Bossiaea obcordata</i>	1					1	1	1			0.1						0.25	0.75	0.025	
<i>Bossiaea scolopendria</i>							0.1	0.1										0.05		
<i>Dillwynia retorta</i>							1											0.25		
<i>Glycine clandestina</i>					1	1	1											0.75		
<i>Hovea linearis</i>						0.1		0.1										0.05		
<i>Mirbelia rubifolia</i>					0.1													0.025		
<i>Platylobium formosum</i>	0.1	1						10	7	5	10	2	1	1	1	1	0.275		8	1.25
<i>Pultenaea paleacea</i>					1													0.25		
<i>Pultenaea tuberculata</i>					1	1	1	0.1										0.775		
<b>Fabaceae Mimosoideae</b>																				
<i>Acacia fimbriata</i>						1	2											0.75		
<i>Acacia linifolia</i>								1	0.1	1			1					0.525	0.25	
<i>Acacia longifolia</i>	1			1												0.5				
<i>Acacia myrtifolia</i>																		0.025		
<i>Acacia suaveolens</i>																		0.025		
<i>Acacia terminalis</i> subsp. <i>angustifolia</i>													1							0.25
<b>Goodeniaceae</b>																				
<i>Goodenia bellidifolia</i>													0.1							0.025
<i>Goodenia hederacea</i>						0.1	0.1	1	1				0.1					0.55		0.025
<b>Haloragaceae</b>																				
<i>Gonocarpus tetragynus</i>	1																0.5			
<i>Gonocarpus teucrioides</i>										1	1	1	0.1						0.775	
<b>Lauraceae</b>																				
<i>Cassyltha pubescens</i>						2	1	1		2	1	1	1	1					1	1.25
* <i>Cinnamomum camphora</i>	1				0.1				0.1	0.1			0.1				0.275	0.025	0.05	0.025
<b>Lobeliaceae</b>																				
<i>Pratia purpurascens</i>	0.1					0.1							0.1	0.1			0.025	0.025		0.05
<b>Myrtaceae</b>																				
<i>Angophora costata</i>	20	10	10	5	15			10	35	35	30	20	25	10	30	20	11.25	6.25	30	21.25

Botanical name	Sub-quadrats										GT Quadrats									
	A-1	A-2	A-3	A-4	B-1	B-2	B-3	B-4	C-1	C-2	C-3	C-4	D-1	D-2	D-3	D-4	A	B	C	D
<i>Conyobia gummifera</i>					5	20	10	20										13.75		
<i>Eucalyptus globoidea</i>							20	10					15					7.5		3.75
<i>Eucalyptus resinifera</i> subsp. <i>resinifera</i>			5		0.1	10	20	5						2			1.25	8.775		0.5
<i>Leptospermum polygalifolium</i> subsp. <i>polygalif</i>	1																0.25			
<i>Leptospermum trinervium</i>							2	1										0.5	0.25	
<i>Melaleuca linariifolia</i>			5														1.25			
<i>Syncarpia glomulifera</i>													15	15	10					10
<b>Nandinaaceae</b>																				
* <i>Nandina domestica</i>		0.1															0.025			
<b>Ochnaceae</b>																				
* <i>Ochna serrulata</i>	1	2	2			0.1	0.1	0.1	0.1				0.1	0.1	1		1.25	0.05	0.025	0.3
<b>Oleaceae</b>																				
* <i>Ligustrum lucidum</i>	0.1	1														1	0.275			0.25
* <i>Ligustrum sinense</i>	1	2	2	1									1	1		1	1.5			0.75
<i>Notelaea longifolia</i>	2		2	2	1	1							1	0.1	1		1.5	0.5		0.525
<b>Oxalidaceae</b>																				
<i>Oxalis exilis</i>													0.1							0.05
<b>Pittosporaceae</b>																				
<i>Billardiera scandens</i>					0.1	0.1	1	0.1	0.1	0.1	0.1	1	0.1					0.325	0.325	0.025
<i>Pittosporum revolutum</i>													5	3	50	40				24.5
<i>Pittosporum undulatum</i>	10	25	10	30			1		0.1		1	1					18.75	0.25	0.525	
<b>Proteaceae</b>																				
<i>Banksia spinulosa</i>	1	1							1	2			2	1			0.5		0.75	0.75
<i>Grevillea buxifolia</i>							1											0.25		
<i>Grevillea sericea</i>					1	1	1											0.75		
<i>Lomatia siliifolia</i>	1	1			0.1	1	1	1	0.1	1	1	1	0.1			1	0.5	0.775	0.775	0.275
<i>Persoonia lanceolata</i>					1	1												0.5		
<i>Persoonia laurina</i> subsp. <i>laurina</i>						0.1											0.025			
<i>Persoonia levis</i>							1	2		2									1.25	
<i>Persoonia linearis</i>									1		1								0.25	
<i>Xylomelum pyriforme</i>							1			1								0.25	0.25	
<b>Ranunculaceae</b>																				
<i>Clematis glycinoides</i>	1	1	2	1					0.1				1	1	1		1.25		0.025	0.75

Botanical name	Sub-quadrats												GT Quadrats							
	A-1	A-2	A-3	A-4	B-1	B-2	B-3	B-4	C-1	C-2	C-3	C-4	D-1	D-2	D-3	D-4	A	B	C	D
<b>Rhamnaceae</b>																				
Pomaderris lanigera													1							0.25
<b>Rubiaceae</b>																				
Opercularia diphylla						0.1	0.1											0.05		
<b>Sapindaceae</b>																				
Dodonaea triquetra								0.1	1	1	1	2	2	1					0.525	1.25
<b>Verbenaceae</b>																				
* Lantana camara	1	1		2								1		2		1				0.75
<b>4. Monocotyledons</b>																				
<b>Anthericaceae</b>																				
Tricoryne simplex													0.1	0.1						0.05
<b>Arecaceae</b>																				
* Syagrus romanzoffiana										0.1						0.1		0.025		0.025
<b>Asparagaceae</b>																				
* Asparagus aethiopicus				1											0.1		0.25			0.025
<b>Colchicaceae</b>																				
Burchardia umbellata													0.1					0.025		0.025
<b>Cyperaceae</b>																				
Carex breviculmis			0.1														0.025			
Cyathochaeta diandra								0.1					1					0.025	0.25	
Gahnia clarkei				1													0.25			
Lepidosperma laterale	3	1		1	2	2	3	2	2	3	3	3	1		3	1	1.25	2.25	2.75	1.25
Ptilothrix deusta										0.1								0.025		
Schoenus apogon				1													0.25			
<b>Iridaceae</b>																				
Patersonia sericea										0.1	0.1	0.1						0.075		
<b>Liliaceae</b>																				
* Lilium formosanum			0.1	0.1														0.05		

Botanical name	Sub-quadrats												GT Quadrats							
	A-1	A-2	A-3	A-4	B-1	B-2	B-3	B-4	C-1	C-2	C-3	C-4	D-1	D-2	D-3	D-4	A	B	C	D
<b>Lomandraceae</b>																				
<i>Lomandra cylindrica</i>							0.1	0.1									0.05			
<i>Lomandra filiformis</i> subsp. <i>filiformis</i>	0.1	0.1						0.1	0.1	0.1			0.1	0.1			0.05	0.075	0.05	
<i>Lomandra gracilis</i>	0.1									0.1							0.025		0.025	
<i>Lomandra longifolia</i>	15	20	1	10				25	10	15	20	5	5	2	10	11.5		17.5	5.5	
<i>Lomandra multiflora</i>	0.1	0.1	1				0.1	0.1		0.1			0.1			0.3	0.05	0.05	0.025	
<i>Lomandra obliqua</i>							0.1										0.025			
<b>Orchidaceae</b>																				
<i>Acianthus</i> sp.								0.1		0.1									0.05	
<i>Calochilus paludosus</i>							0.1											0.025		
<i>Cryptostylis erecta</i>										0.1									0.025	
<b>Phormiaceae</b>																				
<i>Dianella caerulea</i>	1	10	3		1	1	1	1	1	1	1	1	1	1	2	1	3.5	0.75	1	1.25
<i>Dianella revoluta</i>	0.1		0.1		1	0.1				0.1		1					0.05	0.275	0.025	0.25
<b>Poaceae</b>																				
<i>Aristida vagans</i>							0.1	0.1										0.05		
<i>Austrostipa pubescens</i>					2				2	0.1			2	2		1		0.5	0.525	1.25
<i>Eriolasia stricta</i>	3	1	5		25	25	30	15	3	2	3	2	5	10	10	2.25	23.75	2.5	8.75	
<i>Imperata cylindrica</i>	5	10	20	5				3	1	1	0.1	5	5	20	5	10		1.275	7.5	
<i>Microlaena stipoides</i>	15	5	2	2	10	10	10	10	1	1	0.1	2	10	25	10	6	10	1.025	13.75	
<i>Oplismenus aemulus</i>																0.25				
<i>Oplismenus imbecillis</i>		1											0.1			0.25			0.025	
<i>Panicum simile</i>								0.1										0.025		
<i>Paspalidium distans</i>													0.1						0.025	
* <i>Paspalum urvillei</i>										0.1						0.025				
<i>Poa sieberiana</i>													0.1							0.05
<i>Themeda triandra</i>	1	1	1		10	5	15	5	2	1			15	15	2	2	0.75	8.75	0.75	8.5
<b>Smilacaceae</b>																				
<i>Smilax glycyphylla</i>	1							2	5	3	2				0.1	1	0.25		3	0.275
<b>Xanthorrhoeaceae</b>																				
<i>Xanthorrhoea arborea</i>										1									0.5	
<i>Xanthorrhoea media</i>	2	2		1	5	2	1	2				1	0.1	0.1	1	0.1	1.25	2.5		0.325

**Table 4e - Species recorded in sub-quadrats and at supplementary sampling locations in Lambert Park**

Notes: 1. Asterisk (\*) before botanical name signifies exotic species. Hash symbol (#) signifies a non-local native, planted or naturalised.

2. Families are grouped under headings 1. Pteridophytes, 2. Gymnosperms, 3. Dicotyledons, 4. Monocotyledons. One or more of these plant groups may be absent from this site.

3. The numbers in the columns for subquadrats denote percent (%) projected foliage cover in each of the sub-quadrats within the 20 m x 20 m or 10 m x 40 m quadrats. For the spot locations and ground truthing locations, presence only is indicated (X).

4. 'GT locs.' = Ground truthing locations.

Botanical name	Common name	Sub-quadrats							GT locs.			
		1-1	1-2	1-3	1-4	GTa	GTb	GTc	GTd	GTe	GTf	
<b>1. Pteridophytes</b>												
<b>Adiantaceae</b>												
<i>Adiantum aethiopicum</i>	Common Maidenhair Fern											X
<i>Adiantum hispidulum</i>	Rough Maidenhair Fern		0.1									X
<b>Blechnaceae</b>												
<i>Doodia aspera</i>	Prickly Rasp Fern											X
<b>Cyatheaceae</b>												
# <i>Cyathea cooperi</i>	Straw Tree-fern, Scaly Tree-Fern		1				X					
<b>Pteridaceae</b>												
<i>Pteris tremula</i>	Tender Brake											X
<b>Thelypteridaceae</b>												
<i>Cyclosorus dentatus</i>			0.1		0.1							
<b>3. Dicotyledons</b>												
<b>Aceraceae</b>												
* <i>Acer negundo</i>	Box-elder Maple, Box Elder		0.1									
<b>Aphanopetalaceae</b>												
<i>Aphanopetalum resinosum</i>	Gum Vine		1		2							
<b>Araliaceae</b>												
* <i>Hedera helix</i>	Ivy, English Ivy		0.1		0.1							



Botanical name	Common name	Sub-quadrats				GT locs.				
		1-1	1-2	1-3	1-4	GTa	GTb	GTc	GTd	GTf
<b>Asteraceae</b>										
* Bidens pilosa	Cobbler's Pegs	0.1					X			
* Lactuca serriola	Prickly Lettuce, Compass Plant						X			
* Sonchus oleraceus	Common Sow-thistle, Milk-thistle						X			
<b>Basellaceae</b>										
* Anredera cordifolia	Madeira Vine, Lamb's Tail	0.1	0.1	0.1						
<b>Caryophyllaceae</b>										
* Stellaria media	Chickweed						X			
<b>Casuarinaceae</b>										
# Casuarina cunninghamiana	River Oak, River She-oak		10							
Casuarina glauca	Swamp Oak, Swamp She-oak					X				
<b>Celastraceae</b>										
Elaeodendron australe	Red Olive-plum								X	
<b>Convolvulaceae</b>										
Calystegia marginata		0.1								
<b>Euphorbiaceae</b>										
Breynia oblongifolia	Coffee Bush									X
* Euphorbia peplus	Petty Spurge, Radium Plant						X			
Homalanthus populifolius	Bleeding heart, Native Poplar	15								X
<b>Fabaceae Mimosoideae</b>										
Acacia floribunda	White Sally Wattle								X	
Acacia parramattensis	Parramatta Green Wattle						X		X	X
<b>Lamiaceae</b>										
Plectranthus parviflorus	Cockspur Flower									X
<b>Lauraceae</b>										
* Cinnamomum camphora	Camphor Laurel	0.1	0.1							
<b>Malvaceae</b>										
Hibiscus heterophyllus	Native Rosella									X
<b>Meliaceae</b>										
# Melia azedarach	White Cedar, Persian Lilac	0.1								X

Botanical name	Common name	Sub-quadrats				GT locs.				
		1-1	1-2	1-3	1-4	GTa	GTb	GTc	GTd	GTf
# <i>Toona ciliata</i>	Australian Red Cedar									X
<b>Menispermaceae</b>										
<i>Stephania japonica</i> var. <i>discolor</i>	Snake Vine	1	0.1		0.1					
<b>Moraceae</b>										
<i>Ficus coronata</i>	Creek Sandpaper Fig	10	1	2	1	X			X	X
<i>Ficus rubiginosa</i>	Port Jackson Fig, Rusty Fig		15	40	50	X				
<b>Myrtaceae</b>										
<i>Acmena smithii</i>	Lilly-pilly	60	20	40	30	X	X		X	
<i>Angophora floribunda</i>	Rough-barked Apple									X
<i>Callistemon salignus</i>	White Bottlebrush, Pink-tips								X	
# <i>Callistemon viminalis</i>	Weeping Bottlebrush									X
<i>Eucalyptus globoidea</i>	White Stringybark							X		
# <i>Eucalyptus grandis</i>	Flooded Gum		15				X		X	X
<i>Eucalyptus paniculata</i>	Grey Ironbark							X		
<i>Eucalyptus pilularis</i>	Blackbutt					X				
<i>Eucalyptus saligna</i>	Sydney Blue Gum		20	15	X	X			X	X
# <i>Melaleuca quinquenervia</i>	Broad-leaved Paperbark								X	X
<i>Melaleuca styphelioides</i>	Prickly Paperbark						X	X	X	X
<i>Syncarpia glomulifera</i>	Turpentine					X	X	X	X	X
<b>Nandinaeae</b>										
* <i>Nandina domestica</i>	Sacred Bamboo		0.1							
<b>Ochnaceae</b>										
* <i>Ochna serrulata</i>	Mickey Mouse Plant		0.1							
<b>Oleaceae</b>										
* <i>Jasminum polyanthum</i>	Jasmine									X
* <i>Ligustrum lucidum</i>	Broad-leaved Privet, Glossy Privet		0.1							
* <i>Ligustrum sinense</i>	Small-Leaved Privet, Chinese Privet				0.1					
* <i>Olea europaea</i> subsp. <i>cuspidata</i>	African Olive			0.1						
<b>Passifloraceae</b>										
<i>Passiflora herbiflora</i>	Native Passionfruit		10							
* <i>Passiflora subpeltata</i>	White Passionfruit		2							

Botanical name	Common name	Sub-quadrats							GT locs.			
		1-1	1-2	1-3	1-4	GTa	GTb	GTc	GTd	GTe	GTf	
<b>Pittosporaceae</b>												
Bursaria spinosa	Australian Boxthorn						X					
Pittosporum revolutum	Yellow Pittosporum	0.1	2		1		X					
Pittosporum undulatum	Pittosporum	5	20	3	3	X		X		X	X	
<b>Ranunculaceae</b>												
Clematis glycinoides	Headache Vine, Traveller's Joy, Old Man's Beard	1										
<b>Rosaceae</b>												
* Eriobotrya japonica	Loquat	0.1		0.1								
Rubus moluccanus var. trilobus	Molucca Bramble									X		
Rubus rosifolius	Native Raspberry, Rose-leaf Bramble									X		
<b>Rubiaceae</b>												
Morinda jasminoides	Morinda	0.1	1	1	0.1							
<b>Rutaceae</b>												
Melicope micrococca	Hairy-leaved Doughwood	0.1	3									
# Murraya paniculata	Jasmine-orange, Cosmetic Bark	0.1										
<b>Sapindaceae</b>												
# Alectryon tomentosus	Hairy Bird's-eye, Woolly Rambutan										X	
<b>Solanaceae</b>												
* Cestrum nocturnum	Night-scented Jessamine, Lady of the Night	0.1	2		1							
* Physalis peruviana	Cape Gooseberry	0.1										
* Solanum pseudocapsicum	Jerusalem Cherry										X	
<b>Ulmaceae</b>												
* Celtis sinensis	Chinese Hackberry, Chinese Nettle-tree										X	
Trema tomentosa var. aspera	Poison Peach, Peach-leaf Poison Bush	2					X					
* Ulmus parvifolia	Chinese Elm										X	
<b>Vitaceae</b>												
Cissus antarctica	Kangaroo Vine		1									
<b>4. Monocotyledons</b>												
<b>Araceae</b>												
Alocasia brisbanensis	Cunjevoi Lily										X	

Botanical name	Common name	Sub-quadrats				GT locs.				
		1-1	1-2	1-3	1-4	GTa	GTb	GTc	GTd	GTf
<b>Areaceae</b>										
# Archontophoenix sp.				0.1						
Livistona australis	Cabbage Palm, Cabbage-tree Palm									X
* Phoenix canariensis	Canary Island Date									X
<b>Asparagaceae</b>										
* Asparagus aethiopicus	Asparagus Fern			0.1						X
<b>Commelinaceae</b>										
* Tradescantia fluminensis	Trad, Wandering Jew	30	2		X			X		
<b>Cyperaceae</b>										
Cyperus tetraphyllus		1	0.1	0.1						X
<b>Lomandraceae</b>										
Lomandra longifolia	Honey Reed, Spike Mat-rush		1		X			X		X
<b>Philesiaceae</b>										
Geitonoplesium cymosum	Scrambling Lily	0.1								
<b>Phormiaceae</b>										
Dianella caerulea	Blue Flax-lily	1	0.1	1	X					X
<b>Poaceae</b>										
* Ehrharta erecta	Panic Veld-grass	5	2	0.1	1	X	X	X		X
* Lolium perenne	Perennial Ryegrass							X		
Microlaena stipoides	Weeping Grass, Meadow Rice-grass		1							X
Oplismenus aemulus	Australian Basket Grass, Wavy Beard Grass	0.1		0.1						X
Oplismenus imbecillis	Narrow-leaved Basket Grass	0.1		0.1						
* Pennisetum clandestinum	Kikuyu Grass, Kikuyu								X	

**Table 4f - Species recorded in sub-quadrats and at supplementary sampling locations in Outlook Park**

**Notes:**

1. Asterisk (\*) before botanical name signifies exotic species. Hash symbol (#) signifies a non-local native, planted or naturalised.
2. Families are grouped under headings 1. Pteridophytes, 2. Gymnosperms, 3. Dicotyledons, 4. Monocotyledons. One or more of these plant groups may be absent from this site.
3. The numbers in the columns for sub-quadrats denote percent (%) projected foliage cover in each of the sub-quadrats within the 20 m x 20 m or 10 m x 40 m quadrats. For the ground truthing locations, presence only is indicated (X).
4. 'GT locs.' = Ground truthing locations.

Botanical name	Common name	Sub-quadrats				GT locs.		
		1-1	1-2	1-3	1-4	GTa	GTb	GTc
<b>1. Pteridophytes</b>								
<b>Davalliaceae</b>								
# <i>Nephrolepis cordifolia</i>	Fishbone Fern					X		
<b>2. Gymnosperms</b>								
<b>Araucariaceae</b>								
<i>Araucaria heterophylla</i>	Norfolk Island Pine		0.1					
<b>Pinaceae</b>								
* <i>Cedrus deodara</i>	Deodar, Deodar Cedar					X		
<b>3. Dicotyledons</b>								
<b>Acanthaceae</b>								
* <i>Justicia</i> sp.			0.1					
<i>Pseuderanthemum variabile</i>	Pastel Flower	1	1	0.1	0.1	X		
<b>Araliaceae</b>								
* <i>Hedera helix</i>	Ivy, English Ivy	0.1						
<b>Asteraceae</b>								
* <i>Roldana petasitis</i>	Mexican Giant Groundsel						X	
<i>Sigesbeckia orientalis</i>	Indian Weed			0.1				
<b>Bignoniaceae</b>								
* <i>Jacaranda mimosifolia</i>	Jacaranda		2			X		X
<b>Casuarinaceae</b>								
<i>Allocasuarina torulosa</i>	Forest She-oak			2				
<b>Chenopodiaceae</b>								
<i>Einadia trigonos</i>	Fishweed					X		
<b>Convolvulaceae</b>								
<i>Dichondra</i> sp. A	Hairy Kidney Weed			1		X		
<b>Cunoniaceae</b>								
<i>Ceratopetalum apetalum</i>	Coachwood	7						
<b>Elaeocarpaceae</b>								
<i>Elaeocarpus reticulatus</i>	Blueberry Ash				5			
<b>Euphorbiaceae</b>								
<i>Glochidion ferdinandi</i>	Cheese Tree	2						

Botanical name	Common name	Sub-quadrats				GT locs.		
		1-1	1-2	1-3	1-4	GTa	GTb	GTc
Homalanthus populifolius	Bleeding heart, Native Poplar	1				X		
<b>Eupomatiaceae</b>								
Eupomatia laurina	Bolwarra					X		
<b>Fabaceae Mimosoideae</b>								
Acacia parramattensis	Parramatta Green Wattle					X		
<b>Lamiaceae</b>								
Clerodendrum tomentosum	Hairy Clerodendrum					X		
Plectranthus parviflorus	Cockspur Flower		0.1			X		
<b>Lauraceae</b>								
* Cinnamomum camphora	Camphor Laurel		0.1					
Cryptocarya glaucescens	Jackwood	1	1		1			
<b>Meliaceae</b>								
# Melia azedarach	White Cedar, Persian Lilac				0.1			
# Toona ciliata	Australian Red Cedar	0.1				X		
<b>Menispermaceae</b>								
Stephania japonica var. discolor	Snake Vine					X		
<b>Moraceae</b>								
Ficus coronata	Creek Sandpaper Fig	3			10	X		
Ficus rubiginosa	Port Jackson Fig, Rusty Fig	10					X	
<b>Myrtaceae</b>								
Acmena smithii	Lilly-pilly	2	2					
Angophora costata	Sydney Red Gum, Smooth-barked Appl					X		
Backhousia myrtifolia	Grey Myrtle, Ironwood	2						
Eucalyptus acmenoides	White Mahogany					X	X	
Eucalyptus saligna	Sydney Blue Gum	15	15		20			X
Melaleuca styphelioides	Prickly Paperbark		1	5		X		X
Syncarpia glomulifera	Turpentine		60		25	X	X	X
Syzygium australe	Brush Cherry	1		1				
# Syzygium paniculatum	Brush Cherry		1	1				
Tristaniopsis laurina	Water Gum, Kanuka Box					X		
<b>Ochnaceae</b>								
* Ochna serrulata	Mickey Mouse Plant					X		
<b>Oleaceae</b>								
* Ligustrum lucidum	Broad-leaved Privet, Glossy Privet					X		
Notelaea longifolia	Large Mock-olive					X		
<b>Oxalidaceae</b>								
* Oxalis latifolia		0.1		0.1		X		
<b>Passifloraceae</b>								
* Passiflora suberosa	Corky Passionflower	0.1						
<b>Pittosporaceae</b>								
Pittosporum revolutum	Yellow Pittosporum	1				X		
Pittosporum undulatum	Pittosporum		15		7	X		X
<b>Proteaceae</b>								
# Grevillea robusta	Silky Oak			10				
# Stenocarpus sinuatus	Firewheel Tree	0.1						
<b>Rosaceae</b>								
Rubus rosifolius	Native Raspberry, Rose-leaf Bramble	1	1					

Botanical name	Common name	Sub-quadrats				GT locs.		
		1-1	1-2	1-3	1-4	GTa	GTb	GTc
<b>Rutaceae</b>								
Melicope micrococca	Hairy-leaved Doughwood				0.1			
<b>Sapindaceae</b>								
Dodonaea triquetra	Hopbush	0.1						
<b>Solanaceae</b>								
* Solanum pseudocapsicum	Jerusalem Cherry		0.1	0.1				
<b>Sterculiaceae</b>								
# Brachychiton acerifolius	Illawarra Flame-tree, Flame Kurrajong	5	1			X		
<b>Ulmaceae</b>								
* Celtis occidentalis	American Hackberry		1					
Trema tomentosa var. aspera	Poison Peach, Peach-leaf Poison Bush		1					
<b>Verbenaceae</b>								
* Lantana camara	Lantana					X	X	
<b>Vitaceae</b>								
Cayratia clematidea	Slender Grape	0.1		0.1	0.1	X		
<b><u>4. Monocotyledons</u></b>								
<b>Arecaceae</b>								
Livistona australis	Cabbage Palm, Cabbage-tree Palm	2			1			
<b>Asteliaceae</b>								
# Cordyline rubra	Palm-lily					X		
<b>Commelinaceae</b>								
Commelina cyanea	Blue Spiderwort	1	1	1	1	X		
* Tradescantia fluminensis	Trad, Wandering Jew					X		
<b>Cyperaceae</b>								
Cyperus imbecillis				0.1				
Cyperus tetraphyllus		0.1	1					
<b>Iridaceae</b>								
* Dietes grandiflora	Butterfly Iris		1					
* Neomarica sp.	Walking Iris					X		
<b>Lomandraceae</b>								
Lomandra longifolia	Honey Reed, Spike Mat-rush	1	1	1				
<b>Philesiaceae</b>								
Eustrephus latifolius	Wombat Berry					X		
<b>Phormiaceae</b>								
Dianella caerulea	Blue Flax-lily			0.1		X		
<b>Poaceae</b>								
* Ehrharta erecta	Panic Veld-grass	0.1	0.1	0.1		X		
Entolasia marginata	Bordered Panic	0.1		0.1				
Microlaena stipoides	Weeping Grass, Meadow Rice-grass					X		
Oplismenus aemulus	Australian Basket Grass, Wavy Beard	1	1	20	1	X		

Table 5: Species recorded greater than 2 m in height in the supplementary sampling locations, including spot locations, ground truthing locations, and ground truthing quadrats, in the 4 re-surveyed parks (Brush Farm Park, Darvall Park, Field of Mars Reserve, Lambert Park) and 2 previously unsurveyed parks (Bell Park, Outlook Park)

Bell Park

Species	GTa		GTb	
	Number	Height (m)	Number	Height (m)
<i>Acacia parramattensis</i>	X			
<i>Acmena smithii</i>	X			
# <i>Angophora costata</i>	X			
<i>Angophora floribunda</i>	X	heights		heights
# <i>Araucaria bidwillii</i>			X	
# <i>Casuarina glauca</i>	X			
<i>Ceratopetalum apetalum</i>	X			
# <i>Eucalyptus grandis</i>			X	
# <i>Eucalyptus microcorys</i>	X		X	
<i>Eucalyptus paniculata</i>			X	
# <i>Eucalyptus pilularis</i>	X	not	X	not
# <i>Eucalyptus robusta</i>				
# <i>Eucalyptus saligna</i>			X	
<i>Eucalyptus scoparia</i>	X			
* <i>Jacaranda mimosifolia</i>				
* <i>Lantana camara</i>	X			
* <i>Ligustrum lucidum</i>	X	recorded		recorded
* <i>Ligustrum sinense</i>	X			
# <i>Melaleuca styphelioides</i>	X			
* <i>Phoenix canariensis</i>			X	
<i>Pittosporum undulatum</i>	X			
* <i>Quercus robur</i>			X	
* <i>Rothmannia globosa</i>			X	
<i>Syzygium paniculatum</i>	X			
# <i>Toona ciliata</i>	X			
<i>Tristaniopsis laurina</i>	X			



### Bell Park

Species	Spot A		Spot B	
	Number	Height (m)	Number	Height (m)
# <i>Brachychiton acerifolius</i>			1	3
# <i>Eucalyptus microcorys</i>	3	20	7	30
# <i>Eucalyptus robusta</i>			4	20
<i>Ficus rubiginosa</i>	1	17		
# <i>Melaleuca styphelioides</i>			1	7

### Brush Farm Park

Species	GTa		GTb		GTc	
	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Acacia parramattensis</i>			1	18		
<i>Acmena smithii</i>	X	not				
<i>Backhousia myrtifolia</i>			1	5		
# <i>Eucalyptus grandis</i>			1	27		
* <i>Ligustrum lucidum</i>	X					
<i>Melia azedarach</i>					1	15
<i>Pittosporum undulatum</i>	X	recorded	1	17	1	17
# <i>Toona ciliata</i>					1	7

Darvall Park

Species	GTa			GTb		
	Number	Height (m)	Diameter (m)	Number	Height (m)	Diameter (m)
# <i>Brachychiton acerifolius</i>	1	15				
<i>Eucalyptus acmenoides</i>				X	18	
# <i>Eucalyptus grandis</i>			0.4			
<i>Eucalyptus saligna</i>	1	25	1	1	25	0.45
# <i>Lophostemon confertus</i>	X	13				
# <i>Melaleuca styphelioides</i>	1	15				
<i>Pittosporum undulatum</i>				X	7	
<i>Syncarpia glomulifera</i>				1	14	0.3

Species	GTC			GTD		
	Number	Height (m)	Diameter (m)	Number	Height (m)	Diameter (m)
<i>Acmena smithii</i>	X	4		X	6	
# <i>Brachychiton acerifolius</i>				X	6	
<i>Callistemon salignus</i>				1	10	
<i>Elaeodendron australe</i>				X	2	
<i>Eucalyptus paniculata</i>	X	30	0.8			
<i>Eucalyptus saligna</i>	X	30		X	30	0.9
# <i>Melaleuca styphelioides</i>	X	8				
<i>Notelaea longifolia</i>	X	2				
<i>Syncarpia glomulifera</i>	X	7		X	17	
<i>Trema tomentosa</i>	X	2				

Darvall Park

Species	GTe			GTf		
	Number	Height (m)	Diameter (m)	Number	Height (m)	Diameter (m)
<i>Acacia parramattensis</i>				X	15	
# <i>Brachychiton acerifolius</i>	X	9				
<i>Eucalyptus acmenoides</i>	X	25	0.5			
# <i>Eucalyptus grandis</i>				X	20	0.2
<i>Eucalyptus saligna</i>	X	30	0.6			
<i>Eucalyptus tereticornis</i>				X	6	
# <i>Melaleuca styphelioides</i>				X	6	
<i>Notelaea longifolia</i>	X	2				
<i>Syncarpia glomulifera</i>	X	10				

Species	GTg		
	Number	Height (m)	Diameter (m)
# <i>Brachychiton acerifolius</i>	X	7	
# <i>Eucalyptus grandis</i>	X	20	0.6
<i>Eucalyptus tereticornis</i>	X	28	0.6
# <i>Lophostemon confertus</i>	X	9	
<i>Trema tomentosa</i>	X	25	0.7

Field of Mars Reserve

Species	GT QA-1		GT QA-2		GT QA-3		GT QA-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Acacia longifolia</i>			1	3				
<i>Allocasuarina littoralis</i>	5	12	2	12			2	14
<i>Angophora costata</i>	2	17	1	16	1	16	2	16
<i>Breynia oblongifolia</i>			3	4	2	2	3	4
# <i>Cyathea cooperi</i>					3	3	1	3
<i>Elaeocarpus reticulatus</i>	2	3						
<i>Eucalyptus resinifera</i>					3	18		
<i>Glochidion ferdinandi</i>			3	7				
<i>Leptospermum polygalifolium</i> subsp. <i>polygalifolium</i>	1	2						
* <i>Ligustrum lucidum</i>			1	2				
* <i>Ligustrum sinense</i>								
<i>Melaleuca linariifolia</i>					2	13		
<i>Notelaea longifolia</i>	3	3					1	5
<i>Pittosporum undulatum</i>	5	4	9	4	2	4	8	4
<i>Polyscias sambucifolia</i>	4	2						
* <i>Senna pendula</i> var. <i>glabrata</i>			1	3			3	4

Species	GT QB-1		GT QB-2		GT QB-3		GT QB-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Acacia fimbriata</i>			1	5				
<i>Allocasuarina littoralis</i>	40+	8	50+	8	25	8	20	6
<i>Angophora costata</i>	1	14					2	11
<i>Corymbia gummifera</i>	1	10	3	16	2	10		
<i>Eucalyptus globoidea</i>					1	16	1	15
<i>Eucalyptus resinifera</i>	1	2			3	13	1	7
<i>Grevillea sericea</i>								
<i>Leptospermum trinervium</i>					1	3		
<i>Notelaea longifolia</i>					1	2	1	2
<i>Persoonia lanceolata</i>	1	3	1	3				

Field of Mars Reserve

Species	GT QC-1		GT QC-2		GT QC-3		GT QC-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Allocasuarina littoralis</i>	2	4	3	5	3	5	2	6
<i>Angophora costata</i>	4	17	4	18	3	18	2	13
<i>Dodonaea triquetra</i>					2	3	2	3
<i>Elaeocarpus reticulatus</i>			1	5				
<i>Persoonia levis</i>			1	2			1	3
<i>Xylomelum pyriforme</i>					1	2		

Species	GT QD-1		GT QD-2		GT QD-3		GT QD-4	
	Number	Height (m)	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Allocasuarina littoralis</i>	7	10	5	11			1	5
<i>Allocasuarina torulosa</i>			1	9			1	12
<i>Angophora costata</i>	8	14	6	18	6	18		
<i>Breynia oblongifolia</i>							1	2
<i>Dodonaea triquetra</i>	1	2						
<i>Eucalyptus globoidea</i>	1	16						
<i>Eucalyptus resinifera</i>			3	10	1	18		
<i>Notelaea longifolia</i>					1	3		
<i>Pittosporum undulatum</i>	2	2	4	2	5	3	4	4
* <i>Senna pendula</i> var. <i>glabrata</i>					1	3		
<i>Syncarpia glomulifera</i>	2	12			1	10	1	13

Field of Mars Reserve

Species	GTa		GTb		GTc	
	Number	Height (m)	Number	Height (m)	Number	Height (m)
	3 (ArborPlan 317, 314, 312)	heights				
<i>Angophora costata</i>			X	18		
<i>Eucalyptus pilularis</i>			X	15		
<i>Eucalyptus punctata</i>	1	not				
	(ArborPlan 313)				X	20
<i>Eucalyptus resinifera</i>					X	3
<i>Glochidion ferdinandi</i>		recorded			X	6
<i>Hakea salicifolia</i>					X	7
<i>Pittosporum undulatum</i>			X	8	X	4
<i>Polyscias sambucifolia</i>					X	4

Lambert Park

Species	GTa			GTb			GTc		
	Number	Height (m)	Diameter (m)	Number	Height (m)	Diameter (m)	Number	Height (m)	Diameter (m)
<i>Acacia parramattensis</i>				(many)	15				
<i>Acmena smithii</i>				X	7				
<i>Eucalyptus globoidea</i>							(several)	10	
# <i>Eucalyptus grandis</i>				1	25	0.3			
<i>Eucalyptus paniculata</i>							X	12	
<i>Eucalyptus pilularis</i>	1	11	0.25						
<i>Eucalyptus saligna</i>	1	25+	<0.4						
# <i>Melaleuca styphelioides</i>							X	10	

Species	GTd			GTe			GTf		
	Number	Height (m)	Diameter (m)	Number	Height (m)	Diameter (m)	Number	Height (m)	Diameter (m)
* <i>Celtis sinensis</i>				1	8				
<i>Elaeodendron australe</i>				X	5				
<i>Eucalyptus globoidea</i>	1	2							
<i>Hibiscus heterophyllus</i>	1	20							
<i>Homalanthus populifolius</i>				X	5				
# <i>Melaleuca styphelioides</i>				X	3				
<i>Melaleuca quinquenervia</i>	(numerous)	10		X	9				
<i>Pittosporum undulatum</i>				X	3				
<i>Syncarpia glomulifera</i>							X	9	
<i>Ulmus parvifolia</i>	2	not recorded							

### Outlook Park

Species	GTa		GTb		GTc	
	Number	Height (m)	Number	Height (m)	Number	Height (m)
<i>Angophora costata</i>	2	27				
<i>Eucalyptus acmenoides</i>	2	25	1	35		
<i>Eucalyptus saligna</i>					3	30
<i>Ficus rubiginosa</i>			1	10		
<i>Syncarpia glomulifera</i>			10	25	5	25